

Phenolic Compounds of *Rumex* L: Aerial Part Fractions and Essential Oil Results of *In vitro* Screening for Antimicrobial Activity

**G. D. Shermatova^{a*}, Kh. M. Bobakulov^a, B. A. Shamuratov^b,
S. M. Mavlyanov^b, Y. J. Zhang^c, K. A. Eshbakova^a, Sh. S. Azimova^a
and S. A. Sasmakov^a**

^a Institute of the Chemistry of Plant Substances named after Acad. S.Yu. Yunusov Academy of Sciences of the Republic of Uzbekistan, 100170, 77, M. Ulugbek str., Tashkent, Uzbekistan.

^b Institute of Bioorganic Chemistry named after Acad. O.S. Sadykov Academy of Sciences of the Republic of Uzbekistan, 100125, 83, M. Ulugbek str., Tashkent, Uzbekistan.

^c Kunming Institute of Botany, Chinese Academy of Sciences, 650201, 132, Lanhei Road, Heilongtan, Kunming, China.

Authors' contributions

This work was carried out in collaboration among all authors. Author GDS did the conceptualization. Authors BAS, SMM, YJZ, XMB and SAS methodology. Author GDS validation. Authors GDS and XMB formal analysis. Authors GDS, XMB and SAS investigation. Author GDS writing-original draft preparation. Author GDS writing-review and editing. Authors BAS, SMM, YJZ and KAE supervision. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CSJI/2022/v31i1130273

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86738>

Original Research Article

Received 12 March 2022

Accepted 26 May 2022

Published 31 May 2022

ABSTRACT

The aim of the present study to clarify the phytochemical constituents of *Rumex* plants, extraction, separation and structural identification of chemical components, provide theoretical support and scientific basis for its traditional use, further expand the application scope of traditional medicinal plant resources in Uzbekistan, and develop its potential value. Three known anthraquinone derivatives has been isolated from the chloroform fraction extract of *Rumex pamiricus* roots and two known compounds were separated from the ethyl acetate fraction extract of *Rumex conglomeratus* roots. Their structures were elucidated by extensive spectroscopic evidence and chemical methods.

*Corresponding author: E-mail: gdshermatova@gmail.com;

Qualitative analyses of major phenolics by TLC analysis were also evaluated. The ester oil from the aerial parts of *Rumex confertus* and *Rumex pamiricus*, the gasoline, chloroform, ethyl acetate and butanol extracts of *Rumex pamiricus* aerial part and alcohol extract of *Rumex pamiricus* roots were tested *in vitro* for antimicrobial activity.

This study aimed to determine the *in vitro* antimicrobial activity of the medicinal plants traditionally used in Uzbekistan against the microbial strains associated with infectious diseases.

Keywords: *Polygonaceae* L.; *Rumex pamiricus* Rech. f.; *Rumex conglomeratus* Murray.; *Rumex confertus* Willd.; sorrel; dock; phenol; phenolic acid; flavonoid; anthraquinone; *In vitro*; antimicrobial; extract; fraction.

1. INTRODUCTION

In the field of medicinal chemistry natural products research, Uzbekistan is a latter-coming and less active player even though Uzbekistan has a very long history to use local medicinal plants to treat various diseases. In recent years, Uzbekistan government has launched several projects, to support scientists carrying out natural products research. As a result, in the course of our continued searching for bioactive compounds from Uzbekistan and in the territory of Central Asia plants we did find that Uzbekistan possesses a great biodiversity and a great chemical diversity as well.

1.1 Importance of Medicinal Plants

“Herbal remedies play an important role in modern medicine and it appears feasible that the compounds from herbs can be helpful in prevention or treatment of different diseases” [1]. “The interest of natural drugs as adjunctive therapy for acute and chronic diseases has grown significantly in the recent years” [2]. The phenolic compounds are of great importance in terms of various biological activities in the research work in this area. Phenolic compounds are probably the most explored natural compounds due to their potential health benefits as demonstrated in a number of studies. Continuing these studies, we began to study the phenols of the plants *Rumex pamiricus*, *Rumex confertus* and *Rumex conglomeratus* in order to isolate natural compounds from local plant raw materials and study their biological activity (Fig. 1). The priority is this research to study the discovery of biological active substances constituents basis of genus *Rumex* plants in Uzbekistan and study chemical constituents of phenolic compounds.

1.2 Role of Phenolic Compounds

“Phenolic compounds are secondary metabolites that are derivatives of the pentose phosphate,

shikimate and phenylpropanoid pathways in plants. Major polyphenolic compounds found in plants are flavonoids, catechins, epigallocatechin-3-gallate (EGCG), flavonones, iso-flavones, flavanols, anthocyanins, phenolic acids, stilbenes, flavonoids, chalcones, lignans etc. These compounds are secondary plant metabolites and possess antimicrobial, antiviral and anti-inflammatory properties along with high antioxidative activity. These properties make polyphenols interesting for the treatment of various diseases like inflammation, cancer and also used for anti-ageing purposes in cosmetic formulations as well as have nutraceutical applications”. [3]. Phenolic compounds are found in different organs and tissues of plants differ from each other not only in quantity but also in quality. Because their participation in various biochemical processes is determined by the structure and degree of polymerization. Most simple phenolic compounds are easily oxidized and are actively involved in metabolism. This is why they are concentrated in the tissues of leaves, flowers, and growth points, where most biochemical processes are most active [4].

1.3 The genus *Rumex* and Distribution

“The name *Rumex* derived from the Latin word for dart, alluding to the shape of the leaves. It is the largest genus of family *Polygonaceae*” [1]. “Plants of the genus *Rumex* L. (sorrel, dock) are widely distributed in North America, Central and Eastern Europe, Kazakhstan, the Far East and partly in the Caucasia, Russia and East Asia” [5-7]. “This genus includes more than 250 species distributed worldwide. 16 species grow in Uzbekistan. *Rumex pamiricus*, *Rumex Confertus* and *Rumex conglomeratus* are the most common species among them” [2,8,9]. “Since ancient times, concoctions and tea from leaves and roots of *Rumex* L. species have been used to treat various intestinal inflammations” [2,10]. “The herb *Rumex pamiricus* belongs to the family of *Polygonaceae* which is widespread in Central

Asia (Pamir-Alay, Tian Shan, Dzungarian Alatau), Kashgaria. One of the most common type of *Rumex* is present in Uzbekistan (Samarkand and Kashkadarya regions). It grows along wet mountain meadows, along the banks of mountain rivers and lakes. Perennial herbaceous plant reaching 60–100 cm in height” [11]. “Since ancient times, concoction or tea from various parts of this herb has been used in folk medicine to treat diarrhea, dysentery, stercoral ulcer, as appetizer, analeptic medicine for liver, heart, as antihemorrhagic, to treat hepatitis, fever and other diseases” [2]. “The consumption of wild edible plants has been an integral part of human nutrition and traditional medicine since ancient times” [12,13]. “Thus, researchers began to pay more attention to wild flora. Plants of the genus *Rumex* are no exception. In addition, this plant genus is known as a super-producer of secondary phenolic compounds” [14]. “Wild plants are known to be a good source of primary nutritional compounds (proteins, fats, sugars, vitamins, and minerals)” [15]. “Wild plants contain various biologically active components that demonstrated health benefits effects (flavonoids, phenolic acids, anthocyanins, tannins, terpenoids, steroidal saponins, glucosinolates, and so on)” [13].

1.4 Source of Secondary Metabolites

“Higher plants synthesize several thousand known different phenolic compounds and the number of those fully characterized is continually increasing” [16]. “Phenolic compounds are known to have strong antioxidant as well as cardioprotective, immune system promoting, antibacterial, anti-cancer, and anti-inflammatory effects” [14]. “Plants of the *Rumex* genus are rich in secondary metabolites, in particular flavonoids, phenolic acids and anthraquinones, which are likely to be responsible for the medicinal properties attributed to these species” [17]. “The list of **anthraquinones** particularly common in *Rumex* plants includes chrysophanol, physcion, emodin and their glycosides, rhein, nepodin, and so on” [14]. “These compounds also show anticarcinogenic, anti-inflammatory, antiarthritic, antifungal, antibacterial, antioxidant and diuretic activity” [18,19]. “**Flavonoids** are another important class of compounds that determine the therapeutic effect of *Rumex* plants. Derivatives of kaempferol, quercetin, apigenin, luteolin, and catechins, as well as derivatives of benzoic and cinnamic acids, lignans, coumarins, and proanthocyanidins, have been isolated from

various *Rumex* species” [20]. “Flavonoids constitute the largest class of phenolic compounds with more than 3,000 structures. These consist of two aromatic rings bound together by three carbon atoms that form an oxygenated heterocycle. For many years there has been much interest in flavonoids because of their beneficial effects on human health: e.g. antioxidant activity, free radical scavenging capacity, digestive stimulation action, anti-inflammatory, antimicrobial, antiviral, hypolipidemic, antimutagenic effects and anticarcinogenic potential” [21]. “**Phenolic acids** are further divided into hydroxycinnamic acid (HCA) and hydroxybenzoic acid (HBA). Phenolic acids account for about a third of the polyphenolic compounds in our diet and are found in all plant materials, but are particularly abundant in acidic fruits. Caffeic acid, gallic acid, ferulic acid are some common phenolic acids” [22]. “Fruits, vegetables, beverages such as tea, chocolate and wine are the chief sources of phenolic compounds in the human diet” [3, 23]. “This shows their potential as nutritional supplements, feed additives, and medicinal agents” [13, 24].

1.5 Nutritional Aspects and Traditional uses of *Rumex* L.

“Among wild plants, *Rumex* plants have a great potential. They are already widely used as food, fodder, melliferous, and medicinal plants” [7, 25, 26]. “In some countries, the leaves of *Rumex* plants (such as *R. vesicarius*, *R. acetosella*, *R. abyssinicus*, *R. crispus*, *R. induratus*, *R. sanguineus*, *R. obtusifolius*, *R. tuberosus*, *R. thyrsiflorus*, and *R. acetosa*) are used for food, mainly as salads” [18,19]. “According to the literature information, several *Rumex* species are included in the pharmacopoeias of various countries. For example, *R. crispus* is listed in the American Herbal Pharmacopoeia as a general detoxifier and an agent for skin treatment” [27]. “The State Pharmacopoeia of the Russian Federation includes the roots of *R. confertus* as a herbal medicine, which is used in the treatment of liver diseases, dysentery, pulmonary and uterine bleeding, as well as a laxative” [28, 29]. “*Rumex* plants have traditionally been used as edible or medicinal plants in various regions of the world” [14,29]. “Several *Rumex* species have been used in traditional Chinese medicine (TCM) for the therapy of different diseases. *R. dentatus*, found almost everywhere in China, has been employed traditionally for the treatment of many



Fig. 1. *R. pamiricus* Rech. f. (1); *R. confertus* Willd. (2); *R. conglomeratus* Murray (3). Location: Beldersay, Chimgan mountains (Ugam Chatkal National Park), Tashkent region. (Pictures author: G.D.Shermatova)

kinds of bacterial and fungal infections, e.g. dysentery, enteritis and acariasis” [30]. “*R. crispus* has a long history of domestic herbal use in India and Pakistan. It is a gentle and safe laxative and useful for treating a wide range of skin problems (sores, ulcers and wounds). The root of the plant is alterative, mildly tonic, antiscorbutic, cholagogue and astringent, while the seeds effective in the treatment of diarrhea” [31,32]. In Australia *Rumex* species are used for the treatment of stings [33]. The extracts of some *Rumex* species (*R. hymenosepalus* and *R. maderensis*) are used as a “blood depurative” or “blood purifier” [34,35]. “Literature demonstrates that *R. hastatus* is traditionally used in the treatment of sexually transmitted diseases, including AIDS” [36]. “*R. nepalensis* is applied to treat stomachache in Ethiopian regions. *R. vesicarius* is a wild edible Egyptian herb. In folk medicine, it is used as a tonic and analgesic and for the treatment of hepatic diseases, constipation, poor digestion, spleen disorders, flatulence, asthma, bronchitis, dyspepsia, vomiting and piles, among others” [37]. “The roots of *R. confertus* are used for liver diseases,

dysentery, pulmonary and uterine bleeding, as a laxative, for hemorrhoids and anal fissures, externally for burns, wounds, stomatitis, gingivitis, and skin diseases in Russian Federation” [7]. “However, today, their biotechnological potential is becoming evident, and these species can act as a source of biologically active substances. The *Rumex* plants are abundant, undemanding, gain phytomass easily, and have a short vegetative cycle (and, as a consequence, can reproduce frequently throughout the year), thus they have a real advantage among wild plants of the temperate zone. It should also be noted that *Rumex* species have a high potential for regrowth after injury” [14].

2. EXPERIMENTAL PART

2.1 Plant Material

The aerial parts (during the flowering period on May 2020) and roots (on August 2020) of the plants were collected from Botanic Garden, Tashkent, Uzbekistan.



Fig. 2. Some pictures of the collected samples. The leaves of *Rumex pamiricus* Rech. f. (1); the roots of *Rumex conglomeratus* (2); the aerial part of *Rumex confertus* Willd. (3); The process of collecting *R. Pamiricus* Rech. f. (4); the seeds of *Rumex confertus* Willd. (5). Location: Tashkent Botanical Garden named after F. N. Rusanov. (Pictures author: G.D.Shermatova)

2.2 Extraction and Methods

The roots of the herb *Rumex pamiricus* dried at room temperature, in shade. The pounded herb roots were first subjected to extraction in chloroform, then three times in 70% acetone hydrous solution. The acetone extract was distilled under vacuum, the remaining water solution was subjected to extraction with ethyl acetate. Ethyl acetate extracts were collected and were dehydrated by adding anhydrous salt Na_2SO_4 . The dehydrated extract was filtered, its concentration increased under vacuum, the total phenols were precipitated by adding pure hexane to the condensed extract. The created precipitate was washed, and filtered and the extracted total phenols of chloroform and ethyl acetate fractions constituted 3.4% of the herb dry weight.

2.3 Isolation and Results

The roots are the best organs for the accumulation of anthraquinones [2]. The chloroform fraction subjected with column chromatography on KSK silica gel, eluted with a mixture of extraction benzene–ethyl acetate: (50:1, 40:1, 30:1, 20:1 and 10:1). The structure of chrysophanol, emodin and rhein (Fig. 3) was established on the basis of the analysis of the data of MS (Mass spectrometry), ^1H and ^{13}C NMR spectra (Nuclear Magnetic Resonance), and of the DEPT (Distortionless Enhancement of Polarization Transfer), HSQC (Heteronuclear Single Quantum Coherence) and HMBC (Heteronuclear Multiple Bond Correlation) experiments. Qualitative analyses of major phenolics by TLC (Thin Layer Chromatography) analysis were also evaluated.

2.4 Discussion

The anthraquinones, chrysophanol, emodin and rhein have been isolated from other types of *Rumex* in studies before us. According to the

literature: chrysophanol from roots of *R. dentatus* [30], emodin from roots of *R. abssinicus* [38], from leaves of *R. chalepensis* [39], rhein from *R. acetosa* [38], chrysophanol and emodin from roots of *R. crispus* [38], *R. abyssinica*, *R. patientia* [39], *R. nepalensis* [40], and from the aerial parts of *R. acetosa* [41].

2.5 Extraction and Methods

The *Rumex conglomeratus* (20 kg) were extracted with 60% acetone four times at room temperature, each one time one week. After filtration and condensed to little volume by Rotary evaporation machine, the concentrated liquid (most are water) was fractionated by EtOAc (1/4 volume of water) five times, to give two parts, water layer and EtOAc layer. After filtration, the EtOAc layer was dried by Rotary evaporation machine to give 72.4 g of the EtOAc sample.

2.6 Isolation and Results

The EtOAc fraction was then subjected to Diaion HP 20SS column chromatography with MeOH containing increasing proportions of water (6 cm i.d.- 90 cm, 10-100%, 10% stepwise elution, each 3L), to afford seven fractions E1-E7. Fraction E-1 was further fractionated by Sephadex LH-20 column chromatography (7 cm i.d.-110 cm) with 10-100% MeOH(10% stepwise elution, each 3-L) and give to six fractions E1-1-E1-6. And the fraction E1-2 were separated by column chromatography using the MCI gel CHP 20P (3 cm i.d.- 40 cm) with 10-100% MeOH (5% stepwise elution, each 300 ml) to yield caffeic acid (Fig. 3) (0.092 g). Fraction E1-5 was successively applied to a Toyopearl HW 40F column chromatography (2 cm i.d.- 28 cm) with 5-100% MeOH (5 stepwise elution, each 150 ml) to give rosmarinic acid (Fig. 3) (0.252 g).




Fig. 3. Chemical structures of isolated compounds 1–5: rosmarinic acid (1), caffeic acid (2), chrysophanol (3), emodin (4), rhein (5). All known pure compounds were isolated in our laboratory from *Rumex pamiricus* and *Rumex conglomeratus* plants for the first time

2.7 Discussion

Rosmarinic acid and caffeic acid have been isolated from other species and families in previous studies and their properties have been studied. Rosmarinic acid was first isolated in 1958 by Scarpati and Oriente from *R. officinalis* [42], from *Melissa officinalis* L.[43], from *Hyptis atrorubens* Poit [44,45], [46], [47,48,49,50], caffeic acid from the aerial parts of *R. aquatica* [39], rosmarinic acid and caffeic acid from leaves of *R. acetosa*, *R. acetosella*, *R. confertus*, *R. crispus*, *R. maritimus*, *R. obtusifolius* and *R. sanguineus* [14].

2.8 Data of Isolated Compounds

1. **Rosmarinic acid (1)** [44]. C₁₈H₁₆O₈. Yellow amorphous powder. ESI-MS (negative ion) *m/z* 359 [M-H]⁻; UV λ_{max} (CH₃OH): 221, 330 nm; IR (KBr) (cm⁻¹) ν_{max}: 3382, 1697, 1606, 1522; ¹H NMR (CD₃OD): δ 7.52 (1H, d, *J* = 15.5 Hz, H-7'), 7.04 (1H, d, *J* = 2.0 Hz, H-2'), 6.91 (1H, dd, *J* = 8.0, 2.0 Hz, H-6'), 6.75 (1H, d, *J* = 8.0 Hz, H-5'), 6.70 (1H, d, *J* = 2.0 Hz, H-2), 6.69 (1H, d, *J* = 8.0 Hz, H-5), 6.57 (1H, dd, *J* = 8.0, 2.0 Hz, H-6), 6.26 (1H, d, *J* = 15.5 Hz, H-8'), 5.19 (1H, dd, *J* = 10.0, 3.5 Hz, H-8), 3.06 (1H, dd, *J* = 14.5, 5.5 Hz, H-7a), 3.00 (1H, dd, *J* = 14.5, 5.5 Hz, H-7b); ¹³C-NMR (CD₃OD): δ 174.3 (C-9), 169.4 (C-9'), 149.5 (C-4'), 145.7 (C-7'), 145.0 (C-3), 144.9 (C-4), 143.2 (C-3'), 130.4 (C-1), 128.2 (C-1'), 122.2 (C-6), 121.4 (C-6'), 117.6 (C-2), 116.3 (C-5), 116.1 (C-5'), 115.8 (C-8'), 115.3 (C-2'), 77.8 (C-8), 38.9 (C-7).
2. **Caffeic acid (2)** [51], colorless acicular crystals, mp 222.1-225.8°C. HR- ESI-MS *m/z* 179.0378 [M-H]⁻ (calcd for C₉H₇O₄, 179.0344). ¹H NMR (400 MHz, DMSO-d₆, δ, ppm, J/Hz): 7.39 (1H, d, *J*=16.0, H-7), 7.02 (1H, d, *J*=2.0, H-2), 6.94 (1H, dd, *J*=8.0, 2.0, H-6), 6.75 (1H, d,*J*=8.0, H-5), 6.17 (1H, d,*J*=16.0, H-8). ¹³C NMR (125 MHz, DMSO-d₆, δ, ppm): 125.8 (C-1), 115.6 (C-2), 144.3 (C-3), 148.1 (C-4), 115.8 (C-5), 121.1 (C-6), 145.6 (C-7), 114.6 (C-8), 168.2 (C-9).
3. **Chrysophanol (3)** [52]. ¹H NMR (600 MHz, CDCl₃, δ, ppm, J/Hz): 12.14 (1H, s), 12.03 (1H, s), 7.83 (1H, d, *J* = 6.8, H-5), 7.67 (2H, s, H-6, 4), 7.31 (1H, d, *J* = 6.8, H-7), 7.12 (1H, s, H-2), 2.48 (3H, s, 3-Me). ESI-MS *m/z* 255.0 [M + H]⁺.

4. **Emodin (4)** [53]. Orange needles, C₁₅H₁₀O₅, mp 254–256°C. ¹H NMR (600 MHz, acetone-d₆, δ, ppm, J/Hz): 7.54 (1H, s, H-4), 7.23 (1H, d, *J* = 2.3, H-5), 7.12 (1H, s, H-2), 6.63 (1H, d, *J* = 2.3, H-7), 2.44 (3H, s, CH₃). ¹³C NMR (125 MHz, CDCl₃, δ, ppm): 190.9 (C-9), 181.4 (C-10), 165.5 (C-3), 165.2 (C-1), 162.4 (C-8), 148.8 (C-6), 135.8 (C-14), 135.8 (C-11), 124.2 (C-5), 124.1 (C-7), 120.7 (C-13), 113.7 (C-12), 108.8 (C-4), 108.0 (C-2), 21.2 (CH₃).
5. **Rhein (5)** [54]. C₁₅H₈O₆, yellowish powder. ESI-MS *m/z* 307.2 [M + Na]⁺. ¹H NMR (600 MHz, DMSO-d₆, δ, ppm, J/Hz): 8.15 (1H, d, *J* = 1.7, H-4), 7.78 (1H, d, *J* = 1.7, H-2), 7.75 (1H, dd, *J* = 7.8, 1.0, H-5), 7.85 (1H, t, *J* = 8.0, 7.8, H-6), 7.42 (1H, dd, *J* = 8.0, 1.0, H-7). ¹³C NMR (100 MHz, DMSO-d₆, δ, ppm): 161.6 (C-1), 124.3 (C-2), 138.2 (C-3), 118.9 (C-4), 119.6 (C-5), 137.7 (C-6), 124.7 (C-7), 161.2 (C-8), 191.5 (C-9), 181.2 (C-10), 133.4 (4a), 116.4 (8a), 118.4 (9a), 134.1 (10a), 165.6 (COOH).

3. ANTIMICROBIAL ACTIVITY

Medicinal plants are widely used for the treatment of different infectious diseases. Infectious diseases caused by bacteria have a large impact on public health [55]. We studied the leaves and roots of *Rumex confertus* Willd. *in vitro* for antibacterial and fungal activity in the fractions of gasoline, chloroform, ethyl acetate and butanol in our previous work. As a result, it was found that the leaves of the *Rumex confertus* plant, chloroform and ethyl acetate fractions of the root part have antibacterial activity against fungi and positive bacteria [1]. Continuing these studies, the ester oil from the aerial parts of *Rumex confertus* and *Rumex pamiricus*, the gasoline, chloroform, ethyl acetate and butanol extracts of *Rumex pamiricus* aerial part and alcohol extract of *Rumex pamiricus* roots were tested *in vitro* for antimicrobial activity (Table 1). It was studied at the Institute of the Chemistry of Plant Substances named after Acad. S.Yu. Yunusov AS of Uzbekistan, Laboratory of Molecular Genetics by Dr. Sasmakov S.A.

4. RESULTS

As a result, the aerial part of the *Rumex pamiricus* plant, chloroform and ethyl acetate fractions showed moderate antimicrobial activity against gram-positive bacteria.

Table 1. *In vitro* screening results for antimicrobial activity

Name of substances	Inhibition zone diameter (mm)				
	Gram-positive strains		Gram-negative strains		Conditionally pathogenic fungus <i>Candida albicans</i>
	<i>Bacillus subtilis</i>	<i>Staphylococcus aureus</i>	<i>E. coli</i>	<i>Pseudomonas aeruginosa</i>	
Ester oil of <i>Rumex pamiricus</i> Rech. f. aerial part	6	7	na	na	na
Ester oil of <i>Rumex confertus</i> Willd. aerial part	6	6	na	na	na
Alcohol extract of <i>Rumex pamiricus</i> Rech. f. aerial part	10	8	na	na	na
Gasoline extract of <i>Rumex pamiricus</i> Rech. f. aerial part	6	6	na	na	na
Ethyl acetate extract of <i>Rumex pamiricus</i> Rech. f. aerial part	10	12	na	na	na
Alcohol extract of <i>Rumex pamiricus</i> Rech. f. root	na	na	na	na	na
Butanol extract of <i>Rumex pamiricus</i> Rech. f. aerial part	9	9	na	na	na
Chloroform extract of <i>Rumex pamiricus</i> Rech. f. aerial part	9	11	na	na	na
Ampicillin (10 µg/disc)	27	26	nt	nt	nt
Ceftriaxone (30 µg/disc)	nt	nt	26	25	nt
Flucanazole (25 µg/disc)	nt	nt	nt	nt	28

na- not active; nt – not tested

Inhibition zones ≤ 6-8 mm;

Appreciable: 8-14 mm;

Pronounced: 14-20 mm;

Strong: ≤ 20 mm

5. DISCUSSION

Nowadays, the role of secondary metabolites as regulatory and adaptogenic is not questioned. For instance, the wide geographical distribution of the *Rumex* plants can be partly associated with the flexible system of secondary metabolism. In this study, wild plants with relatively uniform growing conditions were used. The collection sites were located in similar climatic and landscape conditions, with a low anthropogenic load. In addition, the plants were analyzed within the same ontogenetic phase- the flowering phase. This point is fundamental, as the level of regulatory secondary compounds can differ significantly at different stages of growth [14].

During the study process, it was observed that the difference in the color of *R. pamiricus*, a plant that grows in mountainous areas, and *R. pamiricus*, which grows in urban areas. That is to say, it was found that the redness characteristic of the genus *R. pamiricus* is not clearly visible in plants growing in the mountainous region (Fig. 1), but is dark red in the city (Fig. 2).

5. CONCLUSION

1. Continuous studies on the chemical composition of *Rumex pamiricus* Rech. f. led to the isolation of anthraquinones: chrysophanol, emodin and rhein from the plant root extract using column chromatography on KSK silica gel. The structure of chrysophanol, emodin and rhein was established on the basis of the analysis of the data of MS, ¹H, and ¹³C NMR spectra, and of the DEPT, TLC, HSQC and HMBC experiments.
3. The 60% acetone extracts of *Rumex conglomeratus* Murray was successively separated by MCI gel CHP 20P, and Toyopearl HW 40F column chromatography to yield two compounds. Their structures were elucidated by spectroscopic analyses as: rosmarinic acid and caffeic acid.
4. The ester oil from the aerial parts of *Rumex confertus* and *Rumex pamiricus*, the gasoline, chloroform, ethyl acetate and butanol extracts of *Rumex pamiricus* aerial part and alcohol extract of *Rumex pamiricus* root were tested for antimicrobial activity. As a result, the aerial part of the *Rumex pamiricus* plant, chloroform and ethyl acetate fractions showed moderate

antimicrobial activity against gram-positive bacteria.

ACKNOWLEDGEMENTS

I would like to express my feelings of gratitude for the kind support and guide of my supervisors Dr. Shamuratov B.A, Prof. Mavlyanov S.M., Prof. Zhang Y.J. Also, I thank Dr. Bobakulov X.M for his assistance in determining the structures of chemical compounds and deeply thank the anonymous reviewers and the editorial committee who contributed to substantially improve the manuscript.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shermatova GD, Zhang YJ, Davranov K. Antibacterial and antifungal activities of rumex confertus willd. International Journal for Research in Applied Science & Engineering Technology. 2021;9/12:1855-1856.
2. Shermatova G. Emodin, an anthraquinone derivative from *Rumex pamiricus* Rech. f. Universum: chemistry and biology. 2022;3/93:28-31.
3. Kumar H, Choudhary N, Varsha, Kumar N, Suman, Seth R. Phenolic compounds and their health benefits: A review. Journal of Food Research and Technology. 2021;2/2:46-59.
4. Shermatova GD, Shamuratov BA, Mavlyanov SM. Tannins from rumex confertus willd. Uzbek Chemistry Journal. 2012;3:11-13.
5. Kołodziejek J. Growth performance and emergence of invasive alien *Rumex*

- confertus in different soil types. Scientific Reports. 2019;9/1:1-13.
6. Kholmatov HKh, Habibov ZN, Olimkhodjaeva NZ. In book: Medicinal herbs of Uzbekistan. Ibn Sino. 1991;93-94.
 7. Podgurskaya VV, Luksha EA, Gushchina ES, Savchenko IA, Korneeva IN, Kalinkina GI. Biological activity of the genus Rumex (Polygonaceae) plants. Chem. Plant Raw Mater. 2021;2:59–78.
 8. Shermatova GD, Shamuratov BA. Flavonoids of Rumex pamiricus. Bulletin of National University of Uzbekistan. 2013; 4/2:232-233.
 9. Rao KNV, Ch S, Banji D. A study on the nutraceuticals from the genus Rumex. Hygeia. J.D.Med. 2011;3/1:76-88.
 10. Babulka P. The Rumex, from ethnobotany to modern phytotherapy (Rumex spp.), [Les Rumex, de l'ethnobotanique á la phytothérapie moderne (Rumex spp.)], Phytothérapie. 2004;5:153-156.
 11. Vvedensky AI, Korovin EP. In book: Flora of Uzbekistan. Publishing House of the Academy of Sciences of the UzSSR. Tashkent. 1953;2:102-113.
 12. Carvalho AM, Barata AM. The consumption of wild edible plants. In book: Wild Plants, Mushrooms and Nuts: Functional Food Properties and Applications; Edited by Ferreira I.C.F.R, Morales P, Barros L. John Wiley & Sons, Ltd. 2017;159–198.
 13. Ceccanti C, Landi M, Benvenuti S, Pardossi A, Guidi L. Mediterranean wild edible plants: Weeds or “New Functional Crops”? Molecules. 2018;23:2299.
 14. Feduraev P, Skrypnik L, Nebreeva S, Dzhobadze G, Vatagina A, Kalinina E, Pungin A, Maslennikov P, Riabova A, Krol O, Chupakhina G. Variability of phenolic compound accumulation and antioxidant activity in wild plants of some rumex species (Polygonaceae). Antioxidants. 2022;11/2:311.
 15. Puccinelli M, Pezzarossa B, Pintimalli L, Malorgio F. Selenium biofortification of three wild species, *Rumex acetosa* L., *Plantago coronopus* L., and *Portulaca oleracea* L., Grown as Microgreens. Agronomy. 2021;11:1155.
 16. Lattanzio V. Phenolic compounds: Introduction. Springer-Verlag Berlin Heidelberg. 2013;50:1543-1580.
 17. Prakash MA, Sharifi RM, Shariati MA, Mabkhot YN, Al-Showiman SS, Rauf A, Salehi B, Zupunski M, Gusain P. et al. Bioactive compounds and health benefits of edible Rumex species. A review. Cell. Mol. Biol. 2018;64:27–34.
 18. Li Y, Jiang JG. Health functions and structure-activity relationships of natural anthraquinones from plants. Food Funct. 2018;9:6063–6080.
 19. Diaz-Munoz G, Miranda IL, Sartori SK, de Rezende DC, Diaz MAN. Chapter 11- Anthraquinones: An overview. In Studies in Natural Products Chemistry; Atta-ur-Rahman. Elsevier. 2018;58:313–338.
 20. Balog K, Svircev E, Lesjak M, Orcic D, Beara I, Franciskovic M, Simin N. Phenolic profiling of Rumex L. species by means of the LC-MS/MS. Planta Med. 2012; 78/11:12.
 21. Samsonowicz M, Regulska E. Spectroscopic study of molecular structure, antioxidant activity and biological effects of metal hydroxyflavonol complexes. Spectrochim. Acta Part A: Mol. Biomol.Spectrosc. 2017;173:757–771.
 22. Spencer JP, Mohsen MM, Minihane AM, Mathers JC. Biomarkers of the intake of dietary polyphenols: strengths, limitations and application in nutrition research. British Journal of Nutrition. 2008;99/1:12-22.
 23. Tsao R. Chemistry and biochemistry of dietary polyphenols. Nutrients. 2010; 2/12:1231-1246.
 24. Faehnrich B, Franz C, Nemaz P, Kaul HP. Medicinal plants and their secondary metabolites - State of the art and trends in breeding, analytics and use in feed supplementation with special focus on German chamomile. J. Appl. Bot. Food Qual. 2021;94:61–74.
 25. Bello OM, Fasinu PS, Bello OE, Ogbesejana AB, Adetunji CO, Dada AO, Ibitoye OS, Aloko S, Oguntoye OS. Wild vegetable Rumex acetosa Linn. Its ethnobotany, pharmacology and phytochemistry. A review. S. Afr. J. Bot. 2019;125:149–160.
 26. Vasas A, Orbán-Gyapai O, Hohmann J. The genus rumex: Review of traditional uses, phytochemistry and pharmacology. J. Ethnopharmacol. 2015;175:198–228.
 27. Pinela J, Carvalho AM, Ferreira ICFR. Wild edible plants: Nutritional and toxicological characteristics, retrieval strategies and importance for today's society. Food Chem. Toxicol. 2017;110:165–188.
 28. Upton R, Graff A, Jolliffe G, Länger R, Williamson E. American herbal pharmacopoeia: botanical

- pharmacognosy-microscopic characterization of botanical medicines; CRC Press: Boca Raton, FL, USA; 2016. ISBN 1420073281.
29. Emshanova SV, Potanina OG, Budanova EV, Chistyakov VV. (Eds.). Gosudarstvennaya farmakopeya rossiyskoy federatsii XIV izdaniye. In [State Pharmacopoeia of the Russian Federation XIV edition]. Ministry of Health of Russian Federation. Moscow, Russia. 2018;4:1844.
 30. Zhang H, Guo Z, Wu N, Xu W, Han L, Li N, Han Y. Two novel naphthalene glucosides and an anthraquinone isolated from *Rumex dentatus* and their antiproliferation activities in four cell lines. *Molecules*. 2012;17/1:843–850.
 31. Pareek A, Kumar A. *Rumex crispus* L.- A plant of traditional value. *Drug Discovery*. 2014;9/20:20–23.
 32. Ahmad SS, Erum S, Khan SM, Nawaz M, Wahid A. Exploring the medicinal plants wealth: a traditional medico-botanical knowledge of local communities in Changa Manga Forest, Pakistan. *Middle-East Journal of Scientific Research*. 2014; 20/12:1772–1779.
 33. Packer J, Brouwer N, Harrington D, Gaikwad J, Heron R, Elders Y.C, Ranganathan S, Vemulpad S, Jamie J. An ethnobotanical study of medicinal plants used by the Yaegl aboriginal community in northern New South Wales, Australia. *Journal of Ethnopharmacology*. 2012;139/1:244–255.
 34. Rivero-Cruz I, Acevedo L, Gurerrero J.A, Martinez S, Bye R, Miranda PR, Franzblau S, Timmermann BN, Mata R. Antimycobacterial agents from selected Mexican medicinal plants. *Journal of Pharmacy and Pharmacology*. 2005;57/9:1117–1126.
 35. Tavares L, Carrilho D, Tyagi M, Barata D, Serra AT, Duarte CMM, Duarte RO, Feliciano RP, Bronze MR, Chicau P, Espirito-Santo MD, Ferreira RB, dos Santos C.N. Antioxidant capacity of Macaronesian traditional medicinal plants. *Molecules*. 2010;15/4:2576–2592.
 36. Sahreen S, Khan MR, Khan RA. Comprehensive assessment of phenolics and antiradical potential of *Rumex hastatus* D. Don. roots. *BMC Complementary and Alternative Medicine*. 2014;14/1:1–11.
 37. El-Hawary SA, Sokkar NM, Ali ZY, Yehia MM. A profile of bioactive compounds of *Rumex vesicarius* L. *Journal of Food Science*. 2011;76/8:1195–1202.
 38. Shafiq N, Saleem M, Kousar S, Sahar M, Hussain SM, Jabeen F. Investigation of genus *Rumex* for their biologically active constituents. *Life Science Informatics Publications*. 2017;2/6:148-165.
 39. Abdul Khabir Khan. Phytochemical screening on the constituents of *Rumex obtusifolius*. Ph.D thesis. 2017;29.
 40. Gautam R, Karkhile KV, Bhutani KK, Jachak SM. Anti-inflammatory, cyclooxygenase (COX)-2, COX-1 inhibitory, and free radical scavenging effects of *Rumex nepalensis*. *Planta medica*. 2010;76/14:1564-1569.
 41. Lee NJ, Choi JH, Koo BS, Ryu SY, Han YH, Lee SI, Lee DU. Antimutagenicity and cytotoxicity of the constituents from the aerial parts of *Rumex acetosa*. *Biological and Pharmaceutical Bulletin*. 2005;28 /11:2158-2161.
 42. Al-Dhabi NA, Arasu MV, Park CH, Park SU. Recent studies on rosmarinic acid and its biological and pharmacological activities. *EXCLI J*. 2014;13:1192-1195.
 43. Caleja C, Barros L, Prieto MA, Barreiro MF, Oliveira M, Beatriz PP, Ferreira ICFR. Extraction of rosmarinic acid from *Melissa officinalis* L. by heat-, microwave- and ultrasound-assisted extraction techniques: A comparative study through response surface analysis. *Elsevier*. 2017;186:297-308.
 44. Abedini A, Roumy V, Mahieux S, Biabiany M, Standaert-Vitse A, Riviere C, Sahpaz S, Bailleul F, Neut C, Hennebelle T. Rosmarinic acid and its methyl ester as antimicrobial components of the hydromethanolic extract of *Hyptis atrorubens* Poit. (Lamiaceae). *Evid Based Complement Alternat Med*. 2013;4:604536.
 45. Gun-Dong Kim, Yong Seek Park, Young-Ho Jin, Cheung-Seog Park. Production and applications of rosmarinic acid and structurally related compounds. *Appl Microbiol Biotechnol*. 2015;99:2083–2092.
 46. Kim JK, Park SU. Flavonoids for treatment of Alzheimer's disease: An up to date review. *EXCLI*. 2021;20:495-502.
 47. Bankova V, Trusheva B, Popova M. Caffeic acid phenethyl ester (CAPE) Natural sources, analytical procedures and synthetic approaches. 2018;71/9:1157-1169.
 48. Alam M, Ahmed S, Elsbali AM, Adnan M, Alam S, Hassan MI, Pasupuleti VR.

- Therapeutic Implications of Caffeic Acid in Cancer and Neurological Diseases. *Front. Oncol.* 2022;12:860508.
49. Koyuncu I. Evaluation of anticancer, antioxidant activity and phenolic compounds of *Artemisia absinthium* L. Extract. *Cell Mol Biol.* 2018;64/3:25-34.
50. Jiang RW, Lau KM, Hon PM, Mak TCW, Woo KS, Fung KP. Chemistry and biological activities of caffeic acid derivatives from *Salvia miltiorrhiza*. 2005;12/2:237-246.
51. Chen S, Wang FF, Chen Y, Li MS, Zhang B, Luo JZ, Song XX, Li JJ, Qin F. Chemical constituents of *Corydalis saxicola*. *Chem Nat Compd.* 2022;58:119-121.
52. Qi J, Zhao P, Zhao L, Jia A, Liu Ch, Zhang L, Xia X. Anthraquinone derivatives from a sea cucumber-derived trichoderma sp. fungus with antibacterial activities. *Chem Nat Compd.* 2020;56:112–114.
53. Zhu YY, Yu G, Wang YY, Xu JH, Xu FZ, Fu H, Zhao YH, Wu J. Antiviral activity and molecular docking of active constituents from the root of *Aconitum carmichaelii*. *Chemistry of Natural Compounds.* 2019;55:189–193.
54. Qian YX, Kang JC, Luo YK, Yang XB, He J, Wang L. Secondary metabolites from endophytic pestalotiopsis microspora and their P-glycoprotein inhibitory activity. *Chemistry of Natural Compounds.* 2021;57:924–926.
55. Vu TT, Kim H, Tran VK, Dang QL, Nguyen HT, Kim H, Kim IS, Choi GJ, Kim JC. *BMC Complement Altern Med.* 2016;16/32.

© 2022 Shermatova et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86738>