



Proposal for a New Classification of Cesarean Sections for Developing Countries: A Multicenter Study in Five Maternity Hospitals in Dakar

**Niang Ndama^{a,b*}, Diouf Abdoul Aziz^{a,b,c}, Gueye Mamour^{a,b},
Faye Dieme Marie Edouard^{a,b} and Diouf Alassane^{a,b,c}**

^a *Gynecological and Obstetrical Clinic, Aristide Le Dantec University Hospital, Dakar, Senegal.*

^b *Cheikh Anta Diop University, Dakar, Senegal.*

^c *Center Hospitalier National de Pikine, Dakar, Senegal.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96464>

Original Research Article

Received: 25/10/2022

Accepted: 30/12/2022

Published: 10/01/2023

ABSTRACT

Objectives: The general objective of this study was to contribute to the study of the classification of cesarean sections. The specific objectives were to propose a new classification of cesarean sections based on the obstetrical complications specified by the WHO and to see the applicability of this classification, especially in Africa.

Methods: This is a prospective, descriptive and analytical study, consisting of a review of the records of patients received at the level of the target maternities and who had a caesarean section during the period from January 1, 2019 to December 31, 2019 (12 months). Our study was conducted at five level 2 and 3 maternity hospitals. All patients who had a cesarean section during the study period were included in the study, as well as those with a uterine rupture discovered during cesarean section in one of the study maternity units in the Dakar region.

*Corresponding author: E-mail: ndamaniang@hotmail.com;

We proposed a classification of cesarean section into four groups. In patients with interrelated causes for cesarean section, the most urgent primary indication was used to classify patients into one of the groups.

Results: Group I represented parturients with a direct obstetric complication; group II was for patients with an indirect obstetric complication; patients with a fetal complication were grouped in group III. Group IV was for prophylactic cesarean sections. With this distribution, we obtained the following proportions. Group I comprised the majority of indications with 40.9%, group II was the least represented with 3.5%; groups III and IV represented 32% and 24% respectively. Group I of our classification encompasses almost all major obstetrical emergencies and thus allows us to understand that our high rates of cesarean section in our facilities are justified.

Conclusion: Direct and indirect obstetrical complications are the main causes of maternal death. Taking these obstetrical complications into account in the cesarean section classification systems would contribute to a better control of maternal deaths.

Keywords: Caesarean section; obstetrical complications; maternal death; caesarean section classification; robson classification.

1. INTRODUCTION

Since 2000, maternal and child health has become a major concern for the international health community and has been chosen to represent two (2) of the eight (8) Millennium Health Goals (MDGs) [1]. This concern is especially relevant in Africa where unfavorable socio-economic conditions expose many pregnancy-related complications, with high maternal and perinatal morbidity and mortality rates. Among the factors contributing to this high rate of maternal and perinatal morbidity and mortality in developing countries is the problem of access to care, particularly to cesarean section.

Caesarean section is an essential surgical procedure in obstetrics, since there are approximately 100,000 caesarean sections per year in France and it remains the most common procedure in the world [2,3].

Since 1985, the WHO has considered the ideal rate of cesarean delivery to be between 10% and 15% [4]. Since that time, cesarean delivery has become increasingly common in both developed and developing countries. However, in recent years, governments and clinicians have expressed concern about the increase in cesarean deliveries and the potential negative consequences for maternal and child health, and the need to reconsider the cesarean section rate recommended in 1985 has been raised [4]. The cesarean section rate differs between the population and hospital levels, and the need for a uniform classification of cesarean sections has been suggested to better control the rate.

Several classifications have been described and used, but none has achieved global consensus.

Among those existing, the WHO proposes to adopt the Robson classification system as the international reference system for the evaluation, monitoring and comparison of cesarean section rates within and between health care facilities [5,6].

However, this Robson classification has limitations because it does not take into account the existence of a surgical pool or certain extreme emergencies in the indication for cesarean section.

It is in this context that we considered it necessary to propose a new classification of cesarean sections taking into account these elements; and as an example we will base ourselves on a prospective multicenter study by classifying the different indications according to our new parameters.

The general objective of this study was to contribute to the study of the classification of cesarean sections.

The specific objectives were:

- to propose a new classification of cesarean sections based on obstetrical complications.
- To see the applicability of this classification especially in Africa.

2. METHODOLOGY

2.1 Type and Period of Study

We conducted a multicenter prospective descriptive and analytical study classifying the indications for cesarean sections in 5 surgical

maternities in Dakar according to the different classifications of cesarean sections and according to obstetric complications. Subjects were selected according to an exhaustive sampling that identified all cases of cesarean section during the period from January 1, 2019, to December 31, 2019, i.e., a period of 12 months. All patients who had a cesarean section during the study period were included in the study, as well as those with a uterine rupture discovered during the cesarean section. Patients referred postoperatively to the selected facilities were not included in the study.

The data collected came from two sources: the first source was the medical records and surgical reports of patients at the following facilities, using a previously established form: the Gynecological and Obstetrical Clinic, the Ouakam Military Hospital, and the Institute of Social Hygiene. The second source for the Centre de Santé Philippe Maguilen Senghor and the Centre Hospitalier National de Pikine came from the File Maker Pro software database used in these two facilities.

For each patient, we collected the following parameters: sociodemographic data, main indications for cesarean section, classification of the main indications according to the different classification systems for cesarean section, and then we classified this cohort according to obstetrical complications.

2.2 Operational Definitions

2.2.1 Direct obstetric complications

These are complications that occur during pregnancy, childbirth, or the postpartum period. When not treated effectively from the outset, they become serious emergencies with a high risk of maternal and fetal death. The seven major direct obstetric complications that account for 85% of maternal deaths are: antepartum and postpartum hemorrhage, prolonged labor/dystocia, puerperal infection, abortion complications, preeclampsia/eclampsia, ectopic pregnancy, and uterine rupture. In our study of a cesarean section classification that constitutes care at the end of pregnancy, direct complications such as abortion complications and ectopic pregnancy are excluded. Similarly, puerperal infection is not taken into account because it occurs after the cesarean section.

The direct complications retained to enable a new classification are: antepartum hemorrhage,

prolonged labor/ dystocia, preeclampsia/eclampsia, and uterine rupture.

2.2.2 Indirect obstetrical complications

These are complications that result from a pre-existing disease or condition that developed during pregnancy without direct obstetric causes, but which was aggravated by the physiological effects of pregnancy.

2.2.3 Fetal complications

These are complications concerning the foeto-annexal state leading to a caesarean section in the parturient.

2.2.4 Classification according to complications

These definitions provide an understanding of the breakdown we want to develop.

Each of these complications will constitute a group, and patients who receive a Caesarean section in an uncomplicated setting will constitute a separate group.

2.3 Data Entry and Analysis

The data collected from the CGO, HMO and IHS were entered using CsPro software. For the CSPMS and CHNP, they were processed directly with FileMaker Pro software.

The analysis was performed with the following software: Excel 2010, Epi info 7.2 and IBM SPSS Statistics 21.

In the descriptive analysis, the qualitative variables were described by frequency tables, bar graphs, and pie charts. Quantitative variables were described by their positional (Mean, median and mode) and dispersion (Standard deviation, extremes) parameters.

The bi-variate analysis allowed us to search for associations between the variables while using the appropriate statistical tests according to their applicability conditions. The alpha risk of error was set at 5% and the CI at 95%.

3. RESULTS

The proportion of mandatory cesarean sections was 48.7% (or 1928 patients) as shown in Fig. 1.

Concerning the distribution of cesarean sections according to the classification of entry into labor,

we noted a predominance of mixed indications (41.3%), which concerned 1633 patients. Fig. 2 illustrates this distribution.

The proportion of caesarean sections in group 5 was higher and represented 27% of the caesarean section indications (1068 patients) as shown in Fig. 3. These were cases of monofetal pregnancies in a scarred uterus with a fetus in cephalic presentation. Transverse and oblique presentations belonging to group 9 were rare in our series (0.9%).

We noted a predominance of caesarean sections with an extraction time of less than 60 minutes, known as "green code", i.e. 42.3% (n=1674). The following Fig. 4 shows the details of this distribution according to the 3 caesarean section codes.

We propose to classify the indications for Caesarean section into four groups.

The first group consists of patients whose indications represent direct obstetric complications at the time of delivery that may cause maternal death.

The second group consists of patients whose indications represent indirect obstetric complications.

The third group includes patients whose indications for Caesarean section are related to fetal complications.

The fourth group includes indications for caesarean sections without the context of complications (prophylactic, convenience...). This is shown in Table 1.

Following this distribution, we obtain with our cohort the different proportions represented in Fig. 5.

Group I is the most represented with 1618 patients (40.9%) with direct obstetric complications at the time of caesarean section.

This was followed by patients with fetus-related Caesarean sections (32%) and those with abnormal pelvis, scarred uterus or elective Caesarean sections (23.6%). Group II was the least represented with 3.5% each. For each group, the distribution of indications for caesarean section is shown in the following Fig. 5.

Caesarean sections for dystocia and severe eclampsia/pre-eclampsia were more represented with 52% and 28% respectively. However, antepartum haemorrhage represented a significant rate of 19%.




Fig. 1. Distribution of caesarean sections according to the Maillet and Boisselier classification




Fig. 2. Distribution of caesarean sections according to classification by entry into labour




Fig. 3. Distribution of patients according to the different groups of the Robson classification




Fig. 4. Distribution of caesarean sections according to the Lucas classification




Fig. 5. Distribution of indications for GROUP I caesarean sections




Fig. 6. Distribution of indications for caesarean sections in GROUP II




Fig. 7. Distribution of indications for caesarean sections in GROUP III




Fig. 8. Distribution of indications for caesarean sections in GROUP IV

Table 1. Distribution of caesarean section indications according to obstetrical complications

Groups	Indications for cesarean section
Group I	Direct obstetrical complications
Group II	Indirect obstetrical complications
Group III	Complications fœtales
Group IV	Uncomplicated cesarean sections

In group II, comprising patients whose main indication is related to indirect obstetrical complications, the indications for arterial hypertension, diabetes and uterine myomatosis are in the majority with 30.7%, 21.4% and 16.4% respectively. Indications for acute fetal distress/non-reassuring fetal status for the fetal complications group were in the majority with 34.5%. Caesarean sections for intrauterine

growth retardation were the least represented with 2.4%. In group IV, indications for caesarean section for scarred/multi-scarred uterus dominated with 58% or 539 patients.

4. DISCUSSION

Maternal death is defined as the death of a woman during pregnancy or during the 42 days following the end of pregnancy, regardless of its duration or location. The cause of death is neither accidental nor incidental. Death is due to, or aggravated by, the pregnancy, or by the necessary care [4]. Death most often occurs as a result of complications. Obstetric morbidity has been defined as any condition in a pregnant woman (regardless of the location and duration of the pregnancy) or in a woman who has been in labour for less than 42 days, due to a cause related to or aggravated by the pregnancy or its management, but not related to an accidental or incidental cause [7]. Nearly 80% of maternal deaths are due to obstetric complications and the therapeutic armoury includes caesarean section to improve the prognosis of these parturients (severe maternal morbidity) [8]. Several classification systems have been proposed but none of them took into account obstetric complications in isolation. In its 2014 statement, the World Health Organization concluded that the Robson classification is the system that best meets current local and international needs for classifying indications for Caesarean section, after conducting a systematic review of different classifications. As with other caesarean section classifications, WHO hopes that this classification will help health care facilities to evaluate the effectiveness of strategies or interventions to optimize caesarean section use [9]. However, is the use of this classification in sub-Saharan Africa appropriate? These direct and indirect obstetric complications represent the main causes of maternal death in this region of the world [7]. To better combat maternal mortality, the WHO has identified seven causes of obstetric complications that can directly cause maternal death and indirect causes of maternal death.

To counter this high mortality and the morbidity linked to these obstetric complications, preventive and curative strategies have been implemented, in particular caesarean sections [10]. It is therefore important to take these complications into account in the classification systems, as for the majority, Caesarean section is part of the curative strategy.

Given the importance of this subject, we felt it necessary to propose a classification taking into account the obstetrical complications developed by the WHO.

To do this, we developed 4 groups numbered from 1 to 4. By analysing the distribution of Caesarean section indications in our cohort according to our classification, we find direct peripartum obstetrical complications in the foreground, followed by fetal complications. This is understandable in our context where antepartum haemorrhage, vasculo-renal syndromes, dystocia and uterine rupture represent the first emergencies and occupy a non-negligible proportion of caesarean sections [11]. In the literature [4,11,8,12,7], these complications are the biggest contributors to maternal mortality. Prual et al in their study found a lethality of 3.3% for haemorrhage, dystocia and complications of hypertension [13].

The low proportion of indirect obstetric complications (3.5%) is explained by two facts. On the one hand, we note the regression of many maternal pathologies associated with pregnancy. For example, this is the case of malaria, the incidence of which in pregnant women represented 115 women in the Dakar region for a total number of 7155 cases in 2019, i.e. a rate of 1.6% [14]. On the other hand, in our classification, cases of indirect complications were only taken into account when they were isolated. If they were associated with direct complications, these took precedence over indirect causes. This may justify the low rate of patients in this group.

In our study and in the majority of publications in the literature regarding Robson's classification, the main contributing groups for caesarean section were groups 5, 1 and 10; with group 5 being the majority [2,15,16,17,18]. Robson's group 5 represents patients with a scarred uterus in a full-term pregnancy, regardless of the mode of induction of delivery. But this group does not differentiate patients coming for emergency labour and therefore potentially dystocic labour from parturients who are not in labour. In our study, we separated these two categories of parturients who had a scarred uterus, some in labour had dystocia while those not in labour had prophylactic caesarean sections. This separation allows us to see those who require more urgent and rapid management. So part of our sample was in group I and part in group IV.

Group 1 of Robson's classification includes primiparous women in spontaneous labour with a full term pregnancy. These primiparous women in spontaneous labour had a caesarean section for dystocia or a fetal complication such as distress in more than 50% of cases. Indeed, in our classification, dystocia was in the majority with a proportion of 52% in group I and cardiac rhythm anomalies were in the majority in group III with 32.54%. Group 10 of Robson's classification represents non term pregnancies. Therefore any woman with a non term pregnancy whatever the reason for caesarean section is in this group. Obstetric complications such as antepartum haemorrhage are major contributors to prematurity, whereas prematurity is not an indication for caesarean section. Some of these parturients with a pregnancy that is not full term are therefore in our group I because of the seriousness of the pathology.

This makes it possible to understand the predominance of group I by comparing our classification with that of Robson.

Ours is perfectly adapted to our context, as is Robson's, but allows us to better understand where the rate of caesarean sections in hospitals is highest. And it shows that although this rate is said to be high in facilities exceeding the WHO standards, it is justified and even indispensable.

5. CONCLUSION

Comparing this new classification with Robson's, we can see that Robson's dominant groups were all found in the 4 groups of our classification but with a different distribution of proportions. Group I of our classification includes almost all major obstetric emergencies and therefore allows us to understand that our high caesarean section rates in our facilities are justified. This allows us to recommend the introduction of this new classification in clinical practice, especially in sub-Saharan Africa, in order to easily compare our Caesarean section indications and see that they are justified.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

All information was collected anonymously. Prior authorization from the hospital management or

the chief medical officer, as well as the approval of the head of the maternity department was obtained at the level of the structures targeted for our study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO | Millennium Development Goals (MDGs) [Internet]. WHO. World Health Organization; [cited 2020 Dec 15]. Available: https://www.who.int/topics/millennium_development_goals/about/en/
2. Beucher G, Dolley P, Lévy-Thissier S, et al. Maternal benefits and risks of attempted vaginal delivery compared to planned cesarean section in case of history of cesarean section. *Journal of Obstetrics Gynecology and Reproductive Biology*. 2012;41:708-26.
3. Racinet C, Favier M. *Caesarean section: Indications, techniques, complications* Masson. Paris 1984. 1984;
4. Betrán AP, Ye J, Moller AB, Zhang J, Gülmezoglu AM, Torloni MR. The increasing trend in caesarean section rates: global, regional and national estimates: 1990-2014. *PLoS one*. 2016 Feb 5;11(2):e0148343.
5. Alsayegh E, Bos H, Campbell K, et al. No 361 - Cesarean on maternal request. *Journal of Obstetrics and Gynecology Canada*. 2018;40:972-7.
6. Vogel JP, Betrán AP, Vindevoghel N, et al. Use of the Robson classification to assess caesarean section trends in 21 countries: a secondary analysis of two WHO multicountry surveys. *The Lancet Global Health*. Elsevier. 2015;3:e260-70.
7. Dumont A, De Bernis L, Bouillin D, et al. Maternal morbidity and qualification of health personnel: comparison of two different populations in Senegal. *Journal of Obstetrics Gynecology and Reproductive Biology*. 2002;31:70-9.
8. CEFORP, UNFPA, MSAS. Rapid assessment of the availability, use and quality of emergency obstetric and neonatal care in Senegal, 2015-2016. Dakar, Senegal: Ministry of Health and Social Action. 2017 Jan;85.

9. Haydar A, Vial PY, Baud D, et al. Evolution of the caesarean section rate in a Swiss university maternity hospital according to the Robson classification. *Rev Med Switzerland*. 2017;13:1846-51.
10. High Authority of Health. Indications for scheduled caesarean at term. January 2012. *La Revue Sage-Femme*. 2012;11: 244-50.
11. Cavallaro FL, Cresswell JA, França GV, et al. Trends in Caesarean delivery by country and wealth quintile: cross-sectional surveys in southern Asia and sub-Saharan Africa. *Bull World Health Organ*. 2013;91:914-922D.
12. Costa ML, Cecatti JG, Souza JP, et al. Using a Caesarean Section Classification System based on characteristics of the population as a way of monitoring obstetric practice. *Reproductive Health*. 2010;7: 13.
13. FM drama. Attraction of primary health care services and HIV testing centers in Dakar (Senegal): Improving the approach by polarization area. *EPS*. 2006;361-71.
14. Diniz SG, Chacham AS. "The cut above" and "the cut below": the abuse of caesareans and episiotomy in São Paulo, Brazil. *Reproductive health matters*. Elsevier; 2004;12:100-10.
15. BOUVIER-COLLE M-H. Maternal mortality. *Obstetrics*. 2001;5:9.
16. Chaillet N, Dumont A. Evidence-based strategies for reducing cesarean section rates: a meta-analysis. *Birth*. Wiley Online Library. 2007;34:53-64.
17. Cissé CT, Ngom PM, Guissé A, et al. Reflections on the evolution of caesarean section rates in Africa: example of the University Hospital of Dakar between 1992 and 2001. *Obstetrics Gynecology & Fertility*. 2004;32:210-7.
18. Diallo FB, Diallo MS, Bangoura S, et al. Cesarean: Factor for reducing fetomaternal morbidity and mortality. *Black African Medicine*. 1998;45:359-64.

© 2023 Ndama et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96464>