



Journal of Advances in Mathematics and Computer Science

Volume 38, Issue 4, Page 33-41, 2023; Article no.JAMCS.96295

ISSN: 2456-9968

(Past name: British Journal of Mathematics & Computer Science, Past ISSN: 2231-0851)

On the Pendant Number of Certain Graphs

P. G. R. S. Ranasinghe^{a*}
and L. R. M. K. R. Jayathilaka^a

^aDepartment of Mathematics, Faculty of Science, University of Peradeniya, Peradeniya, Sri Lanka.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMCS/2023/v38i41756

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96295>

Received: 21/12/2022

Accepted: 09/02/2023

Published: 27/02/2023

Original Research Article

Abstract

The present study investigates the pendant number of certain graph classes; complement, line graphs, and total graphs. The pendant number is the minimum number of end vertices of paths in a path decomposition of a graph. A path decomposition of a graph is a decomposition of it into subgraphs; i.e., a sequence of a subset of vertices of the graph such that the endpoints of each edge appear in one of the subsets and each vertex appears in an adjacent sub-sequence of the subsets.

Keywords: Complement; decomposition; line graph; pendant number; total graph.

2010 Mathematics Subject Classification: 53C25, 83C05, 57N16.

**Corresponding author: E-mail: rajithamath@sci.pdn.ac.lk;*

J. Adv. Math. Com. Sci., vol. 38, no. 4, pp. 33-41, 2023

1 Introduction

The motivation for the present work comes from the study by Sebastian et al. [1], where they introduced the pendant number for certain fundamental graph classes. Our research is carried out specifically to determine this parameter for derived graphs such as complement, line graphs, and total graphs. The pendant number Π_p is the minimum number of end vertices of paths in a path decomposition of G [2]. Graph decomposition problems rank among the most prominent areas of research in graph theory which is a collection of edge-disjoint sub-graphs such that every edge belongs to exactly one sub-graph. All graphs that are considered in this paper are undirected, simple, finite, and connected. The graph theoretic terminologies used in this paper are as appeared in [3] and [4].

2 Materials and Methods

In this section, terminologies of the types of graphs that we consider are presented as appeared in [1].

2.1 Complement Graphs

Definition 2.1. If G is a simple graph with vertex set $V(G)$, its complement \bar{G} is the simple graph with vertex set $V(G)$ in which two vertices are adjacent if and only if they are not adjacent in G .




Fig. 1. A Graph and its complement

For example, Fig. 1. shows a graph and its complement. Note that the complement of a complete graph is a null graph and that the complement of a complete bipartite graph is the union of two complete graphs.

2.2 Line Graphs

Given a graph G with at least one edge, the line graph $L(G)$ is that graph whose vertices are the edges of G , with two of these vertices being adjacent if the corresponding edges are adjacent in G [5]. The operation of going from adjacent edges in G to adjacent vertices in $L(G)$ is indicated in Fig. 2.

2.3 Total Graph

Definition 2.2. The total graph $T(G)$ of a graph G is the graph that has as its vertex set $V(G) \cup E(G)$ with a pair of these elements are adjacent if in the graph G they are either

1. adjacent vertices,
2. adjacent edges, or
3. an incident vertex and edge.




Fig. 2. A Graph G and the corresponding line graph $L(G)$

This is illustrated schematically in Fig. 3. below. It follows from the definition that the total graph of graph G is spanned by the sum (disjoint union) $G + L(G)$ of G and its line graph $L(G)$ [6].




Fig. 3. C_5 graph and its total graph $T(C_5)$

2.4 Pendant Number

Definition 2.3. The pendant number of a graph G , denoted by $\Pi_p(G)$, is the least number of vertices in a graph such that they are the end vertices of a path in a given path decomposition of a graph.

Theorem 2.1. Let G be a connected graph with n vertices. If G has l number of odd degree vertices, then $l \leq \Pi_p(G) \leq n$.

2.5 Pendant Number of Acyclic Graphs

An acyclic graph is a graph having no graph cycles. An acyclic-connected graph is called a tree. Since every path P_n has exactly two pendant number of vertices, we have $\Pi_p(G) = 2$ from the following proposition.

Theorem 2.2. Let T be a tree with $n \geq 2$ vertices. Then $\Pi_p(T) = 2$ if and only if T is a path P_n .

2.6 Pendant Number of Cyclic Graphs

Now, we shall state some properties of cyclic graphs which are essential in our study.

Proposition 2.1. If G is the cycle C_n on $n \geq 3$ number of vertices, then $\Pi_p(G) = 2$.

Theorem 2.3. *If G has a Hamilton decomposition of the first kind, then $\Pi_p(G) = 2$.*

Proposition 2.2. *For a complete graph K_n on n vertices, where n is odd, there will be $\frac{n-1}{2}$ edge-disjoint cycles of length n .*

The following theorem discusses the pendant number of a complete graph of odd order.

Theorem 2.4. *For $n \geq 3$, $\Pi_p(K_n) = 2$ if and only if n is odd.*

3 Results and Discussion

Now, we will provide a detailed study on the pendant number of line graphs, total graphs, and complement graphs as suggested in [1].

3.1 Pendant Number of Complement Graphs

In graph theory, a cyclic graph C_n , sometimes simply known as an n -cycle, is a graph on n nodes containing a single cycle through all nodes. When we consider the complement graphs of cycle graphs, those can be obtained by removing a cycle from the complete graph K_n (top) and in the "standard" circulant graph from the bottom. By observing the pattern, the following table can be constructed.

Table 1. Complement of the cycle graph for $n \leq 10$

n	$V(G)$	$E(G)$	$deg(G)$	Hamiltonian	
				Yes/No	Number of cycles
5	5	5	2	✓	1
6	6	9	3	-	-
7	7	14	4	✓	2
8	8	20	5	-	-
9	9	27	6	✓	3
10	10	33	7	-	-

Theorem 3.1. *Let G be the Cyclic Graph C_n . For $n \geq 5$ number of vertices, the following holds:*

$$\Pi_p(\bar{G}) = 2 \text{ if and only if } n \text{ is odd.} \tag{3.1}$$

Proof. For $n \geq 1$, consider the complement graph of C_{2n+1} . Clearly, it has Hamilton decomposition of the first kind, with n -Hamiltonian cycles. The degree of the complement graph of a cyclic graph is $2n$. Hence, by Theorem 2.3, $\Pi_p(\bar{G}) = 2$.

Conversely, take a complete graph C_{2n+1} on $n \geq 1$ vertices such that $\Pi_p(\bar{G}) = 2$. Assume, if possible, that n is even. Then, all its vertices are of odd degree $n - 1$, a contradiction. Hence, n cannot be even, completing the proof. \square

Theorem 3.2. *For the complement graph of cyclic graph C_{2n} on $n \geq 3$ number of vertices,*

$$\Pi_p(\bar{G}) = 2n. \tag{3.2}$$

Proof. Consider the complement graph of C_{2n} for $n \geq 3$. Note that, \bar{C}_{2n} is a regular graph with $2n + 1$ vertices. Hence \bar{C}_{2n} has odd degree vertices by Theorem 2.1. Therefore the pendant number is $2n$. \square

3.2 Pendant Number of Total Graphs

In this subsection, the pendent number of the total graphs of P_n , K_n , C_n , and $K_{1,n}$ is investigated.

Note that the path graph P_n is a tree with two nodes of vertex degree one.

The path graphs and corresponding total graphs of path graphs can be visualized as follows.




Fig. 4. P_n graph and the corresponding total graph $T(P_n)$

Theorem 3.3. Let G be a tree with $n \geq 3$ number of vertices, then

$$\Pi_p(T(G)) = 2 \text{ if and only if } G \text{ is the path } P_n. \tag{3.3}$$

Proof. Note that, the total graph of P_n has $2n - 1$ number of vertices and it is a connected graph. Also, $T(P_n)$ on $n \geq 3$ has 2 degree 3 vertices. Hence by Theorem 2.1, the pendant number is 2. \square

Let K_n be a complete graph with n vertices and $T(K_n)$ be the total graph of a complete graph K_n . Note that $T(K_n)$ is a strongly regular graph [7].

Theorem 3.4. For the complete graph K_n on $n \geq 3$ vertices,

$$\Pi_p(T(K_n)) = 2. \tag{3.4}$$

Proof. Note that, the total graph of the complete graph K_n is $2(n - 1)$ regular graph. Hence it is clearly a graph with Hamilton decomposition of the first kind. Hence by Theorem 2.3, $\Pi_p(T(K_n)) = 2$. \square

Theorem 3.5. Let G is a star graph $K_{1,n}$ on $n \geq 2$ vertices. If n is even, then

$$\Pi_p(T(G)) = n. \tag{3.5}$$

Proof. Note that, the total graph of a star graph $K_{1,n}$ on $n \geq 2$. When n is even $K_{1,n}$ has n number of odd vertices. Because $K_{1,n}$ is a connected graph by Theorem 2.1. Therefore the pendant number is n . \square

Let's look at the total graphs of C_n . When $n = 5$, the total graph of C_5 can be drawn as Fig. 3. For the general case, the total graph of C_n can be visualized as Fig. 4. [8].

Theorem 3.6. If G is the cyclic graph C_n , then for $n \geq 3$ number of vertices, the following holds:

$$\Pi_p(T(G)) = 2. \tag{3.6}$$

Proof. Note that, the total graph of a cycle C_n is a 4-regular graph. Thus it is clear that C_n is a graph with Hamilton decomposition of the first kind. Hence by Theorem 2.3, the pendant number is 2. \square




Fig. 5. C_n graph and the corresponding total graph $T(C_n)$

3.3 Pendant Number of Line Graphs

In this section, we present our result on the pendant number of line graphs.

Theorem 3.7. *If G is the cycle C_n on $n \geq 3$ vertices, then*

$$\Pi_p(L(G)) = 2. \quad (3.7)$$

Proof. Note that, the line graph of the cycle graph C_n is isomorphic to C_n itself. From Proposition 2.1, if G is the cycle C_n on $n \geq 3$ vertices, then $\Pi_p(G) = 2$. Since $L(C_n) = C_n$, the pendant number of the line graph of C_n is 2. \square

Theorem 3.8. *Let T be a tree with $n \geq 3$ number of vertices. If T is a path P_n , then*

$$\Pi_p(L(T)) = 2. \quad (3.8)$$

Proof. Since it has two pendent vertices, the path graph P_n is a tree with two nodes of vertex degree one. Note that the line graph of P_n is isomorphic to P_{n-1} . Hence by Theorem 2.5, the pendant number is 2. \square

Theorem 3.9. *If G is the star graph S_n on $n \geq 3$ number of vertices, then*

$$\Pi_p(L(G)) = \begin{cases} 2 & \text{if } n \text{ is even,} \\ n & \text{if } n \text{ is odd.} \end{cases} \quad (3.9)$$

Proof. Note that, the line graph of the star graph S_n is isomorphic to the complete graph K_{n-1} . From Proposition 2.2 and Theorem 2.4, the pendant number of a complete graph K_n is equal to n when it has odd order. Furthermore, for $n \geq 3$, $\Pi_p(K_n) = 2$ if and only if n is even. Therefore,

$$\Pi_p(L(G)) = \begin{cases} 2 & \text{if } n \text{ is even,} \\ n & \text{if } n \text{ is odd.} \end{cases} \quad (3.10)$$

\square

Our results can be extended to 3-dimensional graphs as well.

Theorem 3.10. *If G is a tetrahedral graph, then*

$$\Pi_p(L(G)) = 2. \quad (3.11)$$

Proof. Note that, the line graph of the tetrahedral graph is the octahedral graph. The Hamiltonian cycle covers the entire vertex set of G and there are three cycles. Since the pendant number of a cyclic graph C_n is 2 and each C_n has common vertices, the pendant number of $L(G)$ is equal to 2. \square

Fig. 6. shows the degree of the line graph of the bipartite graph for fixed $n = 2$.




Fig. 6. Bipartite graph and corresponding line graph for fix $n = 2$

By observing the graphs and line graphs the following theorem can be obtained.

Theorem 3.11. *If G is a complete bipartite graph $K_{m,n}$ with $m \leq n$, then the degree of $L(G)$ can be generalized as follows:*

$$\text{deg}(L(G)) = \begin{cases} m + n - 2 & \text{if } m < n, \\ 2(n - 1) & \text{if } m = n. \end{cases} \quad (3.12)$$

Theorem 3.12. *If G is the complete bipartite graph $K_{m,n}$ with $m \leq n$, then*

$$\Pi_p(L(G)) = \begin{cases} mn & \text{if } \text{deg}(L(G)) \text{ is odd,} \\ n^2 & \text{if } \text{deg}(L(G)) \text{ is even.} \end{cases} \quad (3.13)$$

Proof. Case 1: $m < n$, if m is odd and n is even or m is even and n is odd
 Now $L(G)$ is a connected graph with odd-degree vertices. Hence all vertices have odd-degree by Theorem 2.1, and $\Pi_p(L(G)) = x$, where x is the number of vertices in $L(G)$ and it is mn .
 Case 2: $m < n$, if both m and n are both odd or both are even

Now $L(G)$ is a connected graph with even degree vertices and there are no odd degree vertices.

Case 3: $m = n$

Here $L(G)$ is a connected graph with n^2 number of even degree vertices.

By considering the above three cases we get, the pendant number as required. \square

Theorem 3.13. *If G is a cubic graph, then*

$$\Pi_p(L(G)) = 2. \quad (3.14)$$

Proof. Note that, the line graph of the cubic graph is the cuboctahedral graph. It is a Hamiltonian, and a regular graph. Hamiltonian cycle covers the entire vertex set of G and there are x cycles that have x -fold symmetry. Since the pendant number of the cyclic graph C_n is 2 and each C_n has common vertices, the pendant number of $L(G)$ is equal to 2. \square

4 Conclusions

The present study aimed to determine the pendant number of certain derived graph classes, including the complement, total, and line graphs. The study can be extended to other general graph classes. Although there is no direct relationship between the pendant number and other well-known graph parameters, it was observed that the pendant number highly depends on the number of odd-degree vertices in a graph. Further studies could compare the pendant number with other graph parameters, such as the domination number and graph diameter. The concept of pendant number can be applied in combination with path decomposition, which has applications in Computer Science and Engineering, including network design [9] and graph algorithms [10]. One of the recent studies along these lines is the study of the pendant number of corona products and rooted products of paths and cycles [11]. Also, in [12], the pendant number of regular graphs, complete r -partite graphs, line graphs, total graphs, and line graphs of total graphs are studied. Our approach is different than to [12] in that we used Hamiltonian cycles to establish our results.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Sebastian JK, Kureethara JV. Pendant number of graphs. International Journal of Applied Mathematics. 2018;31(5):679-689.
- [2] Heinrich K. Path decomposition. Le matematiche. 1992;47(2):241-258.
- [3] West DB. Introduction to Graph Theory. Prentice hall Upper Saddle River. 2001;2.
- [4] Barnes JA, Harary F. Graph theory in network analysis. Social Networks. 1983;5(2):235-244. Elsevier
- [5] Beineke LW, Bagga JS. Line graphs and line digraphs. Springer International Publishing. Developments in Mathematics; 2021. ISBN: 9783030813840 Available: <https://books.google.lk/books?id=-Ip-zgEACAAJ>
- [6] Belardo F, Stanić, Zoran, Zaslavsky T. Total graph of a signed graph; 2019. arXiv preprint arXiv:1908.02001
- [7] Joseph AK. Few results on total graph of complete graph. International Advanced Research Journal in Science, Engineering and Technology. 2019;6(72-78).

- [8] Ali MM, Panayappan S. Cycle multiplicity of total graph of C_n , P_n , and $K_{1,n}$. *International Journal of Engineering, Science and Technology*. 2010;2(2):54-58.
- [9] Path decomposition under a new cost measure with applications to optical network design. Anshelevich E, Zhang L. *Algorithms-ESA 2004: 12th Annual European Symposium*, Bergen, Norway, September 14-17, 2004. *Proceedings 12*. 2004;28-39. Springer
- [10] Steen MV. *Graph theory and complex networks. An Introduction*. 2010;144.
- [11] Sebastian KJ, Kureethara JV, Naduvath S, Dominic C. A study on the pendant number of graph products. *Acta Universitatis Sapientiae, Informatica*. 2019;11(1):24-40.
- [12] Kottarathil J, Naduvath S, Kureethara JV. *The pendant number of line graphs and total graphs*. *Data Science and Security*. Springer Nature Singapore. 2022;457-468. Singapore ISBN: 978-981-19-2211-4

© 2023 Ranasinghe and Jayathilaka; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/96295>