



Impact of Anemia during Pregnancy on Maternal and Perinatal Outcomes

Chandana Narayanaswamy ^{a++*} and Vrinda K Patil ^{a#}

^a Department of OBG, SDM College of Medical Sciences, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113535>

Minireview Article

Received: 14/12/2023

Accepted: 21/02/2024

Published: 27/02/2024

ABSTRACT

This literature review aims to elucidate the impact of anemia during pregnancy on maternal and perinatal outcomes. Anemia, a common condition in pregnant women, has been associated with adverse outcomes such as preterm birth, low birth weight, and maternal mortality. By examining various studies and reports, this review seeks to provide a comprehensive understanding of the correlation between anemia and these outcomes, and the potential strategies for prevention and management.

Keywords: Anemia impact; perinatal outcomes; preterm birth; maternal mortality.

1. INTRODUCTION

Anemia, defined as low hemoglobin concentration, is a significant public health problem affecting nearly half of all pregnant women globally [1]. The prevalence is

highest in low- and middle-income countries (LMICs), with rates over 40% in many regions [2]. Iron deficiency accounts for most cases, while other causes include nutritional deficiencies, infections, and hematologic disorders [3].

⁺⁺ Junior Resident;

[#] Associate Professor;

*Corresponding author: E-mail: dollychinna@outlook.com;

Anemia in pregnancy can adversely impact both maternal and perinatal health. However, associations may depend on the severity of anemia, timing during gestation, and effectiveness of treatment interventions currently in place. This literature review will summarize recent evidence on implications of maternal anemia in pregnancy, with a focus on maternal mortality and morbidity.

2. MATERIALS & METHODS

The materials for this literature review consist of peer-reviewed articles, clinical trials, and reports published in reputable journals and databases. The methods involve a systematic search of databases such as PubMed, Cochrane Library, and Google Scholar using relevant keywords such as “anemia”, “pregnancy”, “maternal outcomes”, and “perinatal outcomes”. The selection criteria include studies that specifically investigate the impact of anemia during pregnancy on maternal and perinatal outcomes. The data from these studies are then analyzed and synthesized to draw conclusions and recommendations. This review adheres to the PRISMA guidelines for systematic reviews and meta-analyses.

3. GLOBAL BURDEN AND ETIOLOGIES

According to the latest estimates, 29% of non-pregnant women globally have anemia, increasing to 40% during pregnancy [1]. Regional prevalence in pregnant women ranges from 17% in high-income regions to 48% in LMICs in Southeast Asia and 65% in sub-Saharan Africa [1]. Approximately half of cases are thought to be attributable to iron deficiency [4].

Iron requirements increase significantly during pregnancy to support fetal development. Without sufficient intake or stores, pregnant women can quickly become deficient, developing iron-deficiency anemia. Additional nutrient deficiencies exacerbating anemia risk include vitamins A, B12 and folate. Infections like malaria, HIV and hookworm contribute in endemic regions. Thalassemias and hemoglobinopathies are prevalent genetic hemoglobin disorders in certain ethnic populations [3].

4. ASSOCIATIONS BETWEEN ANEMIA SEVERITY AND MATERNAL MORBIDITY/MORTALITY

The severity of anemia appears important in determining maternal risks. Rahman et al. reported a 60% increase in the pooled odds of maternal mortality with any anemia, rising to 2.36-fold for severe cases (hemoglobin <7 g/dL) specifically [5]. A 2018 multi-country study found maternal mortality nearly doubled for women with hemoglobin <6 g/dL compared to 6.0-11.9 g/dL after adjustment for confounders [6].

In regards to morbidities, a meta-analysis found that compared to no anemia, mild and moderate cases increased the odds of blood transfusion 1.26 to 2-fold, while severe anemia escalated risk over 8-fold [7]. Higher transfusion rates align with clinical management guidelines. However, for other morbidities like postpartum hemorrhage, sepsis and preeclampsia, associations are less consistent. A 2015 review showed no clear link between anemia and preeclampsia, while other cohort data does suggest a potential U-shaped relationship similar to the recent India trial [8,9].

Anemia during pregnancy requires careful management, as the outcomes can vary depending on the etiology. The symptoms and risks associated with anemia during pregnancy are well-documented, and include fatigue, weakness, and increased susceptibility to infections. Furthermore, anemia can lead to adverse pregnancy outcomes such as preterm birth and low birth weight [10,11,12,13]. Therefore, the management of anemia during pregnancy is crucial. Current treatment strategies focus on the supplementation of iron, cobalamin, and folate [14,15,16]. However, more research is needed to optimize these strategies and to explore new treatment modalities. The ultimate goal is to improve both maternal and perinatal outcomes in the context of anemia during pregnancy.

5. RECENT FINDINGS ON ANEMIA AND MATERNAL OUTCOMES IN RURAL INDIA

Bone et al. conducted an important new study assessing associations between early pregnancy anemia and maternal morbidity in over 11,000 women participating in a trial across Karnataka State, India [11]. Despite very high rates (>85%) of anemia, nearly all women received iron

supplementation by delivery, reflecting India's public health policy for routine antenatal administration.

Overall 5% of women experienced a major morbidity, most often (in order): blood transfusion, antepartum hemorrhage, sepsis. Aligning with guidelines for managing anemia, transfusion rates showed a dose-response increase with severity of deficiency. For other outcomes, there were no differences by anemia status except postpartum hemorrhage, which was significantly higher for severe cases.

Uniquely, the study also assessed impacts on hypertensive disorders of pregnancy. Using blood pressure measurements standardized for preeclampsia research, a U-shaped relationship emerged. Compared to no anemia, mild and moderate deficiency associated with 24-29% lower pregnancy hypertension and preeclampsia risks, despite similar gestational ages. Women with hemoglobin $\geq 11\text{g/dL}$ appeared most vulnerable. Posited explanations include elevated blood viscosity or oxidative stress among non-anemic women. However, further interrogation in cohorts encompassing the full hemoglobin range is warranted to confirm this pattern.

6. CONCLUSION

Anemia in early pregnancy remains highly prevalent globally, especially in LMICs like India. Iron deficiency drives most cases, but increasing prevalence of overweight/obesity may signal a double burden of malnutrition. While evidence supports increased risks of certain morbidities with worsening deficiency, associations with sepsis, hemorrhage and preeclampsia are less consistent. Recent findings of lowered hypertension/preeclampsia odds among mildly anemic women challenge conventional thinking on dose-response harms. Further research should confirm whether treating non-anemic pregnant women poses any cardiovascular risks and also determine if supplemental iron beyond routine needs is advisable. Ensuring adequacy of intake continues as the priority for protecting maternal and infant health.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Stevens GA, et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: A systematic analysis of population-representative data. *The Lancet Global Health*. 2013;1:e16-e25.
2. Balarajan Y, et al. Anaemia in low-income and middle-income countries. *The Lancet*. 2011;378:2123-35.
3. Pavord S, et al. UK guidelines on the management of iron deficiency in pregnancy. *Br J Haematol*. 2020;188:819-30.
4. Bah A, et al. Molecular forms of iron and risk of preeclampsia: A systematic review and meta-analysis. *Pregnancy Hypertens*. 2018;14:122-8.
5. Rahman MM, et al. Maternal anemia and risk of adverse birth and health outcomes in low- and middle-income countries: Systematic review and meta-analysis. *Am J Clin Nutr*. 2016;103:495-504.
6. Daru J, et al. Risk of maternal mortality in women with severe anaemia during pregnancy and post partum: Multilevel analysis. *Lancet Glob Health*. 2018;6:e548-e554.
7. Rukuni R, Knight M, Murphy MF, Roberts D, Stanworth SJ. Screening for iron deficiency and iron deficiency anaemia in pregnancy: A structured review and gap analysis against UK national screening criteria. *BMC Pregnancy Childbirth*. 2015;15:269.
8. Jung J, et al. Effects of hemoglobin levels during pregnancy on adverse maternal and infant outcomes: A systematic review and meta-analysis. *Ann NY Acad Sci*. 2019;1450:69-82.
9. Bone JN, et al. Anemia and adverse outcomes in pregnancy: Subgroup analysis of the CLIP cluster-randomized trial in India. *BMC Pregnancy Childbirth*. 2022; 22:407.
10. Karami M, Chalesghar M, Salari N, Akbari H, Mohammadi M. Global prevalence of anemia in pregnant women: A comprehensive systematic review and meta-analysis. *Maternal and Child Health Journal*. 2022;26:1473–14871.
11. Vinogradova MA, Fedorova TA, Strelnikova EV, Rogachevsky O, Shmakov RG, Polushkina ES. Anemia during the

- pregnancy: The management and outcomes depending on the etiology. *Blood*. 2014;124(21):4830.
DOI: 10.1182/blood.V124.21.4830.4830
12. de Sá SA, Willner E, Pereira TA, de Souza VR, Boaventura GT, de Azeredo VB. Anemia in pregnancy: Impact on weight and in the development of anemia in newborn. *Nutricion hospitalaria*. 2015;32(5):2071-9.
 13. Maka SS, Tondare SB, Tondare MB. Study of impact of anemia on pregnancy. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2017 Nov 1;6(11):4847-51.
 14. Achebe MM, Gafter-Gvili A. How I treat anemia in pregnancy: Iron, cobalamin, and folate. *Blood*. 2017 Feb 23;129(8):940-949.
DOI: 10.1182/blood-2016-08-672246
PMID: 28115368.
 15. Bone JN, et al. Anemia and adverse outcomes in pregnancy: Subgroup analysis of the CLIP cluster-randomized trial in India. *BMC Pregnancy Childbirth*. 2022;22:407.
 16. Rahmati S, et al. Maternal Anemia during pregnancy and infant low birth weight: A systematic review and Meta-analysis. *Int J Reprod Biomed*. 2017;15:125-34.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113535>