



Genetic Variability Studies in the Interspecific Derivatives (*S. indicum* x *S. mulayanum*) of Sesame

**Lakshmi Sheethal H ^a, K T Ramya ^{b*}, Praduman Yadav ^b,
K. Alivelu ^b and Sameer Kumar C.V ^a**

^a Department of Genetics and Plant Breeding, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad (Telangana), India.

^b Division of Genetics and Plant Breeding, ICAR-Indian Institute of Oilseeds Research, Rajendranagar, Hyderabad (Telangana), India.

Authors' contributions

This work was carried out in collaboration among all authors. Author KTR did the conceptualization of the study and execution of the experiments. Author LSH did the conducting experiment, raising the material, data recording. Author PY did the biochemical analysis. Author KA did the statistical analysis and data interpretation. Author SKCV did the overall guidance for the study. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i82726>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120871>

Original Research Article

Received: 26/05/2024

Accepted: 28/07/2024

Published: 30/07/2024

ABSTRACT

The present study aims to assess genetic variability for agronomic characters in the interspecific population developed by hybridization between *Sesamum indicum* (Swetha til) x *Sesamum mulayanum* (IC-43144-1). Experimental materials were raised during late Kharif-2023 at ICAR-Indian Institute of Oilseeds Research, Hyderabad. A total of 80 F₆ interspecific derivatives were

*Corresponding author: E-mail: ramya.kt@icar.gov.in;

evaluated along with four check varieties namely GT-10, Swetha Til, IC-43144-1 (*S. mulayanum*), CUMS-17 in Augmented Randomised Complete Block Design (ARCBDB). Analysis of variance (ANOVA) revealed, significant variability in the interspecific derivatives for all the traits studied viz., days for emergence, days for flowering initiation, days for flower session, days for maturity, plant height, number of primary branches per plant, total number of capsules per plant, capsule length, test weight, harvest index and seed yield per plant. The magnitude of phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) values were higher for the traits viz., total number of capsules per plant, harvest index, seed yield per plant, hence improvement through selection could be possible. Phenological traits such as days for emergence, plant height, number of primary branches per plant, total number of capsules per plant, harvest index and seed yield per plant exhibited a high heritability along with a high genetic advance as per cent of mean, indicating additive gene action in the expression of these traits. Simple phenotypic selection may be effective for improving these characters. So, the selection based on these traits should be prioritized in the selection process of interspecific hybrids of sesame breeding programme.

Keywords: Augmented RCBD; heritability; interspecific derivatives; PCV; GCV; (*S. indicum* x *S. mulayanum*).

1. INTRODUCTION

“Sesame (*Sesamum indicum* L.) is a diploid plant with $2n=2X=26$ chromosomes. It is a self-pollinating crop that belongs to the Pedaliaceae family. It is commonly referred to as benni seed, simsim, til, and gingelly, sesame is one of the ancient oilseed crops believed to have originated in East Africa and India” [1]. “Sesame is regarded as one of the oldest oilseed crops, is widely cultivated in Asia and Africa. Known as the “Queen of Oilseeds,” it is valued for its high-quality oil content. The genus *Sesamum* consists of 30 “accepted” species and 17 species with “unresolved” status” [2]. “Asia has a rich diversity of cultivated sesame while Africa is prosperous in wild relatives” [3]. “Seeds are not only valuable for their nutritional and therapeutic benefits but also provide a significant source of oil (44–57%), proteins (18–25%), carbohydrates (13.5%), and ash (5%)” [4].

“Although genetic variation persists for agronomically important traits in sesame, production levels remain notably low in India. Traditional sesame landraces as well as related wild species are an important source of genetic diversity for breeders and form the backbone of agricultural production. Utilization of wild species of *Sesamum* to create genetic variation is practiced by several workers. Pre breeding is a critical step in crop improvement to bring the wild alleles into cultivated genotypes. Wild relatives of sesame are important reservoir of useful genes and need to be exploited for sesame improvement. These wild species exhibit crossability with the cultivated gene pool to

varying extents and can be utilized for transferring the desirable traits using conventional breeding approaches assisted with modern techniques” Singh et al. [5]. Biswas and Mitra [6] were successful in crossing *S. mulayanum*, *S. laciniatum* with *S. indicum*. Seed dormancy was absent in the F_1 hybrids of *S. indicum* x *S. mulayanum* and the inheritance of flower colour followed 9:3:3:1 ratio in F_2 population Tanesaka et al. [7]. Pre-fertilization barriers between *S. indicum* and *S. mulayanum* were absent Kulkarni et al. [8]. Development of interspecific populations and analysis of genetic variation for quality traits like lignans in sesame helps in developing genotypes with superior oil quality rich in lignans Pathak et al. [9]. Pre-breeding and understanding genetic variability within germplasm is essential for selecting and breeding high-yielding, high-quality cultivars to enhance production. The genetic variability present in the available germplasm may be utilized either for direct selection or for hybridization programme which involves a choice of potential parents that can produce progeny population out yielding the parents in a set of desirable characters i.e., characters related to yield. It is imperative to investigate variability in quantitative traits, considering genetic parameters such as genotypic and phenotypic variances, broad-sense heritability, and genetic advance.

The present study is aimed to understand the extent of genetic variation for agronomic characters in the population developed through interspecific hybridization of sesame (*S. indicum* and *S. mulayanum*).

2. MATERIALS AND METHODS

The present investigation was carried out at ICAR - Indian Institute of Oilseeds Research Rajendranagar, Hyderabad during late *kharif*, 2023. The experiment was laid out in augmented block design with 84 germplasm accessions including four check varieties namely GT-10, Swetha Til, IC-43144-1 (*S. mulayanum*), and CUMS-17. The genotypes were sown by following a spacing of 45 cm between the rows and 15 cm between the plants. The recommended agronomical practices and plant protection measures were adopted for raising a good crop. The observations were recorded on five randomly taken plants for eleven quantitative traits *viz.*, days for emergence, days for flowering initiation, days for flower session, days for maturity, plant height, number of primary branches per plant, total number of capsules per plant, capsule length, test weight, harvest index and seed yield per plant. Analysis of variance (ANOVA) was calculated for eleven quantitative traits by using the formula given by Federer et al. [10]. The genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) were estimated by using the formula given by Burton and de Vane [11]. Heritability in broad sense (h^2_b) was estimated according to the formula suggested by Johnson et al. [12] and Hanson et al. [13]. Genetic advance in absolute unit (GA) and percent of the mean (GAM), assuming selection of superior 5% of the genotypes was estimated in accordance with the methods illustrated by Johnson et al. [12]. All the statistical analysis were done in R program version 0.1.7.9000 using augmented RCBD package developed by Aravind et al. [14].

3. RESULTS AND DISCUSSION

The analysis of variance for ARCBD experimental design is presented in (Table 1a and 1b). In the present study, analysis of variance showed that significant mean sum of squares for all traits for different sources of variation. The source of variation for entries (excluding blocks) were found to be significant for all the traits analysed *viz.*, days for emergence, days for flowering initiation, days for flower session, days for maturity, plant height, number of primary branches per plant, total number of capsules per plant, capsule length, test weight, harvest index and seed yield per plant. This indicated the presence of sufficient amount of genetic variability among the genotypes for all the traits. Similarly, a wide

range of variation for the majority of the traits was reported by Kumar et al. [15] and Kumar et al. [16] in sesame.

The critical difference (CD) (Table 2) for any two test treatments of the different blocks showed marginally higher values than the same two test treatments for the same blocks.

The highest value of coefficient of variation (C.V) was found in case of seed yield per plant (13.19), followed by number of primary branches per plant (9.76), total number of capsules per plant (8.07), capsule length (7.44), harvest index (6.15), Plant height (5.22), while, days for maturity (3.53), test weight (2.72) and days for flower session (1.90) had lower C.V values.

Descriptive statistics such as mean, range, standard deviation, standard error, coefficient of variation, skewness and kurtosis for the quantitative traits studied are presented in (Table 2). Significant coefficient of variation for the traits indicated the variability present in the interspecific derivatives. The range of variation for days for emergence is 4 to 6 days with the mean value of 4, days for flowering initiation ranges from 33 to 46 days with the mean value of 39, days for flower session ranges from 62 to 85 days with the mean value of 75 days, days for maturity ranges from 83 to 110 days with the mean value of 96 days, Likewise, the characters plant height ranged from 90.79 to 194.87 cm with the mean value of 132.12 cm, number of primary branches per plant ranged from 3 to 10 with the mean value of 7, total number of capsules per plant ranged from 16 to 124 with the mean value 58, capsule length ranged from 1.80 to 3.49 cm with the mean value of 2.63, seed yield per plant ranged from 1.56 to 10.19 g with the mean value of 5.48 g, test weight ranges from 2.39 to 3.45 g with the mean value 3.0 g, and harvest index ranges from 6.31 to 35.83 % with the mean value 17.33 % also showed adequate variation in the interspecific population. Maximum variation for traits like total number of capsules per plant, plant height and lowest for days for emergence, capsule length, test weight was observed. Vamshi et al. [17] reported low variation for in capsule length and test weight among the advanced breeding lines derived from *S.indicum* crosses.

Frequency distribution of different quantitative traits was presented in Fig. 1. The traits like plant height, total number of capsules per plant, harvest index showed positive skewness. It

indicates that more proportion of genotypes are found at lower end of distribution. Selection of these genotypes will improve the positively skewed traits. Negative skewness was observed for days for maturity and test weight. Distribution of DFI, DFS, CL, and SYP exhibited mesokurtic distribution, indicating uniform or normal distribution for these traits among the interspecific derivatives, while test weight showed negative kurtosis indicating, tailed towards lower values than the mean and harvest index showed positive kurtosis which was tailed towards higher value than the mean. This results are in agreement with the distribution results of advance breeding lines of *S. indicum* by Vamshi et al. [17].

(Table 3) estimation of variance, coefficients of variability, heritability and genetic advance as per cent of mean (GAM).

The value of Phenotypic Coefficient of Variation (PCV) ranged from 4.74 for days for flowering initiation to 37.7 for seed yield per plant, while Genotypic Coefficient of variation (GCV) varied from 3.95 for days to maturity to 36.67 for harvest index. Higher magnitude of both PCV and GCV was recorded for total number of capsules per plant (37.5) and (36.5), seed yield per plant (37.7) and (35.36) and harvest index (37.26) and (36.67). Similar results of high PCV and GCV for total number of capsules per plant, seed yield per plant were reported by Roy et al. [18], Srikanth and Ghodke [19], Thouseem et al. [20] and harvest index by Takele and Abera [21]. Moderate estimates were recorded for days for emergence (15.26), plant height (15.36) and (14.46) and number of primary branches per plant (16.93) and (13.83). Similar results had been reported for plant height by Durodola et al. [22]. Further Low estimates were recorded for days for flowering initiation (6.12) days for maturity (5.27) and (3.95) and days for flower session (4.74), and (4.35) and test weight (7.6), and (7.09). Similar results were reported by Ranjithkumar et al. [23] and Manepalli [24] for days to maturity and days for flowering initiation. Zeinalzadeh [25] reported for test weight. The trait capsule length recorded moderate PCV (10.68) coupled with low GCV (7.71). These results are in agreement with those of Abate and Mekbib [26] and Kant et al. [27]. In the present study, values of PCV were higher for all traits than corresponding GCV but a very little difference between PCV and GCV were noticed for all the traits, indicating that these were less

affected by environment and selection could be effective for further improvement of these traits.

High heritability (broad sense) was observed for the traits viz., days for emergence (100), days for flowering initiation (100), days for flower session (84.2), plant height (88.56), number of primary branches per plant (66.68), total number of capsules per plant (94.74%), seed yield per plant (87.95), test weight (87.04), harvest index (96.83). These findings are in agreement with Gayathri [28]. Remaining traits such as days for maturity (56.13) and capsule length (52.11) exhibited moderate estimates of heritability (broad sense) reported by Abate and Mekbib [26]. This suggested that the phenotypes were the true representative of their genotypes for these characters and selection based on phenotypic value could be reliable.

The high genetic advance (GA) was recorded in traits viz., plant height (37.08), total number of capsules per plant (43.04), while capsule length (0.3), test weight (0.41) were recorded low genetic advance (GA). Genetic advance as per cent of mean (GAM) was observed high for days for emergence (31.48), plant height (28.07), number of primary branches per plant (23.29), total number of capsules per plant (73.3), seed yield per plant (68.41) and harvest index (74.44). These findings are similar with Sahu et al. [29] and Vamshi et al. [17] for plant height, number of primary branches per plant and total number of capsules per plant. Moderate GAM was observed for days for flowering initiation (12.63), capsule length (11.48) and test weight (13.66). Low GAM was observed for days for flowering session (8.24) and days for maturity (6.11) respectively. Pavani et al. [30] also reported low GAM for days for maturity.

Heritability and genetic advance, when considered together would be more reliable and useful in predicting the resultant effects of selection Johnson et al. [12]. High heritability coupled with high genetic advance as percent of mean was observed for six traits viz., days for emergence, plant height, number of primary branches per plant, total number of capsules per plant, seed yield per plant, harvest index indicating that these traits which may be attributed to the preponderance of additive gene action and possess high selective value and thus, selection pressure could profitably be applied on these traits for their rationale improvement. These results were in agreement with those of Divya et al. [31], Khuntia et al. [32],

Table 1a. Analysis of variance for treatment adjustment of eleven agronomic traits for different source of variation

ANOVA - treatment adjusted												
Source of variation	df	DE	DFI	DFS	DM	PH	NPB	TCP	CL	TW	HI	SYP
Block (ignoring Treatments)	3	1.2 **	0.54 **	38.18**	51.36*	347.15**	2.28*	1746.20**	0.25*	0.03**	113.27**	25.27**
Treatment (eliminating Blocks)	83	0.68 **	8.96**	21.96**	45.05*	392.93**	1.6*	697.2**	0.11*	0.07**	72.33**	3.62**
Treatment: Check	3	4 **	91.67**	200**	360.17**	235.18*	8.25**	3502.06**	1.04**	0.48**	433.7**	11.97**
Treatment: Test and Test vs. Check	80	0.55 **	5.86 **	15.29**	33.23*	398.85**	1.35*	592.01**	0.08 ns	0.06**	58.78**	3.31**
Residuals	9	0	0	2	11.44	47.14	0.47	25.56	0.04	0.01	1.32	0.51

Significance levels: ns(non-significant) $P > 0.05$; * $P \leq 0.05$; ** $P \leq 0.01$, d.f. degrees of freedom,

DE-Days for emergence, DFI-Days for flowering initiation, DFS-Days for flower session, DM-Days for maturity, PH-Plant height (cm), NPB-Number of primary branches per plant, TCP-Total number of capsules per plant, CL-Capsule length (cm), TW-Test weight (g), HI- Harvest index (%), SYP-Seed yield per plant (g)

Table 1b. Analysis of variance for block adjustment of eleven agronomic traits for different source of variation

ANOVA - block adjusted												
Source	df	DE	DFI	DFS	DM	PH	NPB	TCP	CL	TW	HI	SYP
Treatment (ignoring Blocks)	83	0.72**	8.98**	23.33**	46.76*	405.14 **	1.65 *	756.94 **	0.12 *	0.07 **	76 **	4.52 **
Treatment: Check	3	4**	91.67**	200 **	360.17**	235.18 *	8.25 **	3502.06 **	1.04 **	0.48 **	433.76 **	11.97 **
Treatment: Test	79	0.57**	5.85**	12.66 **	26.09 ns	411.93 **	1.42 *	486.14 **	0.08 ns	0.05 **	41.67**	4.27 **
Treatment:Test vs. Check	1	2.85**	8.27**	336.67**	740.03**	378.08 *	0.13 ns	13914.8 **	0.28 *	0.37 **	1714.5 **	1.66 ns
Block (eliminating Treatments)	3	0 ns	0 ns	0.33 ns	3.83 ns	9.56 ns	0.92 ns	93.23 ns	0.1 ns	0.01 ns	11.71 **	0.61 ns
Residuals	9	0	0	2	11.44	47.14	0.47	25.56	0.04	0.01	1.32	0.51

Significance levels: ns(non-significant) $P > 0.05$; * $P \leq 0.05$; ** $P \leq 0.01$, d.f. degrees of freedom,

DE-Days for emergence, DFI-Days for flowering initiation, DFS-Days for flower session, DM-Days for maturity, PH-Plant height (cm), NPB-Number of primary branches per plant, TCP-Total number of capsules per plant, CL-Capsule length (cm), TW-Test weight (g), HI- Harvest index (%), SYP-Seed yield per plant (g)

Table 2. Critical difference and Coefficient of Variation

Comparison	Critical difference										
	DE	DFI	DFS	DM	PH	NPB	TCP	CL	TW	HI	SYP
Control Treatment Means	0.00	0.00	2.26	5.41	10.98	1.1	8.09	0.31	0.13	1.84	1.15
Two Test Treatments (Same Block)	0.00	0.00	4.52	10.82	21.96	2.2	16.17	0.62	0.26	3.67	2.29
Two Test Treatments (Different Blocks)	0.00	0.00	5.06	12.1	24.56	2.46	18.08	0.69	0.29	4.11	2.57
A Test Treatment and a Control Treatment	0.00	0.00	4	9.57	19.41	1.94	14.3	0.55	0.23	3.25	2.03
Descriptive statistics											
Mean	4.94	39.50	75.04	96.85	132.12	7.03	58.80	2.63	3.00	17.33	5.48
Std. Error	0.08	0.28	0.42	0.59	2.20	0.14	2.74	0.04	0.03	0.78	0.24
Std. Deviation	0.77	2.53	3.83	5.42	20.16	1.28	25.10	0.36	0.24	7.13	2.23
Min	4.00	33.00	62.25	83.00	90.79	3.63	16.11	1.80	2.39	6.31	1.56
Max	6.00	46.00	85.75	110.00	194.87	10.62	124.31	3.49	3.45	35.83	10.19
CV	0.1	0.06	1.90	3.53	5.22	9.76	8.07	7.44	2.72	6.15	13.19
Skewness	0.10 ^{ns}	0.00 ^{ns}	0.36 ^{ns}	-0.77**	0.58*	0.05 ^{ns}	0.59*	0.09 ^{ns}	-0.53*	1.03**	0.28 ^{ns}
Kurtosis	1.73**	3.30 ^{ns}	4.18*	3.86 ^{ns}	3.41 ^{ns}	2.87 ^{ns}	2.70 ^{ns}	2.62 ^{ns}	2.89 ^{ns}	3.63 ^{ns}	2.32 ^{ns}



CD-Critical difference, CV-Coefficient variation, DE-Days for emergence, DFI-Days for flowering initiation, DFS-Days for flower session, DM-Days for maturity, PH-Plant height (cm), NPB-Number of primary branches per plant, TCP-Total number of capsules per plant, CL-Capsule length (cm), TW-Test weight (g), HI- Harvest index (%), SYP-Seed yield per plant (g)

Table 3. Estimation of variance, coefficients of variability, heritability and genetic advance as per cent of mean (GAM) for eleven agronomic traits

Trait	Mean	PV	GV	EV	GCV		PCV		ECV	hBS		GA	GAM	
					Values	Category	Values	Category		Values	Category		Values	Category
DE	4.94	0.57	0.57	0	15.26	Medium	15.26	Medium	0	100	High	1.56	31.48	High
DFI	39.5	5.85	5.85	0	6.12	Low	6.12	Low	0	100	High	4.99	12.63	Medium
DFS	75.04	12.66	10.66	2	4.35	Low	4.74	Low	1.88	84.2	High	6.18	8.24	Low
DM	96.85	26.09	14.64	11.44	3.95	Low	5.27	Low	3.49	56.13	Medium	5.91	6.11	Low
PH	132.12	411.93	364.8	47.14	14.46	Medium	15.36	Medium	5.2	88.56	High	37.08	28.07	High
NPB	7.03	1.42	0.94	0.47	13.83	Medium	16.93	Medium	9.78	66.68	High	1.64	23.29	High
TCP	58.8	486.14	460.58	25.56	36.5	High	37.5	High	8.6	94.74	High	43.09	73.3	High
CL	2.63	0.08	0.04	0.04	7.71	Low	10.68	Medium	7.39	52.11	Medium	0.3	11.48	Medium
TW	3	0.05	0.05	0.01	7.09	Low	7.6	Low	2.74	87.04	High	0.41	13.66	Medium
HI	17.33	41.67	40.35	1.32	36.67	High	37.26	High	6.63	96.83	High	12.9	74.43	High
SYP	5.48	4.27	3.75	0.51	35.36	High	37.7	High	13.09	87.95	High	3.75	68.41	High



PV-Phenotypic Variance, GV-Genotypic Variance, EV-Environmental Variance, GCV-Genotypic Coefficient Variation, PCV-Phenotypic Coefficient Variation, hBS -Heritability in a Broad Sense, GA-Genetic Advance and GAM-Genetic Advance Mean

DE-Days for emergence, DFI-Days for flowering initiation, DFS-Days for flower session, DM-Days for maturity, PH-Plant height (cm), NPB-Number of primary branches per plant, TCP-Total number of capsules per plant, CL-Capsule length (cm), TW-Test weight (g), HI- Harvest index (%), SYP-Seed yield per plant (g).




Days for emergence

Days for flower initiation




Days for flower session


Days for maturity




Plant height



Total number of capsules per plant



Number of primary branches per plant



Capsule Length




Fig. 1. Estimation of frequency distribution of quantitative traits in sesame

Thouseem et al. [20] for plant height, number of primary branches per plant, total number of capsules per plant, seed yield per plant and harvest index by Hassen et al [33] and Mahla et al [34].

PCV, GCV, heritability in broad sense and Genetic advance as per cent of mean (GAM) should be considered together while assessing the effect of selection than single parameter alone. Mohanty et al. [35] reported high PCV, GCV, heritability and genetic advance as per cent of mean for harvest index and seed yield per plant as observed in the present investigation.

4. CONCLUSION

Pre-breeding is a very important step in crop improvement which involves inflow of wild alleles into the cultivated species. The present study involving interspecific derivatives developed from cultivated *Sesamum indicum* and wild species *Sesamum mulayanum* gives an insight to the extent of genetic diversity created in the sesame crop using wild species. This study aims to identify and characterize variations in agronomic traits related to seed yield. The present study, strongly indicates, significant variability among the interspecific derivatives developed using wild species *S. mulayanum*, for all traits studied. The results of this study can be used by the sesame researchers for developing improved high yielding sesame genotypes. High heritability along with high genetic advance as percent of mean in the traits viz., days for emergence, plant height, number of primary branches per plant, total number of capsules per plant, seed yield per plant, harvest index were with the predominance of additive type of gene action predicting genetic gain under selection than heritability estimates alone and suggesting that these characters can be improved by simple selection among the interspecific derivatives.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

The authors would like to thank ICAR-Indian Institute of Oilseeds Research, Hyderabad for

financial aid in conducting the research and for providing the germplasm, and other resources and the opportunity to carry out this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bedigian D, and Harlan JR. Evidence for cultivation of sesame in the ancient world. *Economic botany*. 1986;40:137-154.
2. Kalwij JM. Review of The Plant List, a working list of all plant species. *Journal of Vegetation Science*. 2012;23(5):998-1002.
3. Sharma SN, Kumar V, Sharma C, Shukla N, Gupta A, Tripathi A, and Paroha S. Characterization and analysis of genetic diversity in Indian Sesame (*Sesamum Indicum* L.) genotypes. *Indian Journal of Genetics and Plant Breeding*. 2014;74(03):344-352.
4. Myint D, Gilani SA, Kawase M, Watanabe KN. Sustainable sesame (*Sesamum indicum* L.) production through improved technology: An overview of production, challenges, and opportunities in Myanmar. *Sustainability*. 2020;12(9):3515.
5. Singh V, Kumar S, Singh A, Bhaduri, NP, Bhat KV, Lakhanpaul S. Unlocking the potential of genetic resources for improvement of sesame (*Sesamum indicum* L.): The current scenario. *Gene Pool Diversity and Crop Improvement*. 2016, Volume 1, 447-479. DOI: 10.1007/978-3-319-27096-8_15
6. BISWAS AK, MITRA AK. Interspecific hybridization in three species of Sesamum. *Indian Journal of Genetics and Plant Breeding*. 1990;50(04):307-309.
7. Tanesaka E, Umeda E, Yamamoto M, Masuda K, Yamada K, Yoshida M. Inheritance mode of seed dormancy in the hybrid progeny of sesame, *Sesamum indicum*, and its wild relative, *Sesamum mulayanum* Nair. *Weed biology and management*. 2012;12(2):91-97.
8. Kulkarni VV, Ranganatha CN, Shankergoud I. Interspecific crossing barriers in sesame (*Sesamum indicum* L.). *Int J Curr Microbiol App Sci*. 2017;6(10):4894-4900.
9. Pathak N, Bhaduri A, Bhat KV, Rai AK. Tracking sesamin synthase gene expression through seed maturity in wild

- and cultivated sesame species—a domestication footprint. *Plant biology*. 2015;17(5):1039-1046.
10. Federer WT, Raghava Rao D. On Augumented Designs. *Biometrics*. 1975;31:29-35.
 11. Burton GW, and de Vane EH. Estimating heritability in Tall Fescue (*Festuca arundinacea*) from replicated clonal material. *Agronomy Journal*. 1953;45:481-487.
 12. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybean. *Agronomy Journal*. 1955;47:314-318.
 13. Hanson CH, Robinson HF, Comstock RE. Biometrical studies of yield in segregating populations of Korean lespedeza 1. *Agronomy Journal*. 1956;48(6):268-272.
 14. Aravind J, Mukesh Sankar S, Wankhede DP, and Kaur V. augmented RCBd: Analysis of augmented randomised complete block, designs. R package version 0.1.7.9000; 2018.
 15. Kumar H, Singh P, Banga S. Assessment of phenotypic divergence in a collection of sesame (*Sesamum indicum* L.) germplasm. *Crop Improvement*. 2010;37(2):140-148.
 16. Kumar N, Chander S, Punia R, and Kumar D. Correlation studies and path analysis in sesame (*Sesamum indicum* L.). *International Journal of Plant & Soil Science*. 2024;36(7):1043-1047.
 17. Vamshi P, Ramya KT, Prasad MS, Ramana JV, Kumhar SR. Genetic variability studies for yield and yield attributes in breeding lines of sesame (*Sesamum indicum* L.). *Journal of Oilseeds Research*. 2021;38(3):240-243.
 18. Roy B, Pal AK, Basu AK. The estimation of genetic variability and genetic divergence of some advance lines of sesame based on morphological traits. *Plant Science Today*. 2022;9(2):281-287.
 19. Srikanth K, Ghodke MK. Genetic diversity studies in sesame (*Sesamum indicum* L.) genotypes. *The Pharma Innovation Journal*. 2022;11(3):1486-1491.
 20. Thouseem N, Arya K, Gayathri G. Genetic variability, heritability and genetic advance for seed yield and yield associated traits in White seeded sesame (*Sesamum indicum* L.). *The Pharma Innovation Journal*. 2022;11(6):385-388.
 21. Takele F, and Abera G. Variability Study in Ethiopian Sesame (*Sesamum indicum* L.) Genotypes at Western Oromia. *International Journal of Precision Farming*. 2023;1(1): 1-7.
 22. Durodola FA, Azeez MA, Adubi AO. Morpho-agronomic variability, heritability and genetic advance studies in sesame (*sesamum indicum* L.) In nigeria. *Bangladesh Journal of Botany*. 2024;53(1):75-81.
 23. Ranjithkumar G, Bisen R, Kumar V. Genetic variability studies for yield and its attributing traits in dark brown sesame (*Sesamum indicum* L.). *Int. J. Plant Soil Sci*. 2022;34(8):43-49.
 24. Manepalli SB. Genetic studies for yield and its related traits in sesame (*Sesamum indicum* L.) Genotypes; 2022.
 25. Zeinalzadeh-Tabrizi H. Assessment of heritability and genetic efficiency in advanced sesame inbred lines. *Journal of Animal & Plant Sciences*. 2024;34(2):515-523.
 26. Abate M, Mekbib F. Assessment of genetic variability and character association in Ethiopian low altitude sesame (*Sesamum indicum* L.) genotypes. *Journal of Advanced Studies in Agricultural, Biological and Environmental Sciences*. 2015;2(3):55-66.
 27. Kant C, Mishra SP, Mani N, Kumar N. Assessment of genetic variability & some biochemical traits in Sesame (*Sesamum Indicum* L.) genotypes. *Journal of Pharmacognosy and Phytochemistry*. 2021;10(5):252-256.
 28. Gayathri G. Heterosis breeding in sesame (*Sesamum indicum* L.) Doctoral dissertation, Department of Plant Breeding and Genetics, College of Horticulture, Vellanikkara; 2011.
 29. Sahu D, Mishra TK, Pradhan B. Genetic variability, heritability and genetic advance in different genotypes of sesame (*Sesamum indicum* L.). *The Pharma Innovation Journal*. 2022;11(11):1799-1801.
 30. Pavani K, Lal Ahamed M, Ramana JV, Sirisha ABM. Studies on genetic variability parameters in sesame (*Sesamum indicum* L.). *International Journal of Chemical Studies*. 2020;8(4):101-104.
 31. Divya K. Studies on genetic diversity in advanced mutant breeding lines of sesame (*Sesamum indicum* L.). PhD thesis. Professor Jayashankar Telangana State Agricultural University, Hyderabad, India. 2018.

32. Khuntia JB, Dash AP, Kar RK, Lenka B, Pradhan H, Samal A, Pattanayak S, Mishra S, Pani B, Satapathy S. Unveiling Genetic Variability and Character associations in Sesame. *Biochemical & Cellular Archives*. 2024;24(1). yield related traits of sesame (*Sesamum Indicum* L.) genotype. *International Journal of Plant & Soil Science* 36. 2024;(2):197-206. Available:<https://doi.org/10.9734/ijpss/2024/v36i24381>
33. Hassen M. Morphological characteristics of sesame (*Sesamum indicum* L.) genotypes via genetic diversity and characters association in Amibara, Ethiopia. *Ecology and Evolutionary Biology*. 2022;7(3):30-45.
34. Mahla NU, PK Jagtap, HR Patel. Genetic variability and association among yield and 35. Mohanty TA, Singh UM, Satish, Singh SK, Kushwaha N, Singh D. Study of genetic variability, heritability and genetic advance in sesame (*Sesamum indicum* L.). *International Journal of Current Microbiology and Applied Sciences*. 2020;9(2):347-356.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/120871>