



Effect of Weed Management on Weed Growth, Performance, Profitability and Economics of Wheat in Transitional Plain of Luni Basin

Arjun Lal Bijarnia^{a*}, P. C. Yadav^b, A. S. Godara^a, N. V. Suresh^a, D. M. Meena^a
and Sohan Lal Boori^a

^a Agricultural Research Station, Keshwana, Jalore, India.

^b Krishi Vigyan Kendra, Keshwana, Jalore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i1230955

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84199>

Original Research Article

Received 20 January 2022

Accepted 31 March 2022

Published 02 April 2022

ABSTRACT

A experiment was conducted at Agricultural Research Station, Keshwana, Jalore under arid climate conditions in *Rabi* season during 2018-19 with the object of effect of herbicide on weed, wheat growth, yield and economics. The data shows that the treatment 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS recorded lower dry matter of weeds at harvest (18.3 g), maximum growth, spike length (8.5 cm), No. of spikelet (17), No. of seeds per spike (54.7), seed yield (40.1 q) and Straw yield (59.1 q) of wheat, and at par with the treatment Metsulfuron 4 g ha⁻¹ 35 DAS + hand hoeing 45 DAS. The treatment 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS also produced the maximum gross return (Rs 117763), net return (Rs 70886) and B:C ratio (2.51).

Keywords: *Wheat; weed management; growth; yield; economics.*

1. INTRODUCTION

India stands in second position next to china in the world with regard to area and production of

wheat. In India wheat is grown on 34.5 million hectares with total production of 108.75 million tonnes with average productivity of 3152 kg ha⁻¹ [1]. In Rajasthan, it is cultivated on 3.09 million

*Corresponding author: E-mail: abijariya2010@gmail.com;

hectares area with total production of 12.02 million tonnes and average productivity of 3885 kg ha⁻¹ [1]. The productivity of the crop in the western Rajasthan is lower than the potential yield. In Jalore district, it is grown on 0.05 million hectare with the total production 0.12 MT and average productivity is 2339 kg ha⁻¹ is very low as compared to state productivity [2].

Productivity of wheat is very low in due to stiff competition from weeds variable climatic conditions, genotypes, seeding time and practices; and other management practices (Kantwa et al.2015). In the wheat crop initial high soil moisture and free space leads to severe infestation of grassy and broad leaves weeds. Globally yield reduction in wheat due to weeds is 13.1% (T.) or more and also reported that selected wheat varieties incurred 60-65 % biomass loss due to weed infestation.

The insect and disease effect on crop is visible, but the damage done by weeds is often noticed. The total loss caused by various pests in agriculture, weeds account for 37%, followed by insect (29%), disease (22%) and others including nematodes, rodents, mites, birds etc. (12%) weed seeds germinate along with crop seeds or many a time before crop and start competing with crops for vital growth resources like solar radiation nutrients and water resulting in serve yield loss. All together total actual economic loss due to weeds in 16 major crops was estimated as Rs 78,591 crore per annum [3]. However, the total economic losses will be much higher, if all the crops, and indirect effects on weeds on human and animal health, loss of biodiversity, nutrient depletion, reduction in grain quality etc. are taken in to consideration.

2. MATERIALS AND METHODS

2.1 Field Experiment Site, Detail and Soil

A field experiment was conducted during rabi season 2018-19 at Agricultural Research Station, Keshwana, Jalore (Agriculture University, Jodhpur, Rajasthan) to study "Effect of weed management practices on weeds, growth, yield and economics of wheat in transitional plain of luni basin of Rajasthan" Randomized Block Design with three replications was used to design the field experiment. The treatments for this experiment was, Weedy check, Weed free, One hand hoeing at 35 DAS, 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS, Metsulfuron 4 g ha⁻¹ 35 DAS, 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS,

Metsulfuron 4 g ha⁻¹ 35 DAS + hand hoeing 45 DAS. Wheat variety RAJ3077 was sown on 6th December 2018. Sowing was done manually by using 100 kg ha⁻¹ seed rate in a row spaced 22.5 cm.

2.2 Analysis of Weeds, plant Growth and Yield

Weed dry matter of each weed species was taken at initial and harvest from two random spots in each plot by counting the number of weeds per quadrat of 0.25 m² and then it was converted in to m⁻².The observations on plant height, spike length, No. of spikelet per spike branch plant⁻¹, were recorded manually on five selected representative plants and No. of seeds per spike was recorded manually on 10 spike from selected five plants. Harvesting was done 13 April 2019. The seed and straw yield were recorded from the net plot area of each treatment. The test weight was recorded by counting of thousand seeds then weights it. Weed control efficiency (WCE) was calculated by using the following formula suggested by Das and expressed in percentage:

$$WCE = \frac{DMC - DMT}{DMC} \times 100$$

Where, DMC is the dry matter of weeds in control (unweeded) plot and DMT is the dry matter of weeds in treated plot.

2.3 Economics and Statistical Analysis

An economic analysis was done to compare the returns of various treatments of herbicide applications. Gross return determined from seed and straw yield and net return was determined by subtracting the costs of production from gross income. The data of experiment in different observations were statistically analyzed in accordance with the analysis of variance techniques as described by Panse and Sukhatme [4]. The critical difference (CD), were calculated at 5 % level of probability. The elucidate the nature and magnitude of treatments effects, summary table along with SEM_± and CD (p=0.05) were prepared.

3. RESULTS AND DISCUSSION

The experiment field was infested with *Chenopodium album*, *Chenopodium murale*, *Rumex dentatus*, *Asphodelus tenuifolius*,

Melilotus indica, *Cyprus rotendus*, *cenchrus Species* and *Fumaria parviflora*. The effect of different management practices on different weeds was significant (Table 1). All the weed control treatments statistically reduced the dry matter of weeds over weedy check at 60 DAS and harvest. The treatment 2, 4 D @ 0.5 kg a.i. ha⁻¹ at 35 DAS+ hand hoeing 45 DAS significantly reduced the density of Broad leaves weed as well as grassy weeds at 60 DAS and harvest. The weed control efficiency shows that 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS and 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS control the weeds 73.09 and 72.08 respectively over weedy check. Regeneration of *R. dentatus* was noticed in 2,4-D at 0.5 kg a.i./ha applied plots and thus increased the dry matter of the weed as compared to metsulfuron treated plots. These findings were in conformity with those reported by Singh and Ali [4] and Pisal et al. [5] Shivran et al. [6] also observed that Regeneration of *R. dentatus* was noticed in 2,4-D applied plots and thus increased the dry matter. The herbicide 2,4 D is used on a wide variety of terrestrial and aquatic broadleaf weeds Anonymous [7]. It has little effect on grasses. It appears to work by causing uncontrolled cell division in vascular tissue. Abnormal increase in cell wall plasticity, biosynthesis of proteins and production of ethylene occurs in plant tissues following exposure, and these processes are responsible for uncontrolled cell division: Anonymous [8].

The metsulfuron 4 g a.i. /ha PoE was next best after 2,4 D in minimising weed biomass. A significant reduction in weed biomass 68.37 and 68.53 % with Metsulfuron 4 g ha⁻¹ + hand hoeing 45 DAS and Metsulfuron 4 g a.i. ha⁻¹ 35 DAS, respectively. Metsulfuron is generally absorbed by leaves and translocated to growing points of the plant where it stops cell division and inhibiting

the photosynthesis resulting into yellowing of plants.

3.1 Effect on Wheat Growth, Yield Attributes and Yields

The plant population and plant heights are important growth parameters influencing yields which are not only genotypic but also on environmental and management practices. The herbicide application of 2, 4-D and Metsulfuron alone and with hand hoeing did not show any significant effect on wheat at harvest stage,

Being the most important economic component, grain yield of a crop reflects the resultant impact of yield attributes as influenced by herbicidal treatments. The maximum plant height (87.33 cm) was observed with weed free treatment, but the spike length, no. of spikelet per spike and No. of seeds per spike maximum observed with the treatment 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS but which was at par with Metsulfuron 4 g ha⁻¹ 35 DAS + hand hoeing 45 DAS. Weed control measures did not show significant effect on test weight of wheat.

The Applied treatment 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS significantly increased the grain and straw yield over one hand hoeing at 35 DAS, 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS and Metsulfuron 4 g ha⁻¹ 35 DAS but was statistically at par with Metsulfuron 4 g ha⁻¹ 35 DAS + hand hoeing 45 DAS as well as weed free check. The treatment 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS and Metsulfuron 4 g ha⁻¹ 35 DAS + hand hoeing 45 DAS was to extent of 76 and 72 % than the weedy check treatment. These results were in close conformity with the finding of Das [9] and Singh et al. [10].

Table 1. Physico-chemical characteristics of the field (0-30 cm)

Soil parameters	Value
pH	8.02
EC (dS m ⁻¹)	0.45
Organic carbon (%)	0.23
Bulk density (Mg m ⁻³)	1.55
Available N (kg ha ⁻¹)	166
Available P ₂ O ₅ (kg ha ⁻¹)	18
Available K ₂ O (kg ha ⁻¹)	280
Available S (mg kg ⁻¹)	13.44
DTPA extractable Zn (mg kg ⁻¹)	0.17
DTPA extractable Fe (mg kg ⁻¹)	2.70
Soil texture	Sandy loam

Table 2. Effect of weed management on weed dry matter (g m⁻²) at 60 DAS and harvest

Sl. No.	Treatments	Dry matter of grassy weeds at 60 DAS	Dry matter of BL weeds at 60 DAS	Dry matter of total weeds at 60 DAS	Dry matter of grassy weeds at harvest	Dry matter of BL weeds at harvest	Dry matter of total weeds at harvest	Weed control efficiency at harvest
1	Weedy check	6.80	26.67	33.47	8.80	56.87	65.67	0
2	Weed free	0.00	0.00	0	0.00	0.00	0	100
3	One hand hoeing at 35 DAS	4.50	19.18	23.68	5.20	36.47	41.67	36.55
4	2,4 D @ 0.5 kg ha ⁻¹ at 35 DAS	2.00	11.40	13.4	2.53	15.14	17.67	73.09
5	Metsulfuron 4 g ha ⁻¹ 35 DAS	2.20	14.03	16.23	2.93	17.74	20.67	68.53
6	2,4 D @ 0.5 kg ha ⁻¹ at 35 DAS+ hand hoeing 45 DAS	1.70	7.60	9.30	2.63	15.70	18.33	72.08
7	Metsulfuron 4 g ha ⁻¹ 35 DAS + hand hoeing 45 DAS	1.75	8.88	10.63	2.85	17.92	20.76	68.37
	SEm±	0.33	0.76	0.93	0.28	2.72	3.457	-
	CD (P=0.05)	1.03	2.37	2.89	0.88	8.48	10.77	-

Table 3. Effect of weed management on growth yield attributes and yield

Sl. No.	Treatments	Plant population (No.of plants/m ²)	Plant height at harvest (cm)	Spike length (cm)	No. of spikelet per spike	No. of seeds per spike	Test wt (g)	Seed yield (q/ha)	Straw yield (q/ha)
1	Weedy check	20.49	70.67	8.17	12.00	37.67	39.32	22.76	34.80
2	Weed free	21.33	87.33	8.80	16.60	53.67	40.63	36.44	55.67
3	One hand hoeing at 35 DAS	20.87	77.33	8.53	14.17	49.50	41.74	33.33	50.60
4	2,4 D @ 0.5 kg ha ⁻¹ at 35 DAS	21.09	80.57	8.43	15.22	53.33	42.17	32.89	49.33
5	Metsulfuron 4 g ha ⁻¹ 35 DAS	20.56	74.83	7.60	14.44	51.00	41.12	27.02	40.53
6	2,4 D @ 0.5 kg ha ⁻¹ at 35 DAS+ hand hoeing 45 DAS	20.79	86.17	8.53	16.89	54.67	41.58	40.09	59.13
7	Metsulfuron 4 g ha ⁻¹ 35 DAS + hand hoeing 45 DAS	20.75	77.13	8.20	15.11	52.00	41.50	39.11	57.33
	SEm±	0.67	3.34	0.40	0.84	1.61	1.51	1.73	2.73
	CD (P=0.05)	NS	10.39	NS	2.62	5.00	NS	5.38	8.49

Table 4. Effect of weed management on economics of wheat

Sl. No.	Treatments	Gross Monitory (Rs/ha.)	Net Monitory (Rs/ha.)	B:C Ratio
1	Weedy check	67472	26345	1.64
2	Weed free	108003	56876	2.11
3	One hand hoeing at 35 DAS	98626	52499	2.14
4	2,4 D @ 0.5 kg ha ⁻¹ at 35 DAS	97023	55146	2.32
5	Metsulfuron 4 g ha ⁻¹ 35 DAS	79709	37882	1.91
6	2,4 D @ 0.5 kg ha ⁻¹ at 35 DAS+ hand hoeing 45 DAS	117763	70886	2.51
7	Metsulfuron 4 g ha ⁻¹ 35 DAS + hand hoeing 45 DAS	114707	67880	2.45

The minimum value of growth, yield attributes and yield was found with weedy check treatment it might be due to competition by weeds for resources, which made the crop plant incompetent to take up more water and nutrients, consequently growth was adversely affected. Poor growth and less uptake of nutrients in weedy check might be due to less photosynthates, then less assimilates to numerous metabolic sink and ultimately poor development of yield components.

3.2 Economics

The maximum net returns and B:C ratio were recorded under application of 2,4 D @ 0.5 kg ha⁻¹ at 35 DAS+ hand hoeing 45 DAS over all the weed control treatments. This might be due to higher grain and straw yield and lower cost to control of weed. These results are in close agreement with results of shivran *et al.* 2020 who reported that highest net return was observed at metsulfuron 4.0 g/ha + 1 hand weeding and 2,4-D 0.5 kg/ha + 1 hand weeding.

4. CONCLUSION

Hand weeding is a common practice in Rajasthan but it is less efficient, labour intensive, costly and often not done on right time. Hand weeding is generally done when weed infestation is quite visible in the field and at this stage weeds have already done the competition to the crop but without hoeing the soil remains compact.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Department of Agriculture site Government of Rajasthan; 2021.
2. Gharde Y, Singh PK, Dubey RP, Gupta PK. Assisment of yield and economic losses in agriculture due to weeds in India. *Crop Protection*. 2018;107:12-18.
3. Singh A, Jain KK, Upadhyaya SD. Effect of weed control methods on weeds and wheat under Eucalyptus tereticornis based agroforestry system. *International Journal Current Microbial and Applied Sciences*. 2018;7(8):2856–2867.
4. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers*; Indian Council of Agricultural Research: New Delhi, India. 1985;87-89.
5. Pisal RR, Gopinath K, Pandey AK. Evaluation of sulfosulfuron and metribuzin for weed control in irrigated wheat. *Indian Journal of Agronomy*. 2013;51(31):135–138.
6. Shivran H, Yadav RS, Singh SP, Godara SL, Bijarniya AL, Samota SR. Tillage and weed management effect on productivity of wheat in North-West Rajasthan. *Indian Journal of Weed Science*. 2020;52 (2):127-131.
7. Anonymous. *Herbicide Handbook*, 8th ed.; Vencil, W.K. Ed.; Weed science society of America: Lawrence, KS. 2002;113-115.
8. Anonymous. Reregistration Eligibility Decision (RED) 2,4-D: EPA738-R-05-002; U.S. Environmental Protection Agency, Office of Prevention, Pesticides and Toxic Substances, Office of Pesticides Programs, U.S. Government Printing Office: Washington, DC; 2005.

9. Das TK. Weed Science: Basics and Application. Jain Brothers Pub, New Delhi, 1st Ed. 2018; 655. effects on succeeding soybean crop grown on vertisol of Rajasthan. Indian Journal of Weed Science. 2004;36(1-2):34–37
10. Singh P, Ali M. Efficacy of metsulfuron-methyl on weeds in wheat and its residual

© 2022 *Bijarnia et al.*; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84199>