

Effect of Hydroethanol Leaf Extract of *Fleurya Aestuans* on Histological and Renal Function Test of an Alloxan Induced Diabetic Wistar Rats

Charles C. Nwafor ^{a*}, Osakwe F. Uchechwukwu ^a, Mfonido G. Umanah ^a,
Ekikereabasi P. Inyang ^a and Peters K. Dell ^b

^a Department of Human Physiology, Faculty of Basic Medical Sciences, College of Medicine, Madonna University, P.M.B 48, Elele, Nigeria.

^b Department of Human Anatomy, Faculty of Basic Medical Sciences, College of Medicine, Madonna University, P.M.B 48, Elele, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87887>

Original Research Article

Received 07 April 2022

Accepted 15 June 2022

Published 23 June 2022

ABSTRACT

Aim: To examine the effect of *Fleurya aestuans* extract (FAE) on histological and renal function test (RFT) in an alloxan-induced type 2 diabetes rat model.

Study Design: The experiment was conducted in the department of Human Physiology, Madonna University, Nigeria, between August and September 2020.

Methodology: Thirty-five (35) male Wistar rats were split into seven (7) groups at random. Group 1 was the control group (CG). Group 2 was the negative control group (NCG). Group 3 was the extract group with the lowest dose (LDEG). Group 4 was the medium dose extract group (MDEG). Group 5 was the extract group with the highest dose (HDEG). Positive control group 1 (PCG1) was made up of rats from group 6. Positive control group 2 (PCG2) was made up of rats from Group 7. Rats were given chloroform anesthesia and sacrificed by cutting the jugular vein and blood was collected into heparinized bottles for renal function test.

Results: *Fleurya aestuans* extract (FAE) at lowest, medium and highest doses given for 30 days resulted in significant reductions in renal electrolyte markers. In diabetic rats treated with FAE, urea and creatinine clearance were likewise declined. Histological experiments were also used to uncover the extract's ameliorative potential structural impacts on renal damage.

Conclusion: We have correctly anticipated that the bioactive elements of FAE may reduce the progression of renopathy using the ALX model. *F. aestuans* may operate as a possible adjuvant for antidiabetic therapy and should be further researched.

Keywords: *Fleurya aestuans*; alloxan; diabetes; renal function test.

1. INTRODUCTION

Diabetic renal dysfunction (DRD) is one of the most common causes of death among diabetics [1]. The thickening of the basement membrane, the proliferation of mesangial cells, fibrosis, and death of podocytes all contribute to morphological and functional problems. In diabetic individuals, renal cell dysfunction causes increased excretion of urine albumin, urea, uric acid, and creatinine, fluid retention, glomerular lesions, and a reduction in glomerular filtration rate (GFR). Furthermore, chronic hyperglycemia and oxidative stress are two important contributors to the development of diabetic complications. Under normal circumstances, the hexokinase enzyme phosphorylates glucose for energy production, however persistent hyperglycemia causes hexokinase saturation. As a result of this saturation, extra glucose enters pathways like the polyol pathway. Chronic hyperglycemia causes oxidative stress by raising the number of reactive oxygen species in the body (ROS). As a result, changes in the metabolic pathways indicated above cause structural and functional abnormalities in nephrons, eventually leading to diabetic renopathy [2,3].

Many diabetic patients choose complementary and alternative therapies in addition to the existing diabetes interventions such as oral hypoglycemic medications and insulin therapy, which has increased the demand for alternative therapies for diabetes control [4].

Fleurya aestuans is a plant that belongs to the Urticaceae family. It is known by a variety of Nigerian names, including *fiyafiya* and *ofuefue* (Yoruba), *bulsum fage* (Hausa), and *ile-nkita* (Igbo) as one of the unique herbal plants prized for its nutritional and therapeutic benefits [5]. Traditional medicine systems employ the leaves of this plant to treat a variety of ailments. Tetrahydroxyflavone, flavonoids (quercetin and isoquercetin, rutin), terpenes (limonin), terpenoids, polyphenols, fatty acids, tannins, sterols, vitamin C, and tocopherols, among others, may be responsible for *Fleurya aestuans*'

medicinal activity [6]. Antidyslipidemic and anti-inflammatory properties are among the pharmacological properties of *Fleurya aestuans* [6,7]. Due to its widespread use in everyday food, *Fleurya aestuans* could be regarded a functional plant. Furthermore, its pharmacological benefits in treating a variety of ailments encourage its use as a functional plant [7].

The current study was designed to explore hydroethanolic extract of *Fleurya aestuans* leaf for attenuation of diabetic renal failure, following the trend of employing safe natural products along with functional plants, their bioactive ingredients, and therapeutic capabilities.

2. MATERIALS AND METHODS

2.1 Preparation of Extract

Fresh leaves of *Fleurya aestuans* (West Indian wood nettle) were collected from the University of Port-botanic Harcourt's farm and scientifically defined by the herbarium of the Department of Plant Sciences, Faculty of Sciences, University of Port-Harcourt, Choba, Nigeria, with the reference number UPH/P/263. The leaves were washed thoroughly and allowed to dry at room temperature. A plastic jar was used to store the dried leaves. The extract was prepared using Al-Attar and Abu Zeid's [8] method with certain changes. Using a manual granulating machine, the dried leaves were ground into fine powder. 350g of plant powder was weighed and dissolved in 400ml of Water-Ethanol combination (25:75) in an Extraction Jar for 72 hours. It was well macerated during this time in order for the solvent to absorb. The filtrate was then separated from the residue by sieving it through a Whatman No. 1 filter paper. After producing a clear filtrate, it was poured onto an evaporating dish and dried on a steam bath at 45 degrees Celsius. The drying process was closely watched until the mixture transformed into a paste. The crude ethanolic extract of *F. aestuans* leaves obtained yielded 75.3g, which was stored at 4°C in a domestic refrigerator.

2.2 Lethal Dose (LD₅₀) of the Extract

Lorke's [9] approach was used to determine the extract's lethality, with slight modifications.

There were two parts to this method:

Phase A:

Nine (9) male rats were employed in this phase. They were divided into three (3) groups, each with three (3) rats. The hydro-ethanolic extract of the leaves of *Fleurya aestuans* was given to each group of rats in doses of 40, 400, and 600 mg/kg, respectively. The animals are kept under constant observation for 24 hours in order to track their behavior and mortality rates.

Phase B:

Four (4) male rats were divided into four (4) groups of one (1) rat each in this phase. The rats were given 800, 1300, 3500, and 4600 mg/kg of the hydro-ethanolic extract of *Fleurya aestuans* leaves, respectively, and were subsequently monitored for a day for signs of acute intoxication and mortality.

The extract's LD₅₀ was estimated using the following formula:

$$LD50 = \sqrt{(D0 \times D100)}$$

Where;

D0 = The highest dose that resulted in no death,

D100 = Lowest dose that resulted in death.

2.3 Induction of Experimental Diabetes

Diabetes was produced in all of the test groups by injecting 150mg/kg body weight of alloxan monohydrate intraperitoneally after an overnight fast [10]. The accucheck glucometer and test strips were used to measure the blood glucose levels (BGL) on a daily basis. Diabetic rats were employed in the investigation because their blood sugar levels were higher than 200 mg/dl.

2.4 Experimental Design

For the investigation, thirty-five (35) male Wistar rats were employed. The rats ranged in weight from 180.4 to 210.9 grams. They were purchased from the Madonna University Animal House in Elele, Nigeria. Prior to the start of the

experiment, the animals were housed in plastic cages and fed commercial rat feed and water ad libitum for a week to acclimate. The animals were split into seven (7) groups, each with five (5) animals.

Group 1 functioned as the control group (CG), receiving unlimited rat food and water.

Group 2 was the negative control group (NCG), receiving ALX only.

Group 3 was the low dose extract group (LDEG), receiving 50 milligrams per kilogram of body weight of *Fleurya aestuans* leaf extract for 30 days.

Group 4 was the medium dosage extract group (MDEG), receiving 75 milligrams per kilogram of body weight of *Fleurya aestuans* leaf extract for 30 days.

Group 5 was the high dose extract group (HDEG), given 200mg/body weight of *Fleurya aestuans* leaf extract for 30 days.

Group 6 was the positive control group 1 (PCG1) rats, given 600 mg of glibenclamide per kilogram of body weight for 30 days.

Group 7 was the positive control group 2 (PCG2) rats, given 100 mg of tetrahydroxyflavone per kilogram of body weight for 30 days.

2.5 Collection of Blood Samples

Rats were given chloroform anesthesia and sacrificed by cutting the jugular vein. Blood was collected into heparinized bottles and centrifuged for 10 minutes at 3000rpm; the supernatant (serum) was collected and stored in the refrigerator for future use.

2.6 Determination of Sodium (Maruna and Trider Method) mmol/L

The current method is based on the reaction of sodium with a selective chromogen, which results in a chromophore whose absorbance fluctuates in direct proportion to the sodium concentration in the test sample.

2.7 Determination of Potassium (Tiets N.W Method) Unit mmol/L

The amount of potassium in a colloidal suspension is determined by using sodium

tetraphenylboron in a specially prepared combination. The sample's turbidity is proportional to the potassium concentration.

2.8 Determination of Chloride: (Levinson S.S Method) Unit mmol/L

The calorimetric displacement of thiocyanate by chloride from mercuric thiocyanate, as well as the subsequent production of red ferric thiocyanate complexes, is measured.

2.9 Determination of Bicarbonate (HCO₃) (Back Titration Method)

Excess standard HCL reacts with serum HCO₃. Back titrates the leftover HCL with standard NaOH using phenol red as an indicator.

2.10 Determination of Urea (Urease-Berthelot Method) mmol/L

In the presence of urease, urea in serum is hydrolyzed to ammonia. Berthelot's reaction is then used to measure ammonia photometrically.

2.11 Determination of Creatinine (Direct End-Point Method) umol/L

In an alkaline solution, creatinine interacts with picric acid to generate a colored complex. The amount of complex produced is proportional to the concentration of creatinine.

2.12 Histological Examination

Following the procedure indicated by Bancroft and Gamble [11]; the tissues were fixed using 10% formalin for 7 days.

The tissues were dehydrated using ascending grades of alcohol 30% 50% 70% (3-12hr) 90% (3-12hrs) Absolute (3-12 hrs) Absolute 2 (3-12hrs) Absolute 3 (3-12 hrs).

The tissues were cleared (de-alcoholisation) using xylene (15- 30 mins).

The tissue were impregnated using paraffin wax for 2-3hrs

The tissues were embattled with molten paraffin wax in a plastic ice- tray the tray was

immediately transfer to cold water to harden the wax.

The block are attached a holder

The tissue were cut using base-sledge microtome

The section tissues are floated in warm-water and pick-up with a slide.

The slides were incubated to remove the wax end fix the tissues on the slide

The slides were stain with Haematoxylin and Eosin.

The slides were mounted using Canada balsam or DPX

The slides were examined using x 40 obj. lens

2.13 Statistical Analysis


All data was expressed as mean + S.E.M. and statistical analysis was performed. A one-way analysis of variance was used to compare the two groups (ANOVA). When P 0.05, values were considered significant.

3. RESULTS

The extract at lowest (40 mg/kg) and highest lethal dose (4600 mg/kg) did not produce any form of lethality in the study animals. Fig. 1 demonstrated a significant increase in renal function parameters in the negative control group when compared to the control group. The extract at 200mg/kg caused a significant decrease ($p < 0.05$) in plasma sodium ions and creatinin in contrast to the negative control group. When compared to the negative control group, glibenclamide at 600 mg/kg induced a significant decrease ($p < 0.05$) in plasma potassium, sodium, and chlorine ions. When compared to the negative control group, tetrahydroxyflavone at 100mg/kg induced a significant decrease ($p < 0.05$) in soduim, chlorine ions, urea, and creatinin.

3.1 Histology of the Kidney

Plates 1-7 displays the photomicrographs of transverse sections of the rat's kidney in the control group and test groups.



	Control	NCG	LDEG	MDEG	HDEG	(PCG1)	(PCG2)
■ K (mmol/l)	5	6.2	5.8	6.4	5.3	3.8	5
■ Na (mmol/l)	126	163	152	170	136	118	127
■ Cl (mmol/l)	64	65	63	62	61	42	59
■ HCO ₃ (mmol/l)	24	26	24	22	26	24	30
■ Urea (mmol/l)	8.4	9.5	9.2	8.3	5.1	8.7	3.6
■ Creatinin (mmol/l)	128	150	148	130	124	134	125

Fig. 1. Renal function markers




Fig. 2. % diff of renal function markers




Plate 1 - CG (H&E) x 600




Plate 2 - NCG (H&E) x 600




Plate 3 - LDEG (H&E) x 600




Plate 4 - MDEG (H&E) x 600




Plate 5 - HDEG (H&E) x 600




Plate 6 - PCG1 (H&E) x 600




Plate 7 - PCG2 (H&E) x 600

4. DISCUSSION

Diabetic kidney disease is the decline of kidney function over time in people who have diabetes. Protein loss in the urine owing to glomeruli destruction can become large, resulting in a low serum albumin level, which can lead to generalized body swelling (edema) and the nephrotic syndrome. Similarly, the estimated glomerular filtration rate (eGFR) may gradually decrease from a normal of over 90 ml/min/1.73m² to less than 15, indicating that the patient has end-stage renal disease [6]. It normally progresses slowly over time. Pathophysiologic problems begin with blood glucose levels that have been poorly regulated for a long time [12].

Hippocrates, the father of medicine as a logical discipline, researched and summarized many therapeutic characteristics of plants. Throughout history, many traditional systems, including Nigerian traditional medicine systems, have used plants as medicine. A plant must include biologically active components that contribute to improved health or reduced illness risk in addition to being nourishing in order to be medicinally active or useful. These are well-thought-out plants that would be found in a typical diet.

Biomarkers such as serum urea and creatinine were observed to be higher in the negative control group in the current investigation, indicating the development of diabetic nephropathy. The elevated renal parameters were reversed when *Fleurya aestuans* was given orally, and the mitigating effects were dose dependent. In comparison to normal control rats, diabetic rats (NCG) have higher electrolyte concentrations. Treatment with *Fleurya aestuans* extract considerably lowered these levels.

The start of diabetic nephropathy is marked by functional abnormalities such as elevated electrolytes, urea, and creatinine levels. Uremia (excess urea in the urine) is associated with diabetic nephropathy [13]. In diabetic rats (NCG), a significant increase in urine urea and creatinine levels indicates poor kidney function and a change in electrolyte metabolism. As a result, treatment with *Fleurya aestuans* stabilized these levels, demonstrating its therapeutic effect in the treatment of renal dysfunctions.

The histoarchitecture of the kidneys is normal in Plate 1 (CG). Plate 2 (NCG) depicts aberrant

renal corpuscles with glomerular degeneration and an enlarged Bowman's capsule capacity. Plate 3 (LDEG) revealed that the epithelial lining of the renal tubules has been lost. Hypertrophied renal corpuscles (RC) with normal glomeruli and increased Bowman's capsule capacity. Plates 4 (MDEG), 5 (HDEG), 6 (PCG1), and 7 (PCG2) reveal that the renal corpuscles (RC) and renal tubules (RT) of the rat kidney are normal, indicating that the diabetic nephropathy produced by ALX medication has been reversed.

Chronic hyperglycemia and oxidative stress cause structural and functional changes in the body. In ALX-diabetic rats (NCG), structural anomalies in the kidney (basement membrane thickening, mesangial expansion, and hypertrophy) result in an elevated kidney parameters [14]. Oral delivery of FAE to diabetic groups greatly reduced diabetic kidney dysfunctions, which is consistent with Donate research [2].

5. CONCLUSION

Overall, this study suggests that *Fleurya aestuans* leaves could be beneficial in the treatment of diabetic renal impairment caused by hyperglycemia in individuals.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Our institutional ethical committee reviewed and approved the study, which was given the reference number UPH/CEREMAD/REC/MM78/052.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nur Samsu. "Diabetic Nephropathy: Challenges in Pathogenesis, Diagnosis, and Treatment", *BioMed Research International*. 2021;1497449. Available: <https://doi.org/10.1155/2021/1497449>
2. Donate-Correa J, Luis-Rodríguez D, Martín-Núñez E. Inflammatory targets in

- diabetic nephropathy. Journal of Clinical Medicine. 2020;9:458.
3. Chen Y, Lee K, Ni Z, He JC. Diabetic kidney disease: challenges, advances, and opportunities. *Kidney Diseases*. 2020;6: 215–225.
 4. Etukudo, I. *Ethnobotany: Conventional and traditional uses of plants*. First Edition, Verdicts Press. 2003;130.
 5. Adetunji Oluwaseyi Adegoke, Jeremiah Olorunjuwon Olugbami, Ayodeji Mathias Adegoke, Michael Adedapo Gbadegesin, Oyeronke Adunni Odunola. Renoprotective and antidiabetic properties of methanol leaf extract of *Fleurya aestuans* in Wistar Rats. *Journal of Evidence-Based Integrative Medicine*. 2021;26:1-11.
 6. Ganiyat K, Oloyede, Olukoyejo, E, Ayanbadejo. phytochemical, toxicity, antimicrobial and antioxidant screening of extracts obtained from *Fleurya aestuans* (Gaud). *Journal of Medical Sciences*. 2014;14:51-59.
 7. Okereke Stanley C, Elekwa I, Chukwudoruo Chieme S. Preliminary phytochemical screening and gas chromatographic FID evaluation of *Fleurya aestuans* leaf extracts. *International Journal of Current Biochemistry Research*. 2014;2(3):37-43.
 8. Al-Attar AM, Abu Zeid IM. Effect of tea (*Camellia sinensis*) and olive (*Olea europaea* L.) leaves extracts on male mice exposed to diazinon. *Biomedical Research International*. 2013;1–6.
 9. Lorke D. A new approach to practical acute toxicity testing. *Archieve Toxicology*. 1983; 54(4):275-287.
 10. Mbaka GO, Adeyemi OO, Noronha CC, Anunobi CC, Okanlawon AO. Anti-hyperglycemic and hypoglycaemic effects of ethanol root extract of *Sphenocentrumjollyanum* normal and alloxan induced diabetic rabbits. *Brazilia Journal of Morphological Sciences*. 2009; 26 (2):123-127.
 11. Bancroft JD, Gamble M. *Theory and practice of histological techniques*. Churchill Livingstone, Edinburgh. 2002;16-64.
 12. Kopel J, Pena-Hernandez C, Nugent K. Evolving spectrum of diabetic nephropathy. *World Journal of Diabetes*. 2019;10:269–279.
 13. Pérez-Morales RE, del Pino MD, Valdivielso JM, Ortiz A, Mora-Fernández C, Navarro-González JF. Inflammation in diabetic kidney disease. *Nephron*. 2019; (143):12–16.
 14. Duni V, Liakopoulos S, Roumeliotis D, Peschos, Dounousi E. Oxidative stress in the pathogenesis and evolution of chronic kidney disease: untangling Ariadne's thread. *International Journal of Molecular Sciences*. 2019;20:37-11.

© 2022 Nwafor et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87887>