

Aspartame Administration Cum *Moringa oleifera* Seed Extract: Blood Glucose Indices and Renal Toxicity Perspectives

**Christopher Chiedozie Mbadugha^{a*}, Aquaisua Nyong Aquaisua^a,
Emmanuel Uka^b, Ikechukwu Ezeah^c, Ndubuisi Nwodo^d
and Chiemela Chijioke Chinkata^a**

^a Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, University of Uyo, Nigeria.

^b Department of Biochemistry, Faculty of Science, University of Uyo, Nigeria.

^c Department of Surgery, University of Medical Sciences, Ondo State, Nigeria

^d Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Madonna University, Elele Alimini, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88098>

Original Research Article

Received 17 April 2022

Accepted 29 June 2022

Published 04 July 2022

ABSTRACT

Haematological parameters have been implicated in health indices. Aspartame is associated with elevated blood glucose level and toxicity in some body organs by alteration of their tissue histology. This study investigated the ameliorative potential of *Moringa oleifera* seed extract on adult male Albino Wistar rats following aspartame consumption. The rats were separated into nine groups and left to acclimatize for two weeks. The experiment lasted for 28 days during which various administration and measurements were carried out. On the 29th day the rats were anaesthetized; the blood and the kidney harvested for blood glucose analysis and routine histological procedures, respectively. Raw data obtained were expressed as mean \pm standard error of mean (M \pm SEM) and subjected to one way analysis of variance (ANOVA) using primer software (version 3.01) and post-hoc analysis using the Newman-Kuel test. $P < 0.05$ was regarded as being significant. It is observed that administration of ethanolic extract of *M. oleifera* seed immediately after aspartame

*Corresponding author: E-mail: christophermbadugha@uniuyo.edu.ng;

consumption, adequately checkmated rise in blood glucose level and maintain it at the optimum level along the normal reference range. Weight gain is observed within groups and across the groups of rats, suggesting that aspartame plays no useful role in weight loss. N-hexane extract of *M.oleifera* seed has the capacity to curtail the destruction of tissue architecture induced by aspartame, unlike ethanolic extract of *M. oleifera* seed. Hence, this study provides new insight on being cautious in consuming aspartame.

Keywords: Adult male albino wistar rats; aspartame; blood; ethanolic extract of *Moringa oleifera* seed; kidney; n-haxene extract of *Moringa oleifera* seed.

1. INTRODUCTION

The kidney is a bean shaped organ with a convex lateral margin and a concave medial side which houses the hilum, from which the renal artery enters and the renal vein and pelvis exits the kidney. The hilum serves as a passage for the ureter to exit, and blood and lymph vessels enter and exit. The kidney has three major regions: the renal cortex, renal medulla and renal pelvis. The smaller branches, the minor calyces come from each major calyx. Each kidney contains about 1 million functional units called the nephron that consists of simple, single layered epithelium along their entire length. Nephron, is the basic structural and functional unit of the kidney that regulates water and soluble substances in the blood. Distal convoluted tubules and collecting ducts, is the final site for reabsorption in the nephron. It is permeable to water and ion [1].

Moringa oleifera is one of the world's most useful trees, with almost every part of the tree used for food, medications and industrial purposes [2]. *M. oleifera* pod is reported to contain large amount of potassium, vitamins A and C [3,4]. The leaves are the most nutritious parts of the plant being a significant source of Vitamin B, K and C, pro Vitamin A as beta-carotene, manganese and protein among other essential nutrients [5].

Aspartame is a popular dietetic sweetener. Aspartame breaks down in part into phenylalanine, which interferes with the action of an enzyme intestinal alkaline phosphatase (IAP) previously shown to prevent metabolic syndrome (a group of symptoms associated with type 2 diabetes and cardiovascular disease) [6]. People who regularly consume artificial sweeteners are at increased risk of "excessive weight gain, metabolic syndrome, type 2 diabetes, and cardiovascular disease," [7]. Several studies link aspartame to weight gain, increased appetite, diabetes, metabolic derangement and obesity-related diseases. This notion implicating aspartame to weight gain and obesity-related

diseases raises some doubt as regards the legality of marketing aspartame-containing products as "diet" or weight loss aids. The aim of the study is to investigate and further determine the effect of *M. oleifera* seed extract on aspartame-induced renal toxicity. The specific objectives are: To determine the effect of ethanolic and N-Hexane extracts of *M. oleifera* seed on the body weight of adult male albino Wistar rats following aspartame administration. To verify the blood glucose levels of adult male albino Wistar rats following administration of aspartame, ethanolic extract and n-Hexane extract of *M.oleifera* seed.

2. METHODOLOGY

2.1 Experimental Animal Handling

Twenty seven adult male albino Wistar rats weighing between 150-215g were used. The animals were weighed, labeled and kept in wooden cages and allowed to acclimatized under optimum environmental conditions of temperature for 2 weeks prior to the commencement of the research work. They were fed pelletized growers and water ad libitum. The ethics of animal care was fully adopted following the guidelines for the care and use of laboratory animals.

2.2 Preparation of Extract of *M. oleifera* Seed

Fresh *M. oleifera* pods were harvested from the *M. oleifera* tree in the botanical garden. The pods were carefully opened and the seeds collected and powdered using mortar and pestol. The sample was put into the thimble section and both end covered with a cotton wool. A condenser is fixed above with a water inlet and outlet and solvent in the distillation flask below. The flask is kept on a heating mantle and the solvent is heated under reflux. Solute were transferred from the extraction chamber into the reservoir. There was continuous repetition of the process till ethanolic extraction is completed, [8], and the

solid residue is subjected further to n-Hexane extraction.

2.3 Administration of Extracts and Aspartame

Both the ethanolic extract of *M. oleifera* seed and N-Hexane extract of *M. oleifera* seed; and the aspartame were administered as follows: Rats in group A were given distilled water. This group served as the Control. Group B rats were given 0.4 ml (low dose) aspartame for 4 weeks (28 days). Group C rats were given 1.5 ml (high dose) aspartame for 4 weeks (28 days). Group D1, 0.6 ml (low dose) aspartame for the first 3 weeks and on the fourth week, 1.5 ml (low dose) ethanolic extract of *M. oleifera* seed only. Group D2, 0.5 ml (low dose) aspartame for the first 3 weeks and on the fourth week, 1.3 ml (low dose) N-Hexane extract of *M. oleifera* seed only. Group E1, 1.30 ml (high dose) aspartame for the first 3 weeks and on the fourth week, 2.5 ml (high dose) ethanolic extract of *M. oleifera* seed only.

Group E2, 1.5 ml (high dose) aspartame for the first 3 weeks and on the fourth week, 3.0 ml (high dose) N-Hexane extract of *M. oleifera* seed only. Group F1, 1.4 ml (high dose) aspartame followed by 2.8 ml (high dose) ethanolic extract *M. oleifera* seed, for four weeks, and group F2, 1.3 ml (high dose) aspartame followed by 2.5 ml (high dose) N-Hexane extract *M. oleifera* seed for four weeks.

2.4 Termination of Experiment and Tissue Processing

The rats were anaesthetized with chloroform and dissected from the superficial layer of the

abdomen and directed craniomedially to the thorax. The blood was gotten directly from the heart and preserved in an anti coagulant bottle, 10% buffered formalin was transcardially perfused to preserve the tissues. The kidney was carefully harvested and fixed in 10% buffer formalin for 48 hour, to maintain the tissues morphology as faithfully as possible compared to the living state. The tissue processing involved the following stages: dehydration, clearing, infiltration, embedding, sectioning, and staining. Haematoxylin and Eosin staining method for routine histological studies were employed [9].

2.5 Statistical Analysis

One way analysis of variance (ANOVA) using primer software (version 3.01); post-hoc analysis using the Newman-Kuel test was applied to test the significant levels. Data with probability $p < 0.05$ were regarded as significant.

3. RESULTS

3.1 Comparison of Changes in Body Weight of Adult Male Albino Wister Rats

When it comes to blood glucose measurement, the standard and criteria used for rats are slightly comparable and close to that used for human. The normal range of fasting sample is 3.0 to 5.5 mmol/l. Figures below 3.0 mmol/l predisposes to hypoglycemia. Figures above 7.0 mmol/l indicate hyperglycemia, while figures above 10.0 mmol/l point to diabetic condition. Based on data above, Control group A, has blood glucose level within the normal reference range.

Table 1. Blood Glucose Count for Adult Male Albino Wistar Rats Administered Aspartame

Group	M±SEM (Mmol/l)	SD
A	3.42±0.28	0.48
B	0.94±0.01	0.02
C	3.88±0.48	0.83
D1	3.52±1.16	2.01
D2	3.00±2.51	4.35
E1	5.16±0.53	0.92
E2	6.18±2.11	3.65
F1	3.52±0.89	1.54
F2	3.85±0.56	0.97
F = 1.45, P = 0.24, P < 0.5		




Fig. 1. Comparison of changes in body weight of Albino Wistar rats. (data are expressed in mean and standard error of mean, $M \pm SEM$)

Key:

A = rats given distilled water Group A (Control).

B = rats given 0.4 ml (low dose) aspartame for 4 weeks (28 days)

C = rats given 1.5 ml (high dose) aspartame for 4 weeks (28 days)

D1 = rats given 0.6 ml (low dose) aspartame for the first 3 weeks and on the fourth week, 1.5ml (low dose) ethanolic seed extract of *M. oleifera* only.

D2= rats given 0.5 ml (low dose) aspartame for the first 3 weeks and on the fourth week, 1.3 ml (low dose) N-Hexane seed extract of *M. oleifera* only.

E1 = rats given 1.30ml (high dose) aspartame for the first 3 weeks and on the fourth week, 2.5 ml (high dose) ethanolic seed extract of *M. oleifera* only.

E2 = rats given 1.5 ml (high dose) aspartame for the first 3 weeks and on the fourth week, 3.0 ml (high dose) N-Hexane seed extract of *M. oleifera* only.

F1 = rats given 1.4 ml (high dose) aspartame followed by 2.8 ml (high dose) ethanolic seed extract *M. oleifera* for four weeks.

F2 = rats given 1.3 ml (high dose) aspartame followed by 2.5 ml (high dose) N-Hexane seed extract *M. oleifera* for four weeks.

Experimental group B, is an exception in that it points to hypoglycemic state following low dose aspartame administration. In experimental group C, high dose of aspartame administration brought about a slight increase in blood glucose level compared to the control, however, the stated value is still within the normal reference range. In group D1, though there was a slight increase above the control, but still below the value obtained following administration of low dose of aspartame; thus, suggesting that ethanolic extract of *M. oleifera* seed may play a role in lowering blood glucose level, when the

aspartame is administered in low doses. In D2, there was a notable decrease in the blood glucose level, also suggesting that n-Hexane extract of *M. oleifera* seed may play a greater role than the ethanolic extract of *M. oleifera* seed in lowering blood glucose level, when the aspartame is administered in low doses. In E1, the value obtained suggested that ethanolic extract of *M. oleifera* seed, lack the capacity to checkmate the rise in the blood glucose level following aspartame administration in large doses; irrespective of the fact that it maintained it within the normal reference range. The same is

applicable to E2, in which n-hexane extract of *M. oleifera* seed is administered. However, the n-Hexane extract of *M. oleifera* seed is behind the ethanolic extract of *M. oleifera* seed in efficacy. In F1, It is observed that administration of ethanolic extract of *M. oleifera* seed immediately after aspartame consumption, adequately checkmated rise in blood glucose level and

maintain it at the optimum level along the normal reference range. In F2, it was also observed that administration of n-Hexane extract of *M. oleifera* seed, following aspartame consumption equally checkmated rise in blood glucose level but not as effectively done by the ethanolic extract of *M. oleifera* seed.




Fig. 2. Blood glucose count for adult male albino wistar rats administered aspartame

3.2 Hematoxylin and Eosin (H&E) Method for General Demonstration of Kidney




Fig. 3. (H&E method, X100)




Fig. 4. (H&E method, X400)

Cross section of the Kidney Cortex of adult male albino Wistar rat given distilled water. Group A (Control). (H&E method, X400). Section revealed features typical of normal kidney cortex: Glomerular (Bowman's) capsule GmCps. Glomerular capillary GmCp. Proximal Convoluted tubule PrCoTb. Parietal Layer PrLy. Visceral layer VsLy. Distal convoluted tubule DsCoTb. Capsular Space CpSp. Urinary pole UrPI




Fig. 5. (H&E method, X100)




Fig. 6. (H&E method, X400)

Cross section of the Kidney Cortex of adult male albino Wistar rat given 0.40 ml (low dose) aspartame for 4 weeks Group B. (H&E method, X400). Section revealed: Atrophy of collecting tubule AtCITb. Atrophy of urinary pole AtUrPI. Inference: Severely affected.




Fig. 7. (H&E method, X100).




Fig. 8. (H&E method, X400)

Cross section of the Kidney Cortex of adult male albino Wistar rat given 1.50 ml (high dose) aspartame for 4 weeks. Group C. (H&E method, X400). Section revealed: Inflammation of capsular space InfCpSp. Atrophy of distal convoluted tubule AtDsCt. Atrophy of proximal convoluted tubule AtPrCt. Inference: Severely affected.




Fig. 9. (H&E method, X100)




Fig. 10. (H&E method, X400)

Cross section of the Kidney Cortex of adult male albino Wistar rat given 0.60 ml (low dose) aspartame for the first 3 weeks and on the fourth week, 1.50 ml (low dose) ethanolic extract of *Moringa oleifera* seed only. Group D1. (H&E method, X400). Section revealed: Atrophy of Juxtaglomerular cells AtJsGlmCs. Atrophy of Macula densa AtMcDs. Atrophy of glomerular capsule AtGlmCp. Necrosis of glomerular capillaries NcGlmCp. Inference: Severely affected.




Fig. 11. (H&E method, X100)




Fig. 12. (H&E method, X400)

Cross section of the Kidney Cortex of adult male albino Wistar rat given 0.50 ml (low dose) aspartame for the first 3 weeks and on the fourth week, 1.30 ml (low dose) N-Hexane extract of *Moringa oleifera* seed only. Group D2. (H&E method, X400). Section revealed intact: : Glomerular (Bowman's) capsule GmCps. Glomerular capillary GmCp. Proximal Convolved tubule PrCoTb. Capsular Space CpSp. Collecting tubule CITb. Inference: Unaffected.




Fig. 13. (H&E method, X100).




Fig. 14. (H&E method, X400).

Cross section of the Kidney Cortex of adult male albino Wistar rat given 1.30ml (high dose) aspartame for the first 3 weeks and on the fourth week, 2.50 ml (high dose) ethanolic extract of *Moringa oleifera* seed only. Group E1. (H&E method, X400). Section revealed: Atrophy of both proximal and distal convoluted tubules AtCoTb. Inflammation of glomerular arterioles InfGmAt. Inference: Severely affected.




Fig. 15. (H&E method, X100).




Fig. 16. (H&E method, X400).

Cross section of the Kidney Cortex of adult male albino Wistar rat given 1.50 ml (high dose) aspartame for the first 3 weeks and on the fourth week, 3.00 ml (high dose) N- Hexane extract of *Moringa oleifera* seed only. Group E2. (H&E method, X400). Section reveal intact: Glomerular (Bowman's) capsule GmCps. Glomerular capillary GmCp. Capsular Space CpSp. Juxta - glomerular cells JsGlmCs. Inference: Unaffected.




Fig. 17. (H&E method, X100).




Fig. 18. (H&E method, X400).

Cross section of the Kidney Cortex of adult male albino Wistar rat given 1.40 ml (high Dose) aspartame followed by 2.80 ml (high dose) ethanolic extract of *Moringa oleifera* seed for four weeks. Group F1. (H&E method, X400). Section revealed: Edema of collecting tubul EdCoTb. Inflammation of Capsular space InfCpSp. Inference: Slightly affected.




Fig. 19. (H&E method, X100)




Fig. 20. (H&E method, X400)

Cross section of the Kidney Cortex of adult male albino Wistar rat given 1.30 ml (high Dose) aspartame followed by 2.50 ml (high dose) N-Hexane extract of *Moringa oleifera* seed for four weeks. Group F2. (H&E method, X400). Section reveal: Glomerular (Bowman's) capsule GlmCps. Glomerular capillary GmCp. Inference: Unaffected.

In our research, the control group A, has blood glucose level within the normal reference range, as stated above. Same figures were obtained from other experimental groups, namely: C, D1, E1, F1 and F2, with slight discrepancies based on the substance administered, with the exceptions of F2, that tilted toward hyperglycemic state and B toward hypoglycemic condition. Besides, Palmnäs et al [13] in their work opined that "Aspartame consumption increased the fasting glucose concentrations in both the standard feed pellet diet and high-fat groups independent of body composition". Their research result had given credence to the outcome of our own work in which there was gradual increase in blood glucose values across the groups as the experiment progressed; though none exceeded the normal reference range, except in group B, in which the mechanism underlying it is not well understood. The spike in glucose level seen in E2 being as a result of large doses of aspartame administered; and it also worthy to note that irrespective of the types of substances and feed administered, there was

4. DISCUSSION OF FINDINGS

There was observed increase in weight within and across groups of rats used for the research; these findings of ours are corroborated by the work of Daniel and colleagues [10], in which they stated that "even acceptable daily dose of aspartame, as regulated by the United States Food and Drug Administration (FDA), might make you hungrier and lead to weight gain".

The average fasting blood glucose (FBG) of normal Wistar rats was (3.95 +/- 1.31) mmol/L, and the 95% upper limit was 6.2 mmol/L. [11,12].

consistency in the rise in blood glucose level within and across groups.

According to Isabela et al [14]; “ via p53 activation, aspartame inhibited a transcriptional coactivator, the peroxisome proliferator-activated receptor gamma coactivator 1 alpha, a master regulator of glucose and lipid metabolism, probably leading to changes in lipid profile in serum, total lipid accumulation, as well as an impairment in the gluconeogenesis in mouse liver, thus causing hypoglycemia”. The outcome of the work of Isabela and colleagues might be the plausible and possible mechanism and explanation that underlies the hypoglycemic state observed in group B, in our own experiment.

All the stated facts above notwithstanding, the US Food and Drug Administration (FDA), [15], has concluded that it is safe to use aspartame as a general purpose sweetener. The FDA went on to indicate that aspartame is yet to be conclusively implicated in any specific health problems; rare exceptions, being found only in people with phenylketonuria (PKU). Besides, the European Food Safety Authority (EFSA), [16] has stated that people with conditions such as: brain tumours, leukemia, lymphatic and haematopoietic cancers as well as a variety of cancers, are not at increased risk following aspartame consumption.

There was an observed dose dependent alteration of histology of renal tissues and this has a link in cases where there were increases in blood glucose level, thus giving credence to the fact that of diagnostic significance in routine clinical evaluation as regards health condition is hematological parameters. In the experimental group E2, there was increased blood glucose level that predisposed to hyperglycemia; this in fact, introduced a gap in the evidence that hematologic parameters are of diagnostic significance in routine clinical evaluation; however, in same experimental group E2, the N-Hexane extract of *M. oleifera* seed, effectively checkmated the disruptive effect of aspartame on the kidney cells, thus, substantiating the suggestion by Mutasim, et al., [17], that *M. oleifera* may have potential for use as source of natural treatment for diseases such as cancer, hepatocarcinoma cells in vitro. Our research finding is further supported by the work of Julie and Gary, [18]; from which they concluded that the Consumption of ≥ 2 servings per day of artificially sweetened soda is associated with a 2-fold increased odds for kidney function decline in

women. Moreover, Arbind, et al., [19] in their study concluded that over a long time duration, the administration of about 40 mg/kg. bw/day, of aspartame orally may result to oxidative stress, caused by methanol, which itself is an aspartame metabolite and can be injurious to the liver and kidney.

5. CONCLUSION

N-hexane extract of *M.oleifera* seed has ameliorative effects on the destruction of tissue architecture induced by aspartame. Hence, this study provides new insight on being cautious in consuming aspartame.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh GP, Sharma SK. Antimicrobial evaluation of leaf Extract of Moringa oleifera Lam. International Research Journal of Pharmacology. 2012;3:1-4.
2. Khalafalla MM, Abdellatef E, Dafalla HM, Nassrallah AA, Aboul-Enein KM, Lightfoot DA, El-Deeb FE, El-Shemy HA. Active principle from Moringa oleifera Lam leaves effective against two leukemias and a hepatocarcinoma. African Journal of Biotechnology. 2010;9(49):8467-71.
3. Fuglie LJ. The miracle tree: Moringa oleifera: natural nutrition for the tropics. Dakar. 68 pp revised in 2001 and published as the miracle tree: the multiple attribute of Moringa. 1999;172.
4. Fahey JW. Moringa oleifera: a review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. Part 1. Trees for life Journal. 2005;1(5):1-5.
5. Leone A, Spada A, Battezzati A, Schiraldi A, Aristil J, Bertoli S. Cultivation, genetic,

- ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera* leaves: An overview. International journal of molecular sciences. 2015 Jun 5;16(6):12791-835.
Available:<https://doi.org/10.3390/ijms160612791>
6. Gul SS, Hamilton AR, Munoz AR, Phupitakphol T, Liu W, Hyoju SK, Economopoulos KP, Morrison S, Hu D, Zhang W, Gharedaghi MH. Inhibition of the gut enzyme intestinal alkaline phosphatase may explain how aspartame promotes glucose intolerance and obesity in mice. *Applied Physiology, Nutrition, and Metabolism*. 2017;42(1):77-83.
 7. Swithers SE. Artificial sweeteners produce the counterintuitive effect of inducing metabolic derangements. *Trends in Endocrinology & Metabolism*. 2013;24(9):431-41.
DOI: 10.1016/j.tem.2013.05.005
 8. Soxhlet F. Die gewichtsanalytische bestimmung des milchfettes. *Dingler's Polytechnisches Journal*. 1879;232(461-465).
 9. Drury RA, Wallington EA. *Carltons Histological Techniques*, 5th edn. New York, Oxford University Press. 1980;206.
 10. Childs DS, Jatoi A. A hunger for hunger: a review of palliative therapies for cancer-associated anorexia. *Annals of Palliative Medicine*. 2019;8(1):50.
DOI: 10.21037/apm.2018.05.08
 11. Johnson-Delaney CA. *Exotic companion medicine handbook for veterinarians*. Wingers Publishing Incorporated; 1996.
 12. Wei Sheng Yan Jiu. 2010;39(2):133-7, 142.
 13. Palmnäs MS, Cowan TE, Bomhof MR, Su J, Reimer RA, Vogel HJ, Hittel DS, Shearer J. Low-dose aspartame consumption differentially affects gut microbiota-host metabolic interactions in the diet-induced obese rat. *PloS one*. 2014;9(10):e109841.
 14. Finamor IA, Bressan CA, Torres-Cuevas I, Rius-Pérez S, da Veiga M, Rocha MI, Pavanato MA, Pérez S. Long-Term Aspartame Administration Leads to Fibrosis, Inflammasome Activation, and Gluconeogenesis Impairment in the Liver of Mice. *Biology*. 2021 Jan 22;10(2):82.
Available:<https://doi.org/10.3390/biology10020082>
 15. US Food and Drug Administration (FDA). Available:[www.fda.gov/food/ingredient-spacing-labeling/food-additives-ingredients/ucm397725.htm](http://www.fda.gov/food/ingredient-spacing-labeling/food-additives-ingredients-ucm397725.htm)
 16. European Food Safety Authority (EFSA). Available:www.efsa.europa.eu/en/topics/topic/aspartame
 17. Khalafalla MM, Abdellatef E, Dafalla HM, Nassrallah AA, Aboul-Enein KM, Lightfoot DA, El-Deeb FE, El-Shemy HA. Active principle from *Moringa oleifera* Lam leaves effective against two leukemias and a hepatocarcinoma. *African Journal of Biotechnology*. 2010;9(49):8467-71.
 18. Lin J, Curhan GC. Associations of sugar and artificially sweetened soda with albuminuria and kidney function decline in women. *Clinical Journal of the American Society of Nephrology*. 2011;6(1):160-6.
DOI: 10.2215/CJN.03260410. Epub 2010 Sep 30.
 19. Kumar Choudhary A, Selvaraj S, Sheela Devi R. Aspartame induce modification in membrane bound and antioxidant enzymes in liver and kidney of Wistar albino rats. *Current Nutrition & Food Science*. 2014;10(4):275-87.
DOI:10.2174/1573401310666141024215431

© 2022 Mbadugha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/88098>