



Efficacy of Mechanical Seedling Transplanter and Deep Placement of Mixed Fertilizer on Rice Yield

Setara Begum ^{a*}, M. S. Islam ^a, M. M. Rashid ^a, M. R. Manir ^a,
M. H. Rahman ^{b++} and M. A. Hossen ^c

^a Farm Management Division, Bangladesh Rice Research Institute (BRRI), Gazipur-1701, Bangladesh.

^b Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh.

^c Farm Machinery and Post-Harvest Technology Division, Bangladesh Rice Research Institute (BRRI), Gazipur-1701, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAAR/2022/v20i2395

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95065>

Original Research Article

Received: 18/10/2022

Accepted: 26/12/2022

Published: 27/12/2022

ABSTRACT

A field experiment was conducted at the West Byde of BRRI farm, Gazipur during T. Aman and Boro, 2021-22 season to evaluate the efficacy of newly developed mechanical rice transplanter cum fertilizer applicator. Five treatments in Randomized Complete Block Design with three replications. Field experiments was conducted on silty clay loam soil in Gazipur. Individual plot size was 8m×5m with 30 cm buffer zone. The treatments were T₁ = Mechanical transplanting along with 100% fertilizer (Urea, TSP, MoP and Gypsum) deep placement, T₂ = Mechanical transplanting along with 80% fertilizer (80% Urea and 100% TSP, MoP and Gypsum) deep placement, T₃ = Mechanical transplanting along with 70% fertilizer (70% urea and 100% TSP, MoP and Gypsum)

** Deputy Registrar;

*Corresponding author: Email: setu.edu.bd@gmail.com;

deep placement, T₄ = Mechanical transplanting along with 100% fertilizer hand broadcasting (TSP, MoP and Gypsum fertilizer as basal dose and urea fertilizer in three split) and T₅ = Hand transplanting of same seedling of rice transplanter along with 100% fertilizer hand broadcasting (TSP, MoP and Gypsum in basal dose and urea fertilizer in three split) and T₆ = BRRI recommended practice. In this experiment, BRRI dhan89 and BRRI dhan92 was used in T. Aman and Boro seasons, respectively. In the field, saving percentage of urea fertilizer varied from 25% to 28% against the calibration of 30% of urea saving due to the variation of the machine, operational speed and more penetration of the driving wheel during operation. It was found that grain yield was varied with the mode and rate of urea fertilizer application. Mechanical transplanting along with urea fertilizer deep placement (80% of recommended dose) gave significantly higher yield compared to manual transplanting and hand broadcasting of urea. Mechanical transplanting along with deep placement of urea fertilizer is a promising technology for rice farmers in Bangladesh. Manual transplanting is tedious and time consuming as a result mechanical transplanting is seen as a solution of labor problems. No significant differences among the treatments was found in growth parameters. Significant variations were recorded in case of yield. BRRI recommended practice and mechanical transplanting with 80% urea with other fertilizers were produced very similar grain yield. From these results, it might be said that mechanical transplanting with 80% urea fertilizer is recommended with BRRI recommended hand transplanting practice. Urea saving is additional benefit with time savings and low transplanting cost when transplanted with rice transplanter cum fertilizer applicator.

Keywords: Mechanical transplanting; rice transplanter; mixed fertilizer; fertilizer deep placement.

1. INTRODUCTION

“Rice is the dominant and staple food crop, plays an important role in food security of about 160 million people in Bangladesh. It provides nearly 48% of rural employment, about two-third of total calorie supply and one-half of the total protein consumption of the country’s people depend on rice” [1]. “Rice production in Bangladesh is a crucial part of the national economy. Bangladesh has ranked third globally in rice production for the fourth consecutive year with a projected output of 38.4 million tonnes. (‘Food Outlook-June 2022’.) Cultivation area of rice, per unit yield, overall production and cropping intensity expanded appreciably over the last 45 years. Employed labor in the agricultural sector decreased remarkably due to increasing rural to urban migration with the growth of non-farm employment opportunities, inflicting rural labour shortages” [2]. “Half of agricultural GDP and one-sixth of rural household income depends on rice” [3]. “Modernization of rice production is the utmost need to feed the growing population of the country from gradually decreasing land. It mostly depends on the agricultural mechanization that helps to enhance the agricultural and food production of the country” [4]. Due to labor scarcity, there is a need to explore establishment methods for rice that require less labor but still allow the crop to be transplanted on time. The use of a mechanical transplanter is one alternative to address this

issue. Mechanical transplanting of rice is the process of transplanting young rice seedlings, which have been grown in a mat nursery, using a self-propelled rice transplanter. In conventional manual transplanting practice, 8-12 laborers are required to transplant one acre. However, if a self-propelled rice transplanter is used, three people can transplant up to four acres in a day. “Manual transplanting is tedious and time consuming which frequently causes delayed rice transplanting. One month delayed of transplanting reduces yield about 25% while two months delayed reduces yield about 70%” [5]. “A total of 156.2 man-days is required for rice production in one hectare of land from which 44.5 man-days (28.24% of total labor requirement) are consumed by seedling raising and transplanting” [6]. “Thus, mechanized rice transplanting is seen as an answer to the present labor crisis. Rice seedling transplanting using mechanical rice transplanter ensures uniform plant spacing as well as fast and efficient planting also. With this point of view, mixed fertilizer (Urea, TSP, MoP and Gypsum) deep placement mechanisms incorporate with the existing walking type rice transplanter. Deep placement of NPK fertilizer significantly increased grain yields and net economic return across all the rice-growing seasons and years compared to traditional practices” [7]. “because only 30 to 50% N used effectively by the plant from top dressing of urea fertilizer” [8]. “Top dressing of prilled urea fertilizer produced higher

amounts of ammonium N in floodwater compared to deep placement of urea” [9]. “The higher amount of ammonium N in floodwater leads to a higher amount of ammonia volatilization loss” [10]. “Broadcast application of N, P, and K fertilizers increases nutrients concentration in floodwater” [9] but “significant losses occur with surface runoff. Urea fertilizer deep placement (UDP) or fertilizer (Urea, TSP, MoP and Gypsum) deep placement (FDP) is an effective management practice for wetland transplanted rice by increasing its productivity and reducing fertilizer losses” [11,12]. “By using UDP technology, yield also increased significantly with less use of urea about 10-20 as compared to prilled urea” [13]. An endeavor has been undertaken to attach the fertilizer deep placement technology to the prevailing mechanical rice transplanter (both the walking and riding) without sacrificing the merit of transplanting to confirm both the mechanized rice transplanting and mixed fertilizer deep placement simultaneously. From the above discussion this type of research was done to evaluate the efficacy of mechanical rice transplanter with deep placement of mixed fertilizer and to observe the yield and yield contributing parameters.

2. MATERIALS AND METHODS

This experiment was conducted at the West Byde of BIRRI farm, Gazipur during T. Aman and Boro, 2021-22 seasons to evaluate the efficacy of newly developed BIRRI mechanical rice transplanter cum fertilizer applicator. Urea fertilizer along with TSP, MoP and Gypsum fertilizer can be placed and covered in 6-8 cm soil depth during mechanical transplanting using the developed rice transplanter. Randomized Complete Block (RCB) design was followed with three replications. Individual plot size was 8m x 5m along with 30 cm buffer spacing. Treatments of the studies were T₁ = Mechanical transplanting along with 100% fertilizer (Urea, TSP, MoP and Gypsum) deep placement, T₂ = Mechanical transplanting along with 80% fertilizer (80% Urea and 100% TSP, MoP and Gypsum) deep

placement, T₃ = Mechanical transplanting along with 70% fertilizer (70% urea and 100% TSP, MoP and Gypsum) deep placement, T₄ = Mechanical transplanting along with 100% fertilizer hand broadcasting (TSP, MoP and Gypsum fertilizer as basal dose and urea fertilizer in three split) and T₅ = Hand transplanting of same seedling of rice transplanter along with 100% fertilizer hand broadcasting (TSP, MoP and Gypsum in basal dose and urea fertilizer in three split) and T₆ = BIRRI recommended practice. In this experiment, BIRRI dhan87 and BIRRI dhan92 was used in T. Aman and Boro seasons, respectively. Twenty days old mat type seedlings were used in mechanical transplanting at 30 cm x 15 cm spacing whereas spacing of manual transplanting was 20 cm x 20 cm. All intercultural operations were done according to BIRRI recommendation and were same for all treatments. Yield and yield components data were taken at harvesting time.

2.1 Statistical Analysis

Data were analyzed as a single factorial design according to Gomez and Gomez (1984) using Statistix 10 program (Statistix 10 software, 2013). Means were compared with the least significant difference (LSD) test. Simple correlation analysis was carried out with Excel 2010 to determine the relationship of grain yield to yield attributes.

3. RESULTS AND DISCUSSION

Growth and yield contributing parameters such as tiller number, panicle number, filled grain, unfilled grain, and 1000 grain weight were not significantly affected by mechanical transplanting along with fertilizer deep placement and hand transplanting along with hand broadcasting of fertilizer. Mechanical transplanting with 80% urea fertilizer along with 100% TSP, MoP, and Gypsum fertilizer deep placement gave the highest yield in T. Aman season where as this treatment gave same yield compared to BIRRI recommended practice during Boro season.

Table 1. Soil type and characteristics of experimental field

Characteristics	
Soil texture	Silty clay loam
pH	5.4
Organic carbon in % (OC)	1.3
Total N (%)	0.13




Fig. 1. Plant height in cm at days after transplanting as affected by mechanical seedling transplanting with deep placement of mixed fertilizer and hand transplanting with hand broadcasting in T. Aman and Boro, 2021-22 seasons




Fig. 2. Number of tillers per square meter at days after transplanting as affected by mechanical seedling transplanting with deep placement of mixed fertilizer and hand transplanting with hand broadcasting in T. Aman and Boro, 2021-22 seasons

3.1 Plant Height

Plant height was not significantly affected by mechanical seedling transplanting and deep placement of fertilizer in both T. Aman and Boro seasons (Table 1). Plant height of BRR1 recommended practice plot were found taller due to higher seedling age (15 days older than mechanical transplanted seedlings) which was finally similar when compared at maturity stage (Fig. 1).

3.2 Tiller Production

There were significant variations recorded in tiller production at different days after transplanting (Fig. 2) among the mechanical transplanting along with deep placement of fertilizer application and hand transplanting along with hand broadcasting of fertilizer during both T. Aman and Boro seasons (Fig. 1 and Fig. 2).

3.3 Panicle Number

Number of panicle m^{-2} was not varied significantly with the mechanical transplanting along with deep placement of fertilizers and hand transplanting along with hand broadcasting of fertilizer (Table 1).

3.4 Filled Grain Per Panicle and 1000 Grain-wt.

There were no significant differences among the treatments in case of number of filled grain panicle⁻¹ and 1000 grain weight. All the treatments produced statistically similar number filled grain panicle⁻¹ and 1000 grain weight (Table 1).

3.5 Grain Yield

Among different variables, only grain yield were significantly varied among the treatments (Table

2). Deep placement of urea fertilizer as briquette form also gave higher yield compared to hand broadcasting method (Hossen, 2013). In T. Aman season, T₂ (Mechanical transplanting along with 80% fertilizer deep placement) gave the highest grain yield (4.66 t ha⁻¹) followed by T₄, T₃ & T₁ (4.48, 4.38 and 3.31 t ha⁻¹, respectively). However, grain yield of treatment T₁ gave statistically lower yield compared to others. This might be due to the severe attack of criseck just after transplanting. Due to high amount of urea application at transplanting time in treatment T₁ foster the severity of creseck and ultimately yield was hampered.

On the other hand, during Boro season T₆ (BRRR recommended practice), T₂ (mechanical transplanting with 80% urea) and T₁ (mechanical transplanting with 100% urea) produced similar grain yield which was higher than the other treatments (Table 2).

Table 2. Yield and yield contributing parameters of rice as affected by mechanical seedling transplanting with deep placement of mixed fertilizer and hand transplanting with hand broadcasting in T. Aman and Boro, 2021-22 seasons

Treatments	Plant height (Cm)	Panicle m^{-2} (no.)	Filled grain panicle ⁻¹ (no.)	1000 grain wt. (gm)	Grain yield (t ha ⁻¹)
T. Aman season (BRRR dhan87)					
T ₁	127	214	152	23.37	4.17
T ₂	130	213	153	23.00	4.54
T ₃	133	218	157	23.02	4.38
T ₄	131	218.	167	23.33	4.48
Isd (0.05)	NS	0.60	NS	NS	NS
CV%	6	3.32	9.05	1.49	5.72
Boro season (BRRR dhan92)					
T ₁	116	406	113	22.49	7.26
T ₂	111	391	110	22.63	7.32
T ₃	110	417	102	22.39	6.52
T ₄	111	376	108	22.79	6.79
T ₅	113	328	103	21.62	6.14
T ₆	115	384	110	22.54	7.47
Isd (0.05)	NS	NS	NS	NS	0.59
CV%	2.36	7.75	2.57	4.23	4.77

T₁= Mechanical transplanting along with 100% fertilizer (Urea, TSP, MoP and Gypsum) deep placement, T₂= Mechanical transplanting along with 80% fertilizer (80% Urea and 100% TSP, MoP and Gypsum) deep placement, T₃= Mechanical transplanting along with 70% fertilizer (70% urea and 100% TSP, MoP and Gypsum) deep placement, T₄= Mechanical transplanting along with 100% fertilizer hand broadcasting (TSP, MoP and Gypsum fertilizer as basal dose and urea fertilizer in three split) and T₅= Hand transplanting along with 100% fertilizer hand broadcasting (TSP, MoP and Gypsum in basal dose and urea fertilizer in three split). T₆= BRRR recommended practice.

4. CONCLUSIONS

It may be concluded that no significant differences among the treatments was found in growth parameters. Significant variations were recorded in case of yield and BRRl recommended practice and mechanical transplanting with 80% urea with other fertilizers were produced very similar grain yield. From these results, it might be said that mechanical transplanting with 80% urea fertilizer is recommended with BRRl recommended hand transplanting practice. Urea saving is additional benefit with time saving and low transplanting cost (labour cost) when transplanted with rice transplanter cum fertilizer applicator. Mechanical rice transplanter is a promising technology considering the present labor crisis in Bangladesh.

ACKNOWLEDGEMENTS

The authors are thankful to FMPHT Division, Bangladesh Rice Research Institute for providing supports to conduct field experiments. This research did not receive any specific funding.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. BRKB (Bangladesh Rice Knowledge Bank). Rice in Bangladesh; 2018. Available:<http://www.knowledgebank-brrl.org/riceinban.php>
2. Ziauddin ATM, Ahmed. Research priorities in farm machinery, irrigation & water management and post-harvest technology (engineering aspects), agricultural research priority, vision- 2030 and beyond, Bangladesh Agricultural Research Council, 2010; 2012.
3. Miah MAM, Gaihre YK, Hunter G, Singh U, Hossain SA. Fertilizer deep placement increases rice production: evidence from farmers' fields in Southern Bangladesh. *Agronomy Journal*. 2016;108:1–8. Available:<https://doi.org/10.2134/agronj2015.0170>
4. BRRl. Introduction and characterization of BRRl rice varieties. In *Adhunik Dhaner Chas*, 19th ed., Agami Printing and Publication Co., 27 Neelkhet, Dhaka 1205. 2016a;6–15.
5. Rahman F, Shammi SA, Parvin MT, Akter N, Khan MS, Haque S. Contribution of rural women to rice production activities in two different areas of Bangladesh. *Progressive Agriculture*. 2016;27(2):180–188. Available:<https://doi.org/10.3329/pa.v27i2.29329>
6. Hossen MA, Hossain MM, Haque ME, Bell RW. Transplanting into non-puddled soils with a small-scale mechanical transplanter reduced fuel, labour and irrigation water requirements for rice (*Oryza sativa* L.) establishment and increased yield. *Field Crops Research*. 2018;225:141–151.
7. Hossen MA. Mechanization in Bangladesh: Way of modernization in agriculture. *International Journal of Engineering Trends and Technology*. 2019;67(9):69-77. Available:<https://doi.org/10.14445/22315381/IJETT-V67I9P212>
8. Statistix 10 software. An Analytical Software of Statistix 10, Analytical Software, 2105 Miller Landing Rd, Tallahassee, FL 32312, USA; 2013.
9. BRRl. Fertilizer management. In *Adhunik Dhaner Chas*, 19th ed., Agami Printing and Publication Co., 27 Neelkhet, Dhaka 1205. 2016b;24–30.
10. Rochette P, Angers DA, Chantigny MH, Gasser MO, MacDonald JD, Pelster DE, Bertrand N. Ammonia volatilization and nitrogen retention: How deep to incorporate urea? *Journal of Environment Quality*. 2013;42:1635–1642. Available:<https://doi.org/10.2134/jeq2013.05.0192>
11. Kapoor V, Singh U, Patil SK, Magre H, Shrivastava LK, Mishra VN. Rice growth, grain yield, and floodwater nutrient dynamics as affected by nutrient placement method and rate. *Agronomy Journal*. 2018;100:526–536. Available:<https://doi.org/10.2134/agronj2007.0007>
12. Bowen WT, Diamond RB, Singh U, Thompson TP. Farmer and environment benefits derived from deep placement of urea briquettes for flooded rice in Bangladesh. In: Zhaoliang Zhu et al. (ed.) 3rd Int. Nitrogen Conf., Contributed Papers, Inst. of Soil Science, Nanjing, China. 12-16 Oct. 2004. Science Press USA, Monmouth Junction, NJ. 2015;71-76.

13. Hossen MA, Islam AKMS, Zaman MK, Khatun A, Alam MA, Bhuiyan MKA. Mechanical deep-placement of urea fertilizer with a prilled urea applicator saved fertilizer and increased yield for long duration wet land rice variety. Bangladesh Journal of Progressive Science & Technology. 2017;15(2):25–30.

© 2022 Begum et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/95065>