



Breastmilk Jaundice – Could it be more Common than We Realize?

Asinobi Isaac Nwabueze^{a,b*}

^a *Department of Paediatrics, College of Medicine, Enugu State University of Science and Technology, Enugu, Nigeria.*

^b *Department of Paediatrics, Enugu State University of Science and Technology Teaching Hospital, Enugu, Nigeria.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/JAMMR/2023/v35i125034

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99397>

Case Study

Received: 22/02/2023

Accepted: 25/04/2023

Published: 26/04/2023

ABSTRACT

Introduction: Breastmilk jaundice is a benign often undiagnosed condition, occurring in healthy, term exclusively breastfed neonates. It is characterized by unconjugated hyperbilirubinaemia usually appearing from the second week of life, but sometimes within the first week of life, and persisting for more than two weeks. It's specific aetiology is multifactorial and remains a topic of debate. Controversies also exist regarding the best treatment option. The American Academy of Paediatricians recommends continuation of breastfeeding with temporary addition of formula.

Aim: This series aims to increase awareness of this condition.

Presentation of Cases: Three cases of unconjugated hyperbilirubinaemia persisting for greater than two weeks, in term, healthy, exclusively breastfed infants in Enugu State, South-East Nigeria are reported. Investigations did not reveal pathological jaundice, and all cases received phototherapy. In the three cases however, the most significant reduction in serum bilirubin occurred after temporary cessation of breastfeeding, with commencement of formula feeds, and subsequent recommencement of breastfeeding.

Discussion: The three cases reported fulfil established criteria for recognizing breastmilk jaundice. This condition is largely underdiagnosed, with no recent reports in literature. The aetiology remains

*Corresponding author: E-mail: isaacasinobi@esu.edu.ng, isaacasinobi@gmail.com;

a subject of research. However, several genetic and environmental factors have been suggested. Though the total bilirubin did not reach the American Academy of Paediatrics thresholds for phototherapy in any of the cases, this treatment modality was still used. In addition, in variance to recommendations of the American Academy of Paediatricians, all cases were treated by breastfeeding cessation instead of mixed feeding.

Conclusion: Breastmilk jaundice is largely not remembered as a cause of prolonged jaundice. Its awareness and treatment would greatly reduce both parental anxiety and unnecessary treatment given to babies.

Keywords: Benign; exclusive breastfeeding; hyperbilirubinaemia; persistent.

1. INTRODUCTION

Breastmilk jaundice is a condition characterized by unconjugated hyperbilirubinaemia. It occurs in term and otherwise healthy, exclusively breastfed infants and often goes undiagnosed [1,2]. This benign condition is largely a diagnosis of exclusion [1,3]. Despite its benign nature, it still poses a source of anxiety and guilt, especially among first-time mothers [4]. Because it is often not diagnosed, its incidence may be on the increase [4]. Incidence rates as high as 36% have been reported in the literature [5]. Although widely studied, its aetiology remains unknown [2,4]. Affected babies are characteristically healthy, with jaundice usually appearing from the second week of life, but sometimes within the first week [2,3]. Also, jaundice classically persists for more than two weeks with total serum bilirubin within the range of 13 -15mg/dl [2]. Controversies exist regarding the best treatment option for this condition [5,6,7]. Approaches include continuing breastfeeding while monitoring bilirubin levels; temporarily withholding breastfeeding and substituting formula; and alternating breastfeeding and formula [4]. The American Academy of Paediatricians however recommends that breastfeeding should not be discontinued [8]. This is because resumption of breastfeeding after its cessation often does not occur, exposing the child to later feeding difficulties [9].

We report three cases, over a nine-month period, of unconjugated hyperbilirubinaemia persisting for greater than two weeks amongst term, otherwise healthy exclusively breastfed infants in Enugu State, South East Nigeria.

2. PRESENTATION OF CASES

2.1 Case One

ED, a 26-day-old female, presented with yellowness of the eyes noticed on the 6th day of life. There was no fever, abnormal body

movement, reduced activity or excessive/abnormal cry. She was delivered vaginally at term with a birth weight of 3.4kg to a 36-year-old teacher. Antenatal care was commenced at a gestational age of 14 weeks. Booking investigations (VDRL, HBsAg, RVS) were all negative. She was regular with her antenatal visits and compliant with her medications. There were no adverse events during pregnancy. Both mother and father were blood group O-ve. She is the third in a family with two other children. None of her siblings were exclusively breastfed and none had jaundice during the neonatal period. ED was exclusively breastfed. At onset of the jaundice, she was exposed to early morning sunlight for two days before presenting to the first hospital. The total serum bilirubin was 272 μ mol/l (15.9mg/dl) with an unconjugated fraction of 236 μ mol/l (13.8mg/dl). She subsequently received three days of phototherapy before being discharged on the 12th day of life with a total and conjugated serum bilirubin of 231.2 μ mol/l (13.2mg/dl) and 188.7 μ mol/l (11mg/dl) respectively. On the 16th day of life, she presented to the second hospital following persistence of yellowness of the eyes. Total serum bilirubin was 238 μ mol/l (13.9mg/dl) with an unconjugated fraction of 209.1 μ mol/l (12.2mg/dl). Her complete blood count showed no evidence of infection. Other investigations carried out included blood group, blood film for malaria parasite and cell morphology, glucose-6-phosphate dehydrogenase (G6PD) assay, coombe's test, liver function test and serum protein. Admitting diagnosis was Prolonged Hyperbilirubinaemia ? Cause. She was then placed on phototherapy for four days during which time the serum bilirubin gradually reduced. She was discharged on the 19th day of life with a total and conjugated serum bilirubin of 180.2 μ mol/l (10.5mg/dl) and 149.6 μ mol/l (8.7mg/dl) respectively. However, on the 22nd day of life, the mother noticed the skin was becoming yellow again. She, however did not present to the hospital on the advice of her

relative. On the 26th day of life, she presented to our facility after being counselled about the complications of jaundice by a friend. At presentation, she was healthy looking and moderately icteric. There were no significant findings on systemic examination. Her weight was 4.8kg. Total serum bilirubin was 255 μ mol/l (14.9mg/dl) with an unconjugated fraction of 236 μ mol/l (13.8mg/dl). Her blood group was O-ve, Hb 14.6g/dl and the G6PD assay showed normal activity. Her complete blood count showed no evidence of infection. The blood film also showed no malaria parasites. She was commenced on phototherapy and discharged after two days with a total and conjugated serum bilirubin of 219 μ mol/l (12.8mg/dl) and 180.6 μ mol/l (10.5mg/dl), respectively. A diagnosis of Breastmilk Jaundice was then made. The mother was advised to stop breastfeeding and use only formula milk for two days, then recommence breastfeeding and come for a repeat serum bilirubin in four days. On follow-up, ED was only mildly icteric and had a total and unconjugated serum bilirubin of 140.5 μ mol/l (8.2mg/dl) and 122.6 μ mol/l (7.2mg/dl) respectively. She came for follow-up a week later and was feeding well with good activity and no clinically demonstrable jaundice.

2.2 Case Two

NU, a one month old female infant presented with yellowness of the eyes since the 3rd day of life. For this, she was exposed to early morning sunlight for fourteen days with no significant improvement before it was discontinued. No laboratory investigations were carried out. She was delivered through an emergency caesarean section following poor progress of labour. Her birth weight was 3.8kg. Antenatal care commenced from the 11th week of gestation, and booking investigations were normal. Pregnancy was uneventful and she was carried to term. There were no significant peri-partum events. Mother and father's blood groups were A+ve and O-ve respectively. She is the first and only child of the family. She was put to breast within an hour of birth and was exclusively breastfed with adequate lactation. On examination she was moderately jaundiced and her weight was 4.6kg. All other examination findings were unremarkable. At presentation, the total serum bilirubin was 171.2 μ mol/l (10mg/dl), with an unconjugated fraction of 154.3 μ mol/l (9mg/l). Other tests done included

complete blood count, blood film for malaria parasite and cell morphology, blood group, liver function test, coombs (direct and indirect), serum protein and G6PG assay. These were all normal. Her haemoglobin level was 15.6mg/dl. Admitting diagnosis was Prolonged Hyperbilirubinaemia ? Cause.

After receiving phototherapy for four days, the total serum bilirubin reduced to 156.4 μ mol/l (9mg/dl) with an unconjugated fraction of 134.8 μ mol/l (7.9mg/dl). A diagnosis of Breastmilk Jaundice was then made and she was then discharged. Mother was asked to stop breastfeeding, give formula milk only for two days, and then have serum bilirubin repeated in four days. On follow up, she was breastfeeding well and the repeat serum bilirubin showed a total of 126.7 μ mol/l (7.4mg/dl) with an unconjugated fraction of 103.6 μ mol/l (6.1mg/dl).

2.3 Case Three

IA presented on the 14th day of life with yellowness of the skin noticed on the fifth day of life. She was delivered at term, had a birth weight of 2.6kg and weighed 3.15kg at presentation. She was otherwise healthy. She was delivered at term through an elective caesarean section on account of a previous maternal caesarean section. She cried well at birth and there were no significant peri-partum events. Pregnancy was booked at the 17th week of gestation. All booking parameters were normal, mother was both compliant with routine antenatal care medication and regular with antenatal visits. There were no illnesses or adverse events during pregnancy. Mothers blood group was B+ve, while the father's was unknown. She is the second in a family with two children. There was no history of jaundice in the elder sibling. She received water for the first four hours of life and was subsequently exclusively breastfed. At presentation, she was active, afebrile, but moderately icteric. Both general and systemic examination revealed no significant findings. Total serum bilirubin was 251 μ mol/l (14.8mg/dl) with an unconjugated fraction of 189 μ mol/l (11.5mg/dl), her blood group was B+ve and Hb 15.4g/dl. All other investigations (G6PD assay, complete blood count, reticulocyte count, blood film for malaria parasite and cell morphology and liver function test) were within normal limits. The admitting diagnosis was Prolonged Hyperbilirubinaemia ? Cause.

Table 1. Clinical characteristics of cases

	Age at Presentation	Jaundice Onset	Other Symptoms	Gestation	Antenatal Care	Illnesses during Pregnancy	Jaundice in Siblings	Mode of Feeding	Birth Wt	Mode of treatment	Duration of Jaundice
Case 1	26 days	6 th day of life	None	Term	Adequate	Nil	Nil	Exclusive Breast-feeding	4.8kg	Phototherapy Temporary cessation of Breast-feeding	>2 weeks
Case 2	28 days	3 rd day of life	None	Term	Adequate	Nil	Nil	Exclusive Breast-feeding	4.6kg	Phototherapy Temporary cessation of Breast-feeding	>2 weeks
Case 3	14 days	5 th day of life	None	Term	Adequate	Nil	Nil	Exclusive Breast-feeding	3.15kg	Phototherapy Temporary cessation of Breast-feeding	>2 weeks

Table 2. Laboratory parameters of cases

	Parents Blood Group	Peak Total Serum Bilirubin	Total Serum Bilirubin at Presentation (TSB)	Haemoglobin concentration	Other Investigations	Discharge Serum Bilirubin
Case 1	Mother O-ve Father O-ve	15.9mg/dl	14.9mg/dl	14.6g/dl	Normal	8.2mg/dl
Case 2	Mother A+ve Father O-ve	10mg/dl	10mg/dl	15.6mg/dl	Normal	9mg/dl
Case 3	Mother B+ve Father unknown	14.8mg/dl	14.8mg/dl	15.4g/dl	Normal	9.7mg/dl

Serial serum bilirubin values over the next few days showed a slow but gradual reduction. On the 21st day of life, after receiving phototherapy for seven days, her total serum bilirubin was 165.2 μ mol/l (9.7mg/dl) with an unconjugated fraction of 160.6 μ mol/l (9.4mg/dl). The diagnosis was then modified to Breastmilk Jaundice. She was discharged home and asked to stop breastfeeding, give formula milk only for two days, and then have serum bilirubin repeated in three days. On follow up, she was healthy looking with a weight of 3.7kg and a total serum bilirubin was 90.0 μ mol/l (5.3mg/dl) with an unconjugated fraction of 70.4 μ mol/l (4.1mg/dl).

A summary of the clinical and laboratory parameters for the three cases is shown in Tables 1 and 2.

3. DISCUSSION

Breast milk jaundice often goes unnoticed and is thus largely underdiagnosed and under-reported. We have reported three cases occurring in South-East Nigeria, which from our search has hitherto not been reported in literature. In the first case, the diagnosis was made only after the patient was exposed to several investigations and interventions. In the other two cases, the diagnosis was made following recognition of the slow and poor response to phototherapy, and after none of the investigations carried out pointed to a specific diagnosis. Breastmilk jaundice is not usually reported in studies as a cause of jaundice [10,11,12]. As such, its actual prevalence is not widely reported. Recent case reports are lacking, with available reports spanning three to six decades ago [10,13,14,15].

Amongst the cases reported, there was similarity in the onset of yellowness of the eyes and skin, with onsets on the 6th, 8th and 5th days of life. However, common to all was the persistence for more than two weeks. This is consistent with standard definitions stating this condition starts within the second and third weeks and persists beyond three weeks of life [1,11]. Unlike healthy formula fed infants who do not have hyperbilirubinaemia in the third week of life, approximately one third and two third of all breastfed infants have jaundice and hyperbilirubinaemia respectively in the third week of life [1,16]. Breastmilk jaundice is also reported to occur in healthy term babies [4]. This is in contrast to breast feeding jaundice which is seen in lean underfed babies, and which results from insufficient breastmilk, with a resultant reduction

in intestinal emptying and thus an increase in enterohepatic circulation [17,18]. The three cases we reported were all healthy term neonates with normal birth weight.

Only one of the cases had a serum bilirubin above 15mg/dl. It has been recommended that serum bilirubin levels within this level are less likely to be due to breastmilk jaundice alone, and that investigations should be carried out to determine other causes of pathological jaundice [1,11]. In cases one and two, all available investigations were carried out to establish the aetiology and rule out other causes of pathological jaundice. In case three however, a coombs test was not done. In this case, the reticulocyte count did not show evidence of increased haemolysis, which would point towards increased haemolysis as a possible cause for the hyperbilirubinaemia. The aetiology of breast milk jaundice is still a subject of research, as no clear aetiology has been described [5]. The role of several genetic and environmental factors has been suggested [2,4,5,11,15,19]. These include: reduced uptake (mutation in the solute carrier organic anion transporter protein *SLCO1B1*); reduction in bilirubin conjugation (defective uridine diphosphate-glucuronyl transferase (*UGT1A1*) activity, increased concentrations of nonesterified free fatty acids; increased pregnane-3- α -20 beta-diol in breast milk); increased enterohepatic circulation (delayed establishment of enteric flora, increased beta glucuronidase activity in breast milk and reduced intestinal motility from increased levels of epidermal growth factor (EGF) in breast milk); and reduced uptake, metabolism, and excretion of bilirubin from increased levels of inflammatory cytokines in human milk [2,4,11,19,20].

The diagnosis of breastmilk jaundice is one of exclusion [3]. Further investigations including those for genetic causes of persistent jaundice such as Crigler-Najjar syndrome and Gilberts syndromes were not carried out due to lack of facilities [21]. In addition, thyroid function tests should have been carried out, but these were omitted [22]. Values within the normal range for these investigations would have further strengthened the diagnosis of breastmilk jaundice. In all three cases reported, breastfeeding was discontinued and formula feeds given for 48 hours. Breast feeding was then successfully recommenced subsequently. The American Academy of Paediatricians however recommends that unless serum bilirubin exceeds 340 μ mol/l (20mg/dl), breastfeeding

should not be discontinued due to concerns about its subsequent successful recommencement [8]. Below this value, formula feeds should be introduced while breastfeeding continues [7].

4. CONCLUSION

We have reported three cases of breastmilk jaundice amongst otherwise healthy, term exclusively breastfed babies with similar presentation and treatment. Breastmilk jaundice is largely not remembered as a cause of prolonged jaundice in our environment, and is thus grossly under reported. Physicians should be aware of this condition and its treatment, especially after excluding other cases of pathological jaundice. This would greatly reduce both parental anxiety and unnecessary treatment given to babies.

CONSENT

The author declares that written informed consent was obtained from the patient.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Preer GI, Philipp BL. Understanding and managing breast milk jaundice. *Arch Dis Child Fetal Neonatal* Ed. 2011;96:F461-66. DOI: 10.1136/adc.2010.184416
2. Soldi A, Tonetto P, Varalda A, Bertino E. Neonatal jaundice and human milk. *J Matern Fetal Neonatal Med*. 2011;24(1):85-7. DOI: 10.3109/14767058.2011.607612. PMID: 21942599.
3. Bratton S, Cantu RM, Stern M. Breast Milk Jaundice. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2023. Accessed: 24 April, 2023. Available: <https://www.ncbi.nlm.nih.gov/books/NBK537334/>
4. Brooten D, Brown L, Hollingsworth A, Tanis J, Bakewell-Sachs S. Breast-milk jaundice. *J Obstet Gynecol Neonatal Nurs*. 1985;14(3):220-3. Available: doi.org/10.1111/j.1552-6909.1985.tb02230.x
5. Alonso EM, Whittington PF, Whittington SH, Rivard WA, Given G. Enterohepatic circulation of unconjugated bilirubin in rats fed with human milk. *J Pediatr*. 1991;118:425-30. DOI: 10.1016/s0022-3476(05)82162-6
6. Prameela KK. Breastfeeding during breast milk jaundice - A pathophysiological perspective. *Med J Malaysia*. 2019;74(6):527-533. PMID: 31929480.
7. Auerbach KG, Gartner LM. Breastfeeding and human milk: their association with jaundice in the neonate. *Clin Perinatol*. 1987;14(1):89-107. Available: [http://dx.doi.org/10.1016/S0095-5108\(18\)30783-8](http://dx.doi.org/10.1016/S0095-5108(18)30783-8). PMID: 3549117.
8. [Guideline] American Academy of Pediatrics Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics*. 2004;114(1):297-316. DOI: 10.1542/peds.114.1.297
9. Ndu I, Ekwochi U, Osuorah C, Chinawa J, Asinobi I, Eze J, et al. The knowledge and practice of forced-feeding among mothers and caregivers in Enugu, South East Nigeria. *Int J Trop Dis Health*. 2016;11:1-7. DOI: 10.9734/IJTDH/2016/19826
10. Tokunaga A. Neonatal jaundice induced by mother's milk. *Nihon Rinsho*. 1985;43(8):1671-7. Japanese.
11. Seneadza N, Insaideo G, Boye H, Ani-Amponsah M, Leung T, Meek J, Enweronu-Laryea C. Neonatal jaundice in Ghanaian children: Assessing maternal knowledge, attitude, and perceptions. *PLoS One*. 2022;17(3):e0264694. DOI: 10.1371/journal.pone.0264694. PMID: 35239710; PMCID: PMC8893663.
12. Mitra S, Rennie J. Neonatal jaundice: aetiology, diagnosis and treatment. *Br J Hosp Med*. 2017;78(12):699-704. DOI: 10.12968/hmed.2017.78.12.699. PMID: 29240507.
13. Sivelli V, Di Battista C. 11 cases of neonatal jaundice caused by pregnanediol in the mother's milk. *Clin Ter*. 1969;51(2):131-6. Italian. PMID: 5365823.
14. Grgurić J, Zimolo A, Bacić V. Jaundice in an infant due to maintenance on mother's milk. *Lijec Vjesn*. 1974 ;96(12):745-8. Croatian. PMID: 4456105.

15. Lupu V, Dascălu D, Barbu M, Horvath I. Icter cu bilirubinemie indirectă prin lapte matern [Jaundice with indirect bilirubinemia via the mother's milk]. *Pediatrie (Bucur)*. 1992;41(1):67-70. Romanian. Romanian. PMID: 1322729.
16. Ullah S, Rahman K, Hedayati M. Hyperbilirubinemia in Neonates: Types, causes, clinical examinations, preventive measures and treatments: A narrative review article. *Iran J Public Health*. 2016;45(5):558-68. PMID: 27398328; PMCID: PMC4935699
17. Siroosbakht S, Aminian AR, Rezakhaniha B. Risk factors of early breastfeeding jaundice: How can gynecologists and nurses be effective in reducing it?. *J Compr Ped*. 2020;11(3):e103578. Available:<https://doi.org/10.5812/compreped.103578>.
18. Kemper AR, Newman TB, Slaughter JL, Maisels MJ, Watchko JF, Downs SM, et al. Clinical practice guideline revision: Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics*. 2022 Sep 1;150(3):e2022058859. DOI: 10.1542/peds.2022-058859. PMID: 35927462.
19. Kumral A, Ozkan H, Duman N, Yesilirmak DC, Islekel H, Ozalp Y. Breast milk jaundice correlates with high levels of epidermal growth factor. *Pediatr Res* 2009;66:218–21. DOI: 10.1203/PDR.0b013e3181ac4a30
20. Ince Z, Coban A, Peker I, Can G. Breast milk beta-glucuronidase and prolonged jaundice in the neonate. *Acta Paediatr*. 1995 Mar;84(3):237-9. DOI: 10.1111/j.1651-2227.1995.tb13621.x. PMID: 7780242.
21. Memon N, Weinberger B, Hegyi T, Aleeksunes LM. Inherited disorders of bilirubin clearance. *Pediatr Res*. 2016; 79: 378–386. Available:<https://doi.org/10.1038/pr.2015.247>.
22. Kayıran SM, Gürakan B. Correlation of third day TSH and thyroxine values with bilirubin levels detected by a neonatal screening system. *Medical journal of Bakirköy*. 2010;6:117-120.

© 2023 Nwabueze; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/99397>