



Soil and Nutrient Losses under Cultivated Bush and Climbing Beans on Terraced Humid Highland Slopes of Southwestern Uganda

G. Gabiri^{1*}, J. A. Obando¹, M. M. Tenywa², J. G. Majaliwa³, C. L. Kizza²,
A. Zizinga³ and R. Buruchara⁴

¹Department of Geography, Kenyatta University, P.O.Box 43844, 00100, Nairobi, Kenya.

²Department of Agricultural Production, Makerere University, P.O.Box 7062, Kampala, Uganda.

³Department of Environmental Sciences, Makerere University, P.O.Box 7062, Kampala, Uganda.

⁴International Centre Tropical Agriculture (CIAT), P.O.Box 6247, Kampala, Uganda.

Authors' contributions

This work was carried out in collaboration with all authors. Author GG initiated the research and conducted it. Authors JAO, MMT and JGM designed the study and supervised the research. Authors GG and AZ managed the literature searches and reviewed the manuscript drafts and participated in the data collection. Author CLK analysed the soil properties in relation to the field and statistical analyses. Author RB provided part of the technical input in the overall research project. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JSRR/2015/18113

Editor(s):

(1) Corina Carranca, National Institute of Agricultural Research and Veterinary (INIAV), Research Unit of Environment and Natural Resources (UIARN), Portugal.

Reviewers:

(1) Anonymous, Universidad Del Mar Campus Puerto Escondido, Chile.

(2) Anonymous, Tribhuvan University, Nepal.

(3) Vineeta Rai, Dept of Biosciences and Bioengineering, IIT Bombay, India.

Complete Peer review History: <http://sciencedomain.org/review-history/9937>

Original Research Article

Received 4th April 2015

Accepted 13th June 2015

Published 29th June 2015

ABSTRACT

Highland regions characterized with terraced slopes in East Africa are faced with serious soil and nutrient losses that have affected agricultural productivity. However, limited studies have analysed the soil and nutrient losses on terraced slopes of these humid highland regions. Therefore, this study was conducted to assess the effects of climbing beans over the traditional bush beans' cultivation on soil erosion control in Bufundi sub catchment, Kabale District, South-western Uganda. Runoff trap approach was employed to assess erosion and the sites were characterized

*Corresponding author: Email: geofreygabiri@gmail.com;

for soil physico-chemical properties. Terraced slopes planted with climbing beans showed significantly lower ($P < 0.05$) annual runoff and soil loss of $36 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$ and $121 \text{ kg ha}^{-1} \text{ yr}^{-1}$ over bush beans, $248 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$ and $548 \text{ kg ha}^{-1} \text{ yr}^{-1}$, respectively. Annual soil nitrogen, phosphorus and potassium losses were significantly higher from bush beans ($1.87 \text{ kg N ha}^{-1} \text{ yr}^{-1}$; $0.6 \text{ kg P ha}^{-1} \text{ yr}^{-1}$ and $0.12 \text{ kg K ha}^{-1} \text{ yr}^{-1}$) over climbing beans ($0.49 \text{ kg N ha}^{-1} \text{ yr}^{-1}$; $0.1 \text{ kg P ha}^{-1} \text{ yr}^{-1}$; and $0.03 \text{ kg K ha}^{-1} \text{ yr}^{-1}$). The total economic value of bush beans due to erosion was higher ($75424 \text{ Ug shillings/ha}$) than that of climbing beans ($15597 \text{ Ug shillings/ha}$). Generally, runoff, soil and nutrient losses were very low from both bean types. However, climbing beans were superior to bush beans in controlling runoff, soil and nutrient losses confirming their effectiveness in controlling erosion down the catchment.

Keywords: Economic value; erosion; macronutrients; hydraulic conductivity; soil texture.

1. INTRODUCTION

Sustaining agricultural productivity and livelihoods in the tropical humid highland ecosystems is increasingly becoming an enormous challenge [1,2,3]. The fragility of the slopes coupled with the increasing population and limited livelihood opportunities has led to increased soil erosion [4,5,6,7]. A compendium of literature reveals high soil erosion in the highlands [7,8,9,10,11,12,13] emphasizing soil and nutrient losses as a key land degradation challenge in most of the humid high land ecosystems of East Africa; that has resulted in negative effects on soil fertility. Soil and nutrient losses have been estimated to affect 50% of the total arable land productivity area in the East African humid highlands [14].

In the Ugandan perspective, soil and nutrient losses have been reported to be the most severe and extensive form of land degradation in the hilly and mountainous farming systems leading to loss of nutrients, and contributing to the declining agricultural productivity [15,16,17,18]. According to [19], Uganda's highland regions experience soil loss of more than 30 t ha^{-1} on an annual basis. Soil and nutrient losses impose a series of onsite and offsite environmental effects, which include eutrophication and excess turbidity to the drainage network and reservoirs [20]. These impacts increase the maintenance costs for terraces and drainage channels/streams and filling of ephemeral gullies as well as the cost of replacing the lost nutrients mainly nitrogen (N) and phosphorus (P) [21].

According to [12], efforts on soil and nutrient loss control in the humid highlands started back during the British colonial period, when terraces and other conservation measures were force fully introduced in East Africa including SW Uganda. Despite these conservation measures,

challenges of soil and nutrient losses have persisted ostensibly due to poor farming practices, expensive technologies, high population growth, etc. Therefore, a search for supporting technologies has been on-going, together with research on innovative methods that can enhance food productivity and environmental management [7,10,13,22,23]. One such venture is the Sub Saharan Africa challenge program with the concept of Integrated Agricultural Research for Development (IAR4D) developed and implemented by Forum for Agriculture Research in Africa (FARA) [24] which has a main focus to improve farmers' productivity without compromising the environment [25]. The IAR4D innovation platform has spearheaded bean studies in the Lake Kivu humid highlands in south-western (SW) Uganda. Planting of climbing beans in pure stands was introduced within the south-west highlands and on the slopes of Mountain Elgon so as to increase productivity [26]. Although the land under bean cultivation constitutes 20% of the total land area in Uganda, still there is a lack of information on the effectiveness of climbing beans in controlling the soil and nutrient losses in comparison with the traditional bush beans.

Despite the fact that bean studies have been conducted in the East African humid highlands with the example of SW Uganda, these studies have largely centered on improving yields, improving resistance to drought, pests and diseases, low soil fertility, less cooking time varieties and improving nutritional value [27,28, 29,30]. There has been limited attention on the contribution of these different bean types in controlling soil erosion and studies that have explored erosion in the cropping systems have largely remained broad on annual crops [31,32,33].

Therefore, the main objective of the study was to assess soil and nutrient loss variability on the

terraced slopes cropped with climbing and bush beans within the Sub Saharan Africa challenge program innovation platforms (IPs) in SW Uganda. The study was mainly comparative between these two bean types and limited to erosion assessment and its economic implications. This study did not consider crop yields, pests and diseases since these aspects have already been described by several authors in the field of bean production programs [27,29, 34,35,36].

2. MATERIALS AND METHODS

2.1 Description of the Study Area

The study was undertaken in a humid highland region of Bufundi sub catchment located in Kabale district, south-western Uganda, between latitudes 1° and 1° 34"S and longitudes 29° 18"E and 30° 9"E (Fig. 1). The Bufundi highlands are part of the East African pre-Holocene volcanics bordering Lake Bunyonyi [37]. The terrain is




Fig. 1. Map of Bufundi sub catchment

dominated by hills coupled with degrading terraces and valleys with most slopes ranging between 12 to 50% but may go as high as 80% [38]. The soils in Bufundi are typically Luvisols, Histosols and Ferrasols (Fig. 2) with declining fertility due to continuous cultivation and susceptible to extreme soil erosion [38]. Histosols are very rich in organic matter (>14%) with incompletely decomposed plant residues, with or without clay, sand and silt [39]. In these organic layers, the infiltration increases, hence less susceptible to soil erosion [40]. Luvisols have strong accumulation of clay in the B-horizon with high cation exchange capacity and moderate weathering and Luvisols with high silt content are susceptible to structure deterioration if tilled in wet conditions and on steep slopes, they are easily eroded [39,41]. Ferrasols are highly weathered soils rich in sesquioxide clays and with low cation exchange capacities [39]. Ferrasols have distinctive soil structure with a weak micro-aggregates, but strong macro-aggregates hence this structure makes Ferrasols fairly porous and less susceptible to erosion than other soils in the tropics [42]. The catchment has a bimodal rainfall with average annual rainfall between 900 mm to 2200 mm with a mean annual temperature of 16.7°C [43]. The major land uses in the sub-catchment include small scale farmland and woodlots (*Eucalyptus*).

2.2 Runoff, Soil and Nutrient Losses Determination

In order to determine runoff, soil and nutrient losses, data collection was conducted during the long rain season (February- July, 2012) and short rain season (September - December, 2012). Three sites with slope elevation ranging between 23% and 29% at the mid-slope of the hill at least 800 m apart were identified for the study. The soil classification at these sites was Acric Ferrasols. Within each site, six erosion plots were constructed within the terraces with each plot measuring 2 m by 9 m to measure runoff using a pipe sampler [45]. Erosion plots were sealed off using several iron sheets measuring 3 m long and 0.3 m high. From the six erosion plots installed in each site, two treatments (bush beans and climbing beans) were planted at a spacing of 50 cm * 25 cm and 50 cm * 20 cm, respectively in each three erosion plots. Farmers managed the plots like any other field in order to represent the management across the catchment A small container was tied at the end of the erosion trap to capture 4% of the total runoff through the slit. However, each plot was calibrated with a known volume of water to establish its runoff transmission coefficient which was later used to determine the actual




Fig. 2. The major soil type of Bufundi sub catchment. Source: [44]

total plot runoff [46]. Total runoff (water and sediments) collected in the container and measured using a measuring cylinder following a rainfall event. Total runoff was then put in clean plastic bottles and periodically delivered to the Department of Soil Science laboratory at Makerere University for subsequent analyses. Event runoff was multiplied by the coefficient of transmission to get the plot runoff per rainstorm event, and then extrapolated to a hectare basis, by multiplying the obtained value with 555.6. Seasonal runoff loss was obtained by summing up the different runoffs registered for different rainfall events and the two seasonal runoff collections were eventually summed up to obtain the annual loss per ha.

In the laboratory, each runoff sample was homogenized and divided into two portions. The first portion was filtered using filter paper and the filtrate was oven dried at 105°C for 24 hours for sediment determination. The amount of soil loss in the plot was obtained by multiplying the sediment concentration by runoff and transmission coefficient and then expressed on ha basis. Seasonal soil loss was obtained by summing up all the seasonal collections and then annual soil loss obtained by summing up the two seasonal soil losses. The remaining portion of the runoff sample was also filtered for first portion and the filtrate air dried and analyzed for macro nutrients: N, P and potassium (K) using the procedures from [47]. Seasonal nutrient losses were determined by summing up all the seasonal collections and then annual nutrient loss obtained by summing up the two seasonal nutrient losses.

2.3 Soil Physical and Chemical Characterization

In order to characterize soil physical and chemical properties of the study sites, soil samples were taken within each erosion plot to determine saturated hydraulic conductivity using constant head method and bulk density using core oven drying method [48] at the end of each bean growing season. Two soil samples were collected using soil core measuring 5 cm long by 5cm internal diameter in each soil depths of 0-15 cm and 15 -30 cm. Two composite soil samples from each soil depth at each site and treatment were taken at planting and at harvesting to assess the characteristics of soil chemical properties under the two different bean types in the study area. Soil organic carbon (SOC), total N, available P and exchangeable bases: K,

sodium (Na), calcium and magnesium (Mg) were analyzed using [47].

2.4 Rainfall Measurement

In order to measure the amount of rainfall received at each study site, a manual recording rain gauge was installed. The amount of rainfall events was summed and a number of rainy days were manually recorded. Annual rainfall was determined by summing up the two seasonal rainfalls.

2.5 Data Analysis

Data were entered into Microsoft Excel and imported into Genstat Discovery Version 13 for statistical analysis. One and two way analysis of Variance (ANOVA) were performed in order to ascertain whether there are differences in the means of runoff, soil and nutrient losses between the two bean types and the experimental sites.

3. RESULTS OF THE STUDY

3.1 Rainfall Characteristics within the Study Area

Mean monthly rainfall ranged from 0.5 mm to 591.8 mm and monthly rainy days ranged from 1 to 11 days. The highest mean monthly rainfall was observed in the long rainy season (February to July) coupled with the highest number of rainy days (Fig. 3). The rainfall and rainy day trends are similar to those reported by [49] in the southwestern region.

3.2 Soil Physic-chemical Characteristics

Soils cropped with beans were mainly clay loam, except the top soil for bush beans which was sandy clay loam. However, the clay and sand contents varied significantly ($P<0.05$) among sites (Table 2).

Soil bulk densities and saturated hydraulic conductivity at the top soil and sub soil were moderately low and varied significantly ($P<0.05$) among sites, bean types and sites (Table 2). Climbing bean registered higher saturated hydraulic conductivity than bush beans for all the experimental sites and depths and there was more water movement at a depth of 0-15 cm due to the high soil organic matter content (Table 2).

The soils in the study area had sufficient nutrients required for normal bean growth (Table

1) though with slight variations between the bean types and depth. For example, soil organic matter ranged between 2.84% and 3.80% with the highest observed in climbing beans at the top soil (0-15 cm) while the least in bush beans at the sub soil (15-30 cm) (Table 1). Soil organic matter was significantly different ($P < 0.05$) between depths, bean types and sites (Table 2). The highest soil organic matter content was registered in climbing beans plots at a depth of 0-15 cm (Table 2).

Nitrogen and available P did not vary significantly with either bean types or soil depths, except among sites where N and P were significantly different ($P < 0.01$) (Table 3). However, mean available P was higher in climbing beans and at a depth of 0-15 cm (Table 1).

Exchangeable K was also significantly higher ($P < 0.05$) in climbing beans than bush beans (Table 1). When compared among the experimental sites, exchangeable K, Ca and Mg varied significantly ($P < 0.05$) between depths, bean types and sites (Table 2). Generally, all parameters were above their respective critical values.

3.3 Runoff, Soil and Nutrient Losses from Climbing and Bush Beans

The mean annual runoff varied significantly ($P < 0.05$) between climbing beans ($36 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$) and bush beans- ($248 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$) (Fig. 4). Runoff generation from each bean growing season was also significantly different between climbing and bush beans at $P < 0.05$ (Fig. 5). Bush beans contributed with the highest seasonal runoff (363 and $161 \text{ m}^3 \text{ ha}^{-1} \text{ season}^{-1}$) than climbing beans (55 and $22 \text{ m}^3 \text{ ha}^{-1} \text{ season}^{-1}$) during the long and short rainy seasons, respectively (Fig. 5). The percentage rainfall that ended up as runoff ranged from 3.6% to 21.9% in the short and long rainy season periods, respectively (Fig. 6). The low runoff to rainfall ratio in the short over the long rainy season is likely to be due to low runoff observed in the short rainy season from both bean types. The low percent rainfall that ended up as runoff from climbing beans over bush beans is due to the ground cover characteristics noted by [30] and other several authors [50,51,52,53] who have concluded that the higher the canopy/ground cover, the lower runoff coefficient due to increased infiltration rate. Therefore, climbing beans' ground cover reduced runoff down the catchment, hence translating into low runoff coefficient.




Fig. 3. Mean monthly precipitation and total rain days in the sub-catchment in 2012

Table 1. Chemical properties of soils under climbing and bush beans

Depth (cm)	Bean type	pH	N	SOM	Av. P	Exch. K	Ca	Mg	Na
			%		mg/kg	cmol(+)/kg			
0-15 top	Climbing beans	5.30	0.18	3.80	20.30	0.62	4.27	2.04	0.05
	Bush beans	5.20	0.15	3.60	16.90	0.50	4.18	2.20	0.05
15-30	Climbing beans	5.20	0.15	3.39	16.60	0.56	3.96	2.00	0.05
	Bush beans	5.00	0.15	2.84	9.10	0.42	3.68	1.90	0.05
Bean type (LSD _{0.05})		ns	ns	ns	ns	0.12	ns	ns	ns
Depth (LSD _{0.05})		ns	ns	0.58	ns	ns	ns	ns	ns

ns = mean not significant at P≥0.05. SOM= Soil Organic Matter; Av. P= Available Phosphorus; Exch. K= Exchangeable Potassium




Fig. 4. Annual runoff from terraced slopes cropped with climbing and bush beans




Fig. 5. Runoff from the different rain seasons during the study period

Annual soil loss varied significantly ($P < 0.05$) from 121 kg ha⁻¹ yr⁻¹ in climbing beans to 548 kg ha⁻¹ yr⁻¹ in bush beans (Fig. 7). Seasonal soil loss also varied significantly between the two bean

types ($P < 0.05$). Bush beans contributed the highest amounts of soil loss in each bean cultivation season: 1008 kg soil loss ha⁻¹ season⁻¹ for long rainy season period and 311 kg soil loss ha⁻¹ season⁻¹ for short rainy season

period as compared to climbing beans which registered 413 kg soil loss ha⁻¹ season⁻¹ for long rainy season period and 29 kg soil loss ha⁻¹ season⁻¹ for short rainy season period (Fig. 8).




Fig. 6. Percentage runoff to rainfall




Fig. 7. Annual soil loss from terraced slopes cropped with climbing and bush beans

Table 2. Interactions of soil physico-chemical properties with sites, bean types and soil depths

Depth (cm)	Bean type	Site	Clay	Sand	SOM	N	Av.P	BD	Ksat	Ca	Mg	Na	K	pH
0-15			%				Mg/kg	g/cm ³	cm/hr.	Cmol(+)/kg				
0-15	Climbing beans	1	22.0	48.0	2.83	0.11	2.2	1.41	4.33	2.30	1.17	0.04	0.24	4.81
		2	20.7	45.7	3.89	0.19	23.0	1.42	11.03	3.80	1.68	0.03	0.53	5.23
		3	28.7	35.3	4.07	0.17	35.6	1.34	7.33	6.72	3.28	0.07	1.10	5.86
	Bush beans	1	22.0	46.0	2.29	0.15	4.5	1.41	2.84	2.70	1.39	0.05	0.41	4.74
		2	18.7	57.3	3.81	0.19	30.9	1.45	8.23	4.14	2.22	0.04	0.61	4.96
		3	27.7	34.7	5.29	0.21	15.3	1.35	10.52	5.70	3.08	0.06	0.47	5.82
15-30	Climbing beans	1	22.0	49.7	2.67	0.12	3.4	1.42	3.85	2.50	1.27	0.04	0.28	5.03
		2	23.7	52.0	3.87	0.14	18.8	1.41	6.89	3.19	1.72	0.04	0.45	4.86
		3	27.3	33.7	3.64	0.20	27.6	1.35	6.05	6.18	3.01	0.06	0.96	5.82
	Bush beans	1	24.0	45.7	1.26	0.13	1.8	1.39	0.32	2.37	1.18	0.04	0.39	4.72
		2	20.7	55.7	3.59	0.15	15.0	1.43	6.28	3.25	1.95	0.06	0.42	4.72
		3	27.7	37.7	3.66	0.18	10.4	1.35	5.65	5.45	2.58	0.05	0.45	5.57
Bean type (<i>P</i> < .05)		ns	ns	ns	ns	ns	ns	<0.05	ns	ns	ns	<0.001	<0.001	
Site (<i>P</i> < .05)			<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	ns	<.001	<0.001
Depth (<i>P</i> < .05)			ns	ns	<0.001	ns	ns	ns	<0.001	<0.05	<0.001	ns	<0.001	<0.05
Bean type*Site (<i>P</i> < .05)			<0.05	<0.05	<0.001	ns	ns	<0.05	<0.001	0.05	<0.001	ns	<0.001	ns
Bean type*depth (<i>P</i> < .05)			ns	ns	ns	ns	ns	ns	ns	ns	<0.05	ns	ns	ns
Bean type*depth*site (<i>P</i> < .05)			ns	ns	ns	ns	ns	ns	<0.001	ns	ns	ns	ns	ns

ns = mean not significant at $P \geq 0.05$. SOM= Soil Organic Matter; BD= Bulk density; Ksat =Saturated hydraulic conductivity; Ca =Calcium; Mg = Magnesium; Na= Sodium

Table 3 shows the annual nutrient losses from climbing and bush beans from the study area. Bush beans contributed significantly to the highest annual N and P losses from both runoff and sediment losses than climbing beans. Potassium loss did not vary significantly from both bean types; however, bush beans had the highest losses than climbing beans. The annual N, P and K losses for the study area were 0.49, 0.1, 0.03 kg ha⁻¹ yr⁻¹ and 1.87, 0.6 and 0.2 kg ha⁻¹ yr⁻¹ from climbing and bush beans, respectively.

Economic valuation for the imposition of monetary value was based on the nutrients losses within the transported soil materials and the calculation was adopted from [54]. The calculation of economic value is the total loss of nutrients (N, P and K) in transported soil material and is converted into the economic value based on the local price of urea, tri super phosphate (TSP), and potassium chloride (KCl). Bush beans had the highest value as compared to climbing beans (Table 4).




Fig. 8. Soil losses for the different rainy seasons during the study period

Table 3. Annual nutrient losses from cultivated climbing beans and bush beans

	Bean type	Nutrient loss (kg/ha-1)		
		N	P	K
Runoff/water	Climbing beans	0.001	0.32	0.24
	Bush beans	0.09	2.09	0.37
Soil/sediment	Climbing beans	0.49	0.10	0.03
	Bush beans	1.87	0.60	0.20
P<0.05 (runoff/water)		<0.05	<0.001	ns
P<0.05 (soil/sediment)		<0.001	ns	ns

ns = mean not significant at P>0.05

Table 4. Total economic value of soil erosion in the catchment

Nutrients	Nutrients losses (kg/ha/yr.)	
	Climbing beans	Bush beans
Nitrogen	0.49	1.96
Phosphorus	0.42	2.69
Potassium	0.27	0.57
Economic value (Ug shilling/ha) of nutrient losses based on price of:		
Urea (46% N)	5337	21304
TSP(0:20:0)	7560	48420
KCL (48% K)	2700	5700
Total economic value (Ug shillings/ha)	15597	75424

Economic value of bush beans is higher than that of climbing beans

4. DISCUSSION OF THE RESULTS

4.1 Soil Physic-chemical Characteristics

The low bulk density and high saturated hydraulic conductivity (Ksat) at the top soil is due to the modification by the rich organic matter at the surface due to littering effect of bean plants. This is in agreement with the earlier findings of [55-58,59] who reported a low bulk density in rich organic soils. The high bulk density in the sub soil is due to the relatively more developed soil structure of mineral soils than organic soils. Climbing beans had lower bulk density and higher Ksat than the bush beans due to the high level of littering from the high vegetation cover and the extensive rooting system characteristic of climbing over the bush beans [27]. This leads to the formation of macro pores as result of decaying effects of the leaves [60] and hence increased infiltration thus leading to a reduction of runoff in the catchment. These changes increase crop yield and environmental protection due to nutrient retention in the long rainy season.

4.2 Runoff, Soil and Nutrient Losses from Climbing and Bush Beans

The relatively high runoff from bush beans could be attributed to the low hydraulic conductivity observed and the limited surface vegetation cover which takes a shorter period than climbing beans whose vegetation cover takes 200 days to stand before maturity [30,34]. That could have acted as runoff breakers hence reducing the impact of rainfall energy to the soil surface [61,62]. According to [63], climbing beans have extensive root systems with greater exploration and mechanical support as well as better ability at binding the soil particles than the bush beans. Thus the climbing beans have better capacity in improving the soil structure and hence reducing the runoff and soil losses than bush beans. The climbing beans have a growth height of 2 m high compared with the 30 cm of bush beans. As well as possessing a well and higher aerated surface cover, the green canopy cover which takes longer (about 6 weeks) before flowering as compared with bush beans which retain a green cover for up to 2 – 3 weeks before flowering [34]. These characteristics aid climbing beans to intercept rainfall and increases infiltration on the soil surface [64] and as a result, reduced runoff and soil losses down the catchment. High runoff from bush beans with limited vegetation cover over climbing beans is in agreement with observations from [51,65,66] under the agricultural lands.

The runoff values from both bean types tallied within the range reported by [67,68] and contrasted with [69,70] who reported high values. However, all these runoff values are very low to the tropical African context under the mountainous agricultural lands due to low rainfall erosivity and high infiltration rates.

The high soil loss from bush beans was due to the high runoff registered by these bean types translating into increased soil losses as also reported by [51,71]. The lower vegetation cover in bush beans than climbing beans [30,34] also caused increased soil losses from bush beans due to the fact that soil particles under bush beans' stands were more exposed to detachment by rain drops, and further transportation along with runoff as also observed by [46,72,73]. The annual soil loss from bush beans and climbing beans are within the range of those reported by [67] and in contrast with [74,75] on mountaneous environments.

The factors that favored runoff could have attributed to the trend in soil loss from these two types of beans. As also reported by [22], low bulk density and high organic matter content resulting into high saturated hydraulic conductivity could have favored low runoff and soil losses from climbing beans. The annual soil loss values in the study area were generally too small as compared to those reported by [70] and below the proxy tolerable limit of $5 \text{ t ha}^{-1} \text{ yr}^{-1}$ used in Uganda. Hence the observed plot scale soil loss values indicated that the situation in catchment has not yet reached the critical levels, as also reported by [72,76,77] under the agricultural crop lands. Appraisals by [78] on erosion in the highland regions of Uganda concluded that soil erosion rates are much less than would be expected even in steep slopes due to the low rainfall erosivity, rainfall intensity, high infiltration rates and low soil erodibility due to favorable soil mineralogy and a stable soil structure as reported by [70]. However, if the soil losses are not curbed, they can lead to economic losses in terms of reconstructing terraces, draining streams/channels and replenishing the soil fertility especially if bush beans are continuously cultivated.

The nutrient losses are generally very low for all bean types, in agreement with those reported by [70,79,80,81]. The smaller vegetation cover under bush beans compared with climbing beans, in agreement with the study of [30] caused higher nutrient losses from bush beans.

This also confirms the observations [82] in Tihany Peninsula who reported that higher vegetation cover caused low nutrient losses. The [83,84] studies indicated that increase in vegetation cover reduced nutrient losses. Therefore, N, P, K losses from runoff and sediment are very detrimental to agricultural production in terms of soil fertility loss and water resources degradation down the catchment. Hence management options have to be devised to reduce on the runoff and soil losses in the study area.

5. CONCLUSION

The soils in the study area had a moderate to rapid Ksat which allows an easy water movement within the soil. There were generally low soil runoff and nutrient losses from both climbing and bush beans compared with those reported in similar highlands. However, bush beans experienced significantly more runoff and soil loss than climbing beans. Therefore, climbing beans if grown in the highlands can serve a dual purpose: of providing food and controlling soil erosion. This will help to reduce the rate of land degradation in the high intensive agricultural highlands of southwestern Uganda. If traditional bush beans are cultivated, then other soil and water conservation practices must be considered in order to reduce the runoff and soil losses. Water infiltration and nutrient leaching studies should be carried out in order to further explain the low level of runoff and soil loss on the steep slopes in the highlands of southwestern Uganda.

ACKNOWLEDGMENTS

We acknowledge DAAD for funding the study under the program capacity building in integrated watershed management in East Africa. Special thanks to the farmers who hosted the experiments in Bufundi sub-catchment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Palm C, Blanco-Canqui H, De Clerck F, Gatere L, Grace P. Conservation agriculture and ecosystem services: An overview. *Agric. Ecosyst. Environ*; 2013. DOI:10.1016/j.agee. 2013.10.010.
2. Tiftonell P, Giller KE. When yield gaps are poverty traps: The paradigm of ecological intensification in African smallholder agriculture. *Field Crops Res.* 2013;143: 76–90. DOI:10.1016/j.fcr.2012.10.007.
3. Pender J, Place F, Ehui S. Strategies for sustainable agricultural development in the East African highlands. *Int. Food Policy Res. Inst. IFPRI.* 1999;41.
4. Wickama J, Okoba B, Sterk G. Effectiveness of sustainable land management measures in West Usambara highlands, Tanzania. *Catena.* 2014;118: 91–102. DOI:10.1016/j.catena.2014.01.013
5. Nyssen J, Poesen J, Haregeweyn N, Parsons T. Environmental change, geomorphic processes and land degradation in tropical highlands. *CATENA.* 2008;75:1–4. DOI:10.1016/j.catena.2008.04.010.
6. Angima SD, Stott D E, O'Neill MK, Ong CK, Weesies G.A. Soil erosion prediction using RUSLE for central Kenyan highland conditions. *Agric. Ecosyst. Environ.* 2003; 97:295–308. DOI:10.1016/S0167-8809(03)00011-2.
7. Nkonya E. Soil conservation practices and non-agricultural land use in the South western highlands of Uganda; 2002.
8. Nyssen J, Poesen J, Deckers J. Land degradation and soil and water conservation in tropical highlands. *Soil Tillage Res.* 2009;103:197–202. DOI:10.1016/j.still.2008.08.002.
9. Muchena FN, Onduru DD, Gachini GN, de Jager A. Turning the tides of soil degradation in Africa: capturing the reality and exploring opportunities. *Land Use Policy.* 2005;22:23–31. DOI:10.1016/j.landusepol.2003.07.001.
10. Pender J. Development pathways for hillsides and highlands: some lessons from Central America and East Africa. *Food Policy.* 2004;29:339–367. DOI:10.1016/j.foodpol.2004.07.005.
11. Den Biggelaar C, Lal R, Wiebe K, Eswaran H, Breneman V, Reich P. The Global Impact of Soil Erosion on Productivity*: II: Effects on Crop Yields and Production over time, in: *Advances in Agronomy.* Academic Press. 2003;49–95.
12. Carswell G. Continuities in Environmental Narratives: The Case of Kabale, Uganda, 1930-2000. *Environ. Hist.* 2003;9:3–29. DOI:10.3nv197/096734003129342755.

13. (FAO) Food and agricultural organisation. Soil fertility management in support of food security in Sub-Saharan Africa. Food & Agriculture Org; 2001.
14. Ovuka M. More people, more erosion? Land use, soil erosion and soil productivity in Murang'a district, Kenya. Land Degrad. Dev. 2000;11:111-124. DOI:10.1002/(SICI)1099-145X(200003/04)11:2<111::AID-LDR371>3.0.CO;2-I.
15. Bamutaze Y, Tenywa MM, Majaliwa MJG, Vanacker V, Bagoora F, Magunda M, Obando J, Wasige JE. Infiltration characteristics of volcanic sloping soils on Mt. Elgon, Eastern Uganda. CATENA. 2010;80:122-130. DOI:10.1016/j.catena.2009.09.006.
16. Tenywa MM, Bekunda M. Managing soils in Sub-Saharan Africa: Challenges and opportunities for soil and water conservation. J. Soil Water Conserv. 2009; 64:44A–48A. DOI:10.2489/jswc.64.1.44A.
17. Majaliwa MJG, Magunda MK, Tenywa MM, Isabirye M. Effect of topography on soil erodibility in Kakuuto micro-catchment, Uganda. East Afr. Agric. For. J. 2005;69: 19–27. DOI:10.4314/eaafj.v69i1.1802.
18. Nkonya E. Strategies for Sustainable Land Management and Poverty Reduction in Uganda. Intl Food Policy Res Inst; 2004.
19. (IFPRI) International Food Policy Research Institute. Strategies for sustainable land management and poverty reduction in Uganda. International Food Policy Research Institute; 2004.
20. Moshood K. Influence of watershed activities on the water quality and fish assemblages of a tropical African reservoir. Int. J. Trop. Biol. 2009;57:707-719.
21. Martínez-Casasnovas JA, Ramos MC. The cost of soil erosion in vineyard fields in the Penedès–Anoia Region (NE Spain). Catena, Soil Erosion Research in Europe. 2006;68:194–199. DOI:10.1016/j.catena.2006.04.007.
22. Siriri D, Tenywa MM, Raussen T, Zake JK. Crop and soil variability on terraces in the highlands of SW Uganda. Land Degrad. Dev. 2005;16:569–579. DOI:10.1002/ldr.688.
23. Sporton D. Cultivating success in Uganda: Kigezi farmers and colonial policies, Carswell G. The British institute in eastern Africa, Nairobi; James Currey, Oxford; Fountain Publishers, Kampala; Ohio University Press, Athens, 2007. ISBN 978–1–84701–601–0 (paperback), xii +258pp. Land Degrad. Dev. 2009;20:587-588. DOI:10.1002/ldr.904.
24. Adekunle A, Fatunbi A, Nyamwaro S, Buruchara R. Integrated Agricultural Research for Development: From Concept to Practice. Forum for Agricultural Research in Africa (FARA); 2013.
25. (FARA) Forum for Agricultural Research in Africa. Status of social, financial, human and physical capital in Lake Kivu Pilot Learning Site, Sub-Saharan Challenge Programme Accra Ghana; 2009. Accessed on September 15, 2011. Available:http://www.fara-africa.org/media/uploads/library/docs/ssacp/SSACP_AR2008_To_Eric_14Sep2010.pdf
26. (CIAT) International Center for Tropical Agriculture. Annual report, Eco-efficient Agriculture for the Poor; 2008.
27. Buruchara R, Chirwa R, Sperling L, Mukankusi C, Rubyogo JC, Mutonhi R, Abang MM. Development and delivery of bean varieties in Africa: The Pan-Africa bean research alliance (PABRA) model. Afr. Crop Sci. J. 2011;19:227–245. DOI:10.4314/acsj.v19i4.
28. Kimani JM, Kimani PM, Githiri SM, Kimenju JW. Genetic variation for tolerance to low soil nitrogen (N) in common beans (*Phaseolus vulgaris* L.); 2003.
29. Graham PH, Ranalli P. Common bean (*Phaseolus vulgaris* L.). Field Crops Res. 1997;53:131–146. DOI:10.1016/S0378-4290(97)00112-3.
30. Sperling L, Muyaneza S. Intensifying production among smallholder farmers: the impact of improved climbing beans in Rwanda. Afr. Crop Sci. J. 1995;3:117–125.
31. Isabirye M, Ruyschaert G, Van linden L, Poesen J, Magunda MK, Deckers J. Soil losses due to cassava and sweet potato harvesting: A case study from low input traditional agriculture. Soil and Tillage Research. 2006;92:96–103.
32. Lufafa A, Tenywa MM, Isabirye M, Majaliwa MJG, Woomer PL. Prediction of soil erosion in a Lake Victoria basin catchment using a GIS-based Universal Soil Loss model. Agricultural Systems. 2003;73:883–894.
33. Majaliwa JGM, Magunda MK, Tenywa MM, Semalulu O. Soil erosion and pollution loading from agricultural land in Bukoora

- sub-catchment, Uganda Journal of Agricultural Sciences. 2003;305-312.
34. Buruchara R. Background information on Common Beans (*Phaseolus vulgaris* L.) in Biotechnology, Breeding & Seed Systems for African Crops; 2007. Available:<http://www.africancrops.net/rockefeller/crops/beans/index.htm>
 35. Nchimbi-Msolla S, Misangu R. Seasonal distribution of common bean (*Phaseolus vulgaris* L.) Bruchid species in selected areas in Tanzania. Bean/Cowpea Collaborative Research Support Program – East Africa Proceedings: Bean Seed Workshop Arusha, Tanzania; 2000.
 36. Teverson D. Participatory promotion of disease resistant and farmer acceptable phaseolus beans in the Southern Highlands of Tanzania. R 7569 (ZA 0374). Final technical report, 1 January 2000 – 31 May 2003.
 37. Taylor D. Environmental change in montane southwest Uganda: A pollen record for the Holocene from Ahakagyazi Swamp. The Holocene. 1993;3:324–332.
 38. (LKPLS) Lake Kivu Pilot Learning Site. Baseline Survey Report. Status of Natural Resources and their Management in the Lake Kivu Pilot Learning Site; 2009.
 39. (FAO) Food and Agricultural organisation. World reference base for soil resources 2014: International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports, 106. 2014.
 40. Lafond JA, Bergeron Piette É, Caron J, Thérault Rancourt G. Evaluating fluxes in Histosols for water management in lettuce: A comparison of mass balance, evapotranspiration and lysimeter methods. Agric. Water Manag. 2014;135:73–83. DOI:10.1016/j.agwat.2013.12.016.
 41. Basic F, Kisic I, Butorac A, Nestroy O, Mesic M. Runoff and soil loss under different tillage methods on Stagnic Luvisols in central Croatia. Soil Tillage Res. 2001;62:145–151. DOI:10.1016/S0167-1987(01)00214-8.
 42. Brunner A, Park S, Ruecker G, Dikau R, Vlek PL. Catenary soil development influencing erosion susceptibility along a hillslope in Uganda. CATENA. 2004;58:1–22.
 43. Wortmann CS, Eledu CA. An agroecological zonation for Uganda: Methodology and Spatial Information. CIAT; 1999.
 44. Majaliwa JG, Tenywa MM, Rao KPC, Musana B, Fungo B, Buruchara R. Soil fertility in relation to landscape position and land use / cover types: A Case Study of the Lake Kivu Pilot Learning site; 2009.
 45. Ngigi SN, Savenije HHG, Thome JN, Rockström J, de Vries FWTP. Agro-hydrological evaluation of on-farm rainwater storage systems for supplemental irrigation in Laikipia district, Kenya. Agric. Water Manag. 2005;73: 21–41. DOI:10.1016/j.agwat.2004.09.021.
 46. Kizza CL, Majaliwa JGM, Nakileza B, Eilu G, Bahat I, Kansiiime F, Wilson J. Soil and nutrient losses along the chronosequential forest recovery gradient in Mabira Forest Reserve, Uganda. Afr. J. Agric. Res. 2013; 8:77–85.
 47. Okalebo JR, Gathua KW, Woomer PL. Laboratory methods of soil and plant analysis: A working manual. Tropical Soil Biology and Fertility Programme; 1993.
 48. Shukla MK. Soil physics: An Introduction. CRC Press; 2013.
 49. Niringiye CS, Kyamanywa S, Ssekabembe CS. Effect Of Plant Population On Yield Of Maize And Climbing Beans Grown In An Intercropping System. Afr. Crop Sci. J. 2005;13:83–93. DOI:10.4314/acsj.v13i1.27904.
 50. Bamutaze, Y. Patterns of water erosion and sediment loading in Manafwa in catchment on Mt. Elgon, Eastern Uganda; PHD dissertation, Department of Geography, Geo-information and Climatic Science, Makerere University (Uganda). 2011;255.
 51. Nunes AN, de Almeida AC, Coelho COA. Impacts of land use and cover type on runoff and soil erosion in a marginal area of Portugal. Appl. Geogr. 2011;31:687–699. DOI:10.1016/j.apgeog.2010.12.006.
 52. Gimeno-García, Vicente A, Rubio J. Influence of vegetation recovery on water erosion at short and medium-term after experimental fires in a Mediterranean shrubland Eugenia. Catena. 2007;69:150–160.
 53. Xinxiao Y, Xuexia X, Jianla J, Manliang Z, Yuanyuan X. Effects of vegetation cover and precipitation on the process of sediment produced by erosion in a small watershed of loess region. ACTA ECOLOGICA SINICA. 2006;26(1).
 54. Anasiru R, Rayes M, Budi Setiawan S. Economic valuation of soil erosion on cultivated drylands in Langge Sub-

- watershed, Gorontalo, Indonesia. *J. Nat. Sci. Res.* 2013;3.
55. Gajić B. Physical properties and organic matter of Fluvisols under forest, grassland, and 100 years of conventional tillage. *Geoderma.* 2013;200–201:114–119. DOI:10.1016/j.geoderma.2013.01.018.
 56. Zhou ZC, Gan ZT, Shanguan ZP, Dong ZB. Effects of grazing on soil physical properties and soil erodibility in semiarid grassland of the Northern Loess Plateau (China). *CATENA.* 2010;82:87-91. DOI:10.1016/j.catena.2010.05.005.
 57. Keller T, Håkansson I. Estimation of reference bulk density from soil particle size distribution and soil organic matter content. *Geoderma.* 2010;154:398–406. DOI:10.1016/j.geoderma.2009.11.013.
 58. Celik I, Gunal H, Budak M, Akpınar C. Effects of long-term organic and mineral fertilizers on bulk density and penetration resistance in semi-arid Mediterranean soil conditions. *Geoderma.* 2010;160:236–243. DOI:10.1016/j.geoderma.2010.09.028.
 59. Arvidsson J. Influence of soil texture and organic matter content on bulk density, air content, compression index and crop yield in field and laboratory compression experiments. *Soil Tillage Res.* 1998;49: 159–170. DOI:10.1016/S0167-1987(98) 00164-0.
 60. Grosbellet C, Vidal-Beaudet L, Caubel V, Charpentier S. Improvement of soil structure formation by degradation of coarse organic matter. *Geoderma.* 2011; 162:27-38. DOI:10.1016/j.geoderma.2011.01.003.
 61. Zhang W, An S, Xu Z, Cui J, Xu Q. The impact of vegetation and soil on runoff regulation in headwater streams on the east Qinghai–Tibet Plateau, China. *Catena.* 2011;87:182–189. DOI:10.1016/j.catena.2011.05.020.
 62. Gysels G, Poesen J, Bochet E, Li Y. Impact of plant roots on the resistance of soils to erosion by water: a review. *Prog. Phys. Geogr.* 2005;29:189–217. DOI:10.1191/0309133305pp443ra.
 63. Gicharu GKA comparative study of nitrogen fixation in selected cultivars of bush and climbing beans inoculated with different rhizobia strains. Kenyatta University Master's thesis; 2011.
 64. Morgan RPC. In Rody N, and Soonil R. 2010. *Soil Erosion and Conservation*, 3rd ed. Blackwell Publishing, Oxford; 2005.
 65. Xu QX, Wang TW, Cai CF, Li ZX, Shi ZH, Fang RJ. Responses of Runoff and Soil Erosion to Vegetation Removal and Tillage on Steep Lands. *Pedosphere.* 2013;23: 532–541. DOI:10.1016/S1002-0160(13)60046-6.
 66. Zhou P, Luukkanen O, Tokola T, Nieminen J. Effect of vegetation cover on soil erosion in a mountainous watershed. *Catena.* 2008;75:319–325. DOI:10.1016/j.catena.2008.07.010.
 67. Bamutaze Y. Patterns of water erosion and sediment loading in Manafwa in catchment on Mt. Elgon, Eastern Uganda; PHD dissertation, Department of Geography, Geo-information and Climatic Science, Makerere University (Uganda). 2010;255.
 68. Sasal MC, Castiglioni MG, Wilson MG. Effect of crop sequences on soil properties and runoff on natural-rainfall erosion plots under no tillage. *Soil Tillage Res.* 2010; 108:24–29. DOI:10.1016/j.still.2010.03.010.
 69. Taylor R, Howard KWF. The influence of tectonic setting on the hydrological characteristics of deeply weathered terrains: Evidence from Uganda. *J. Hydrol.* 1999;218:44–71.
 70. Tukahirwa JMB. Status and trends in soil and water conservation in Uganda. In Tenywa, M. M and Bekunda, M. A (Eds.), *Proceedings of the workshop on the past, present and prospects of soil and water conservation in Uganda.* Kampala: Soil and Water Conservation Society; 1996.
 71. Bagoora FDK. Soil loss in Rukiga highlands in Eastern Kabale, Western Uganda. PhD, Makerere University, Kampala, Makerere University Institute of Environmental Science; 1998.
 72. Molina A, Govers G, Vanacker V, Poesen J, Zeelmaekers E, Cisneros F. Runoff generation in a degraded Andean ecosystem: Interaction of vegetation cover and land use. *Catena.* 2007;71:357–370. DOI: 10.1016/j.catena.2007.04.002.
 73. De Vente J, Poesen J. Predicting soil erosion and sediment yield at the basin scale: Scale issues and semi-quantitative models. *Earth-Sci. Rev.* 2005;71:95–125. DOI:10.1016/j.earscirev.2005.02.002.
 74. Kimaro DN, Poesen J, Msanya BM, Deckers JA. Magnitude of soil erosion on the northern slope of the Uluguru Mountains, Tanzania: Interrill and rill erosion. *Catena.* 2008;75:38–44. DOI:10.1016/j.catena.2008.04.007.

75. Jamu DM, Chimphamba JB, Brummett RE. Land use and cover changes in the Likangala catchment of the Lake Chilwa basin, Malawi: implications for managing a tropical wetland. *Afr. J. Aquat. Sci.* 2004; 28:123–135.
76. Bolwig S. Land use change and soil degradation in the southwestern highlands of Uganda. A Contribution to the Strategic Criteria for Rural Investments in Productivity (SCRIP) Program of the USAID Uganda Mission. *Int. Food Policy Res. Inst.* 2033 K Str. NW Wash. DC 20006; 2002.
77. Kosmas C, Danalatos N, Cammeraat L, Chabart J, Diamantopoulos J, Farand R, Gutierrez L, Jacob A, Marques H, Martínez-Fernandez J, Mizara N, Moustakas N, Nicolau J, Oliveros C, Pinna G, Puddu R, Puigdefabregas J, Roxo M, Simao A, Stamou G, Tomasi N, Usai D, Vacca A. (The effect of land use of runoff and soil erosion rates under Mediterranean conditions. *Catena.* 1997;29:45–59.
78. Carswell G. *Cultivating success in Uganda: Kigezi farmers & colonial policies.* James Currey Publishers; 2007.
79. Zöbisch MA, Richter C, Heiligtag B, Schlott R. Nutrient losses from cropland in the Central Highlands of Kenya due to surface runoff and soil erosion. *Soil Tillage Res.* 1995;33:109–116. DOI:10.1016/0167-1987(94)00441-G.
80. Silgram M, Jackson D, Bailey A, Quinton J, Stevens C. Hillslope scale surface runoff, sediment and nutrient losses associated with tramline wheelings. *Earth Surf. Process. Landf.* 2010;35:699–706. DOI:10.1002/esp.1894.
81. Khan F, Bhatti AU, Khattak RA. Soil and nutrient losses through sediment and surface runoff under maize mono-cropping and maize-legume, intercropping from up-land sloping field. *Pak. J. Soil Sci. Pak;* 2001.
82. Centeri CS. The roles of vegetation cover in the control of soil erosion on the Tihany peninsula. *ActaBotanicaHungarica.* 2002; 44(3–4),285–295.
83. Durán ZVH, Rodríguez PCR, Peinado M FJ, de Graaf J, Martínez JRF, Flanagan DC. Environmental impact of introducing plant covers in the taluses of terraces: Implications for mitigating agricultural soil erosion and runoff. *Catena.* 2011;84:79–88.
84. Kothyari BP, Verma PK, Joshi BK, Kothyari UC. Rainfall–runoff–soil and nutrient loss relationships for plot size areas of Bhetagad watershed in Central Himalaya, India. *Journal of Hydrology.* 2004;293: 137-150.

© 2015 Gabiri et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/9937>