



Tensor Fascia Lata (TFL) Muscle Flap Associated With a Skin Graft, as a Salvage Solution in Deep Abdomino-Perineal and External Genitalia Burns

**A. H. Hissein^{a++*}, J. L. Mokako^{a++}, J. Benyoussef^{a++},
A. Meftah^{a++}, S. Sabur^{a++}, A. Chakir^{a++}, A. El Harti^{a++}
and M. Diouri^{a#}**

^a *Plastic and Reconstructive Surgery, CHU Ibn Rochd of Casablanca, Hassan II University, Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103105>

Case Report

Received: 10/05/2023
Accepted: 14/07/2023
Published: 22/07/2023

ABSTRACT

A burn corresponds to a destruction of the skin and/or deep subcutaneous structures caused by a thermal, electrical or chemical agent. When it is deep, it poses the problem of reaching the underlying noble structures. Burning by hot contact in the context of an epileptic seizure, starting from the mutilations they cause, often requires the use of shreds to cover the losses of the substances generated. It causes significant morbidity and mortality, hence the importance of urgent and adequate initial management. The abdomino-perineal region and the external genitalia (EG)

⁺⁺ *National Center for Burns;*

[#] *Head of the National Center for Burns;*

^{*}*Corresponding author: E-mail: hisseine1@gmail.com*

are rarely affected. We report a case of deep burn of the abdomino- perineal region and EG occurring in the context of a work accident in which the exposure of underlying noble elements required the use of Tensor Fascia Lata (TFL) muscle Flap. It was a 31-year-old man, chronic smoker at 10 PA and epileptic for 8 years, having stopped his treatment, who was admitted to hour 2 (H2) for his burn by hot contact which occurred following an epileptic attack. The patient had fallen on a hot welding instrument in the context of an informal work accident. Initially transferred to a local hospital, he was sent to us for care. Examination found a 3rd degree burn affecting the abdomino - inguino -pubic regions, the sheath of the penis and the scrotum. The initial management consisted of hemodynamic stabilization, the realization of a relief incision at the level of the sheath of the penis and local care. An early excision of abdomino-perineal on day 3 had been performed having exposed the spermatic cords, followed by coverage on day 10 by a Tensor Fascia Lata muscle Flap of the inguinal region - pubis, split – thickness skin graft of the sheath of the penis. The postoperative course was simple.

Keywords: *Deep burn; hot contact burn; abdominal-perineal burn; Tensor Fascia Lata muscle flap; EG skin graft.*

1. INTRODUCTION

The deep burn poses the problem of reaching the underlying noble structures. Burning by hot contact in the context of an epileptic seizure, starting from the mutilations they cause, often requires the use of shreds to cover the losses of the substances generated. It causes significant morbidity and mortality, hence the importance of urgent and adequate initial management. The abdomino-perineal and EG regions are rarely affected. Being at the urogenital and digestive crossroads, these regions are characterized by the presence of digestive and urogenital orifices as well as the risk of secondary infection that they entail.

Genital and perineal burns have serious physical, functional, sexual and psychological consequences that have a direct influence on quality of life [1]. The American Burn Association classifies these burns as major burns. They are very worrying, not only for the victim, but also for the treating surgeon [2, 3, 4].

Reconstruction can be very demanding because it is necessary to preserve the sexual and

excretory functions of the perineal area, to prevent the occurrence of postoperative eventration [5].

2. CLINICAL CASE

He was a 31-year-old man, chronic smoker at 10 PA and epileptic for 8 years, having stopped his treatment and admitted to hour 2 (H2) for his burn by hot contact which occurred following an epileptic attack.

The patient had fallen on a hot welding instrument (Fig. 1) as part of an informal work accident. Initially transferred to a local hospital, he was sent to us for treatment.

The examination found a 3rd degree burn involving the inguino -pubic regions, the sheath of the penis and the scrotum (Fig. 2).

The initial treatment consisted of hemodynamic stability, performing a relief incision at the level of the sheath of the penis (Fig. 3) and local treatment using silver sulfadiazine- type flammazine cream associated with sheets of tulle gras.



Fig. 1. Automatic electric heating system for the welding machine



Fig. 2. Third-degree burns on the inguinal-pubic regions, the shaft of the penis, and the scrotum



Fig. 3. Showing the discharge incisions on the shaft of the penis



Fig 4. Early excision by avulsion of the abdomino- perineal areas, the shaft of the penis, exposing the spermatic cord

We proceeded to an early excision (day 3 of the burn) of the abdomino-perineal areas and of the sheath of the penis exposing the spermatic cord (Fig. 4). The sheath of the penis was grafted on day 2 of the excision (Fig. 5) with split – thickness skin graft. The exposure of the

spermatic cords was covered on day 10 (D10) with a flap of the tensor fascia lata allowing easy bilateral coverage of the inguino -pubic region and skin grafting of the adjacent areas (Fig. 6). The postoperative course was simple.

We noted no necrosis of the flap, nor lysis of the skin grafts. The delbet blade was removed at the first dressing change on day 3 (D3). The skin graft staples were removed on day 5, and the donor area staples were removed on day 10.

The patient had received very early rehabilitation on postoperative day 10. Discharge was permitted on day 15. Follow-up was carried out in outpatient consultations on day 30, 3 months, and 6 months. Skin coverage was satisfactory, with good skin flexibility and trophicity. There was a resumption of sexual function and good socio-professional reintegration. The patient reported being satisfied with the results (Fig. 7).

3. DISCUSSION

Deep abdomino -perineal burns associated with external genitalia are rare [1, 6]. They are often encountered in extensive burns. The external genitalia have important anatomical, functional and physiological functions for the human body.

Their burns lead to serious physical, functional, sexual and psychological consequences that have a direct influence on the quality of life [1, 4, 6, 7].

Abdomino-perineal and EG burns are immediately considered a serious factor with a risk of secondary infection and sepsis. They require transfer and care in a specialized center [8].

The circumstances of the occurrence of these burns, such as epileptic seizures, are in themselves a pejorative element due to prolonged contact with the injuring agent [9]. Initial management in a specialized center should guarantee not only vital functions but also adequate local treatment. From admission and in parallel with the continuation of resuscitation measures, a thorough initial clinical examination to search for associated trauma is required in all cases.



Fig. 5. Split-thickness skin graft on the shaft of the penis. Note the exposure of the spermatic cords



Fig. 6. The coverage of the spermatic cords with a TFL flap bilaterally and Split-thickness skin grafting of adjacent areas

Maintenance of fluid and electrolyte balance is important and should be adapted to the patient's metabolic and renal status. Cutaneous relief incisions, particularly in our patient at the sheath of the penis, must be performed urgently within 6 hours before signs of tissue damage appear [10]. In our patient, they were performed at hour 2 (h2).

Early excision-grafting is necessary in order to reduce the risk of infection and improve local trophic status [11]. Our patient benefited an early excision by avulsion of the abdomino- perineal

areas and of the sheath of the penis exposing the spermatic cord (Fig. 4). The sheath of the penis was grafted on day 2 of the excision (Fig. 5).

When there is exposure of a noble element, namely nerve, tendon, vessels, joint or bone, the provision of local, regional or distant pedicled or free flaps is necessary [12]. In our patient, the exposure of the spermatic cords was covered on D10 by a flap of the TFL allowing easy coverage of the inguino -pubic region bilaterally and skin grafting of the adjacent areas (Fig. 6).



Fig. 7. 6 months postoperative front, strict left and right profiles, and 3/4 left and right profiles

The spermatic cord is made up of arteries, veins, lymphatics, nerves and the vas deferens. The testicular artery supplies blood to the testicles, while the vas deferens is the excretory duct of the testicles. Damage or injury to the cord at any point in its course can affect testicular blood flow and/or interrupt the continuity of the vas deferens [13, 14, 15]. It is clear here that coverage by a flap is necessary. Furthermore, the fragility of this inguino-crural region requires a strong reconstruction. The choice of the flap of the tensor fascia lata seemed judicious to us because of the sufficient tissue it provides. Other flaps, such as the difficult-to-dissect anterolateral thigh flap, could be used, but its rotational axis does not allow for bilateral coverage of the inguino-pubic region. Artificial dermis, due to its

cost and risk of infection, did not appeal to our choice, and muscle flaps are not indicated in this area [16, 17, 18].

The fascia lata flap allowed for effective and bilateral coverage of the exposed spermatic cords. The postoperative course was uneventful.

It is important to initiate early functional rehabilitation and physiotherapy in order to prevent the development of sequelae. Our patient underwent early rehabilitation starting on the 10th postoperative day. Discharge was permitted on the 15th day. Follow-up was conducted through outpatient consultations at 30 days, 3 months, and 6 months. The skin coverage was satisfactory, with good skin suppleness and

trophicity. Copulation function was restored, and there was a good socio-professional reintegration. The patient reported being satisfied with the results. Psychological follow-up was also provided.

4. CONCLUSION

Deep abdomino-perineal burns and burns of the external genitalia are rare but serious, especially since they occur during an epileptic seizure. Early excision-graft and coverage by a flap of the noble structures are necessary. The use of a tensor fascia lata flap in the case of exposed spermatic cords is an interesting option. It allows effective coverage and simple technique.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Carrillo-Cordova LD, Carrillo-Esper R, Carrillo-Cordova JR. Burns to the genitalia and perineum: General considerations and urologic management. *Rev Mex Urol*. 2017;77(1):36–49.
2. Tresh A, Baradaran N, Gaither TW, Fergus KB, Liaw A, Balakrishnan A, Hampson LA, Breyer BN. Genital burns in the United States: Disproportionate prevalence in the pediatric population. *Burns*. 2018;44(5):1366-1371.
3. Abel NJ, Klaassen Z, Mansour EH, Marano MA, Petrone SJ, Hounq AP, et al. Clinical outcome analysis of male and female genital burn injuries: a 15-year experience at a Level-1 Burn Center. *Int J Urol*. 2012;19:351–8.
4. Umar Farooq Baba, Ahmad Shah R, Rasool A, Kaul A, Hafeez Wani A. Perineal burns epidemiology, severity factors, management issues and outcome. *Burns Open*; 2022. DOI: 10.1016/j.burnso.2022.03.003
5. Sharma Rk, Parashar A. The management of perineal wounds. *Indian J Plast Surg*. 2012;45:352-63.
6. Sajad W, Hamid R. Outcome of split-thickness skin grafting and multiple z-plasties in postburn contractures of groin and perineum: a 15-year experience. *Plast Surg Int*. 2014;2014:358526. 10.1155/2014/358526. Epub 2014 May 21.7- PMID: 24967100; PMCID: PMC4055381.
7. Thakur JS, Chauhan CGS, Diwana VK, Chuahan DC, Thakur A. Perineal burn contractures: an experience in a tertiary hospital of a Himalayan state. *Indian J Plastic Surgery*. 2008;41(2):190.
8. Oksüz M, Deliaga H, Topkara A, Koçak OF. An unexpected long-term complication of genital burn in a child: Secondary cryptorchidism. *Ulus Travma Acil Cerrahi Derg*. 2018;24:85–7.
9. Gagnani A, Müller BR, Oliveira AF, Ferreira LM. Burns and epilepsy – review and case report. *Burns*. 2015;41(2):e15-e18. DOI:10.1016/j.burns.2014.08.004
10. Bakkali H, Ababou K, Bellamari H, Ennouhi A, Nassim Sabah T, Achbouk A, Moussaoui A, Fouadi FZ, Siah S, Ikhrai H. La prise en charge des brûlures électriques: a propos de 30 cas. *Ann Burns Fire Disasters*. 2009;22(1):33-6.
11. Janzekovic Z. A new concept in the early excision and immediate grafting of burns. *J Trauma*. 1970;10(12):1103–8.
12. Zhu ZX, Xu XG, Li WP. Experience of 14 years of emergency reconstruction of electrical injuries. *Burns*. 2003;29:65-72.
13. Firoozabadi R, Stafford P, Routt M. Risk of spermatic cord injury during anterior pelvic ring and acetabular surgery: an anatomical study. *Arch Bone Jt Surg*. 2015;3(4):269-73.
14. Gray H. *Gray's Anatomy*. Philadelphia: Lea & Febiger; 1985.
15. Raman J, Goldstein M. Intraoperative characterization of arterial vasculature in spermatic cord. *Urology*. 2004;64(3):561-4.
16. Song Z, et al. Abdominal wall reconstruction following resection of large abdominal aggressive neoplasms using tensor fascia lata flap with or without mesh

- reinforcement. *Hernia : The Journal of Hernias and Abdominal Wall Surgery*. 2018;22(2):333-341.
17. Iwai, Miho, et al. Successful resection of growing teratoma syndrome in the abdominal wall with reconstruction using tensor fascia lata muscle flap: A case report and literature review. *Gynecologic Oncology Reports*. 2023;101161.
18. Williams JK, Carlson GW, deChalain T, Howell R, Coleman JJ. Role of tensor fasciae latae in abdominal wall reconstruction. *Plast Reconstr Surg*. 1998;101(3):713-8.

© 2023 Hissein et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103105>