



# **Influence of Sulphur and Zinc on Growth, Yield Attributes and Economics of Cowpea (*Vigna unguiculata* L.)**

**Anju Kundu <sup>a+++</sup>, Biswarup Mehera <sup>b#</sup> and Prateek Kumar <sup>at</sup>**

<sup>a</sup> Department of Agronomy, Naini Agricultural institute, SHUATS, Prayagraj, Uttar Pradesh, India.

<sup>b</sup> Naini Agricultural institute, SHUATS, Prayagraj, Uttar Pradesh, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2023/v13i102970

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105637>

**Original Research Article**

**Received: 01/07/2023**

**Accepted: 05/09/2023**

**Published: 12/09/2023**

## **ABSTRACT**

The experiment was laid out in Randomized Block Design with nine treatments each replicated thrice on the basis of one year of experimentation. The treatments which are T1: 2% Sulphur+0.5% zinc, T2: 2% Sulphur+ 1% Zinc, T3: 2% Sulphur + 1.5% Zinc, T4: 3% Sulphur + 0.5% Zinc, T5: 3% Sulphur +1% Zinc, T6: 3% Sulphur +1.5% Zinc, T7: 4% Sulphur + 0.5% Zinc, T8: 4% Sulphur+ 1% Zinc, T9: 4% Sulphur +1.5% Zinc, T10 N:P: K 20:40:40 kg/ha. The results showed that application of 4% Sulphur+ 1.5% Zinc was recorded significantly higher plant height (79.17 cm), nodules/ plant (30.93), Plant dry weight (22.83 g/plant), No. of pods/ plant (12.14), Test weight (100.23 g), maximum crop growth rate (8.82g/m<sup>2</sup>/day). However, higher Seed yield (1.32 t/ha), Stover Yield (3.48 t/ha) gross returns (INR84971.50/ha), net return (INR60594.40/ha) and benefit cost ratio (1.28) were obtained with application of 4% Sulphur+1.5% Zinc as compared to other treatments.

<sup>++</sup>M.Sc. Scholar;

<sup>#</sup>Dean;

<sup>†</sup>Ph.D Scholar;

<sup>\*</sup>Corresponding author: E-mail: [anjukundu181@gmail.com](mailto:anjukundu181@gmail.com);

**Keywords:** Sulphur; zinc; growth; yield; cowpea; economics.

## 1. INTRODUCTION

Cowpea (*Vigna unguiculata* L.) commonly referred to as "Lobia," serves as a versatile plant, functioning as a pulse, fodder, and green manure crop. With its origins tracing back to the Asian and African tropics, cowpea boasts a lengthy history as one of the earliest pulse crops in these regions. Renowned for its protein abundance and diverse nutrient content, it has earned the moniker of "vegetable meat." This plant holds particular significance as a valuable protein source for economically disadvantaged populations, standing as a timeless staple in human diets and likely having been cultivated since the Neolithic era. At its core, cowpea seeds stand out as a primary reservoir of protein and carbohydrates. These seeds assume a critical role in human nutrition and also serve as a nourishing feed for livestock. Notably, the protein within cowpea seeds showcases an impressive concentration of essential amino acids such as lysine and tryptophan, setting it apart from cereal grains. The mature cowpea seed contains 24.8% protein, 63.6% carbohydrate, 1.9% fat, 6.3% fibre, 7.4 ppm thiamine, 4.2 ppm riboflavin and 28.1 ppm niacin [1]. "Cowpea boasts a remarkable versatility, finding utility as a catch crop, mulch crop, intercrop, mixed crop, and green crop. Within the realm of nutritional security, pulses hold a crucial position, and cowpea is no exception. Its cultivation footprint spans across Africa, Asia, North and South America, Australia, as well as the central and southern regions of Europe. Operating as a leguminous plant, cowpea plays a pivotal role by harnessing significant levels of atmospheric nitrogen to fulfill its own needs. In India it is cultivated mainly in UP, MP, Bihar, Punjab, Haryana, Rajasthan, HP, etc. where it is grown for both vegetable and pulse purposes and is a highly remunerative crop" [2].

It is asserted that in India, pulses are cultivated on an area of 29.36 mha with a production and productivity of 24.51 mt and 835 kg per hectare [3]. "Wide spread deficiency of micronutrient has become major constraint for achieving higher genetic potential yield along with quality of crop. Among the micronutrients, the deficiency of Zinc and sulphur has become wide spread in recent years all over the country. In India next to nitrogen, phosphorus, potassium and sulphur 5th important yield limiting nutrient is Zinc which accounts for 48 per cent of Indian soils" [4].

"Plants require essential nutrients for better growth and yield. Sulphur is a major nutrient for healthy crop growth. It is the fourth major nutrient after NPK" [5]. "Sulphur is the constituent of the cysteine, amino acids, methionine and chlorophyll" [6]. "Oilseed crops require higher amounts of S for their growth and oil yield because it is critical in the oil synthesis and formation of bold grains for oil production in the oil seeds crops" [7].

"Zinc is one of the seventh plant micronutrient, involved in many enzymatic activities of the plant. It functions generally as a metal activator of enzymes. It is reported that, Zinc improves crop productivity almost as much as major nutrients. Besides increasing crop yield, it increases the crude protein content, amino acids, energy value and total lipid in chickpea, soybean, blackgram, etc. Zn deficiency can also adversely affect the quality of harvested products, plants susceptibility to injury by high light or temperature intensity and infection by fungal diseases can also increase. Zinc seems to affect the capacity for water uptake and transport in plants and also reduce the adverse effects of short periods of heat and salt stress" [8]. "The accumulation of Zinc in edible parts of plant serves as Zinc source for primary consumers. Unfortunately, about 50% of Indian soils are deficient in Zinc and sulphur expected to further increase up to 63% by 2025 which imparting Zinc malnutrition in population especially in children" (Singh, 2006); [9]. "One third of the world population is reported at the risk of Zinc malnutrition due to inadequate dietary intake of Zinc resulting from wide spread hidden hunger of Zinc in seeds and feeds" [10]. "Zinc being essential nutrient plays a significant role in stomatal regulation and reducing the tensions of less water by creating ionic balance in plants system and is involved in various physiological processes such as synthesis of protein and carbohydrates. Similarly, S application improves growth, and enhances stress tolerance in plants and improves grain production. Sulphur and Zinc are the essential plant micronutrients and their importance for crop productivity is similar to that of major nutrients. Both play an important role in the basic plant functions like photosynthesis, protein and chlorophyll synthesis" [11]. Keeping all these facts in view, the present investigation entitled, "Effect of sulphur and zinc on growth and yield of cowpea (*Vigna unguiculata* L.)".

## 2. MATERIALS AND METHODS

The field experiment was conducted during *Zaid* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P.). The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 8), low level organic carbon (0.62%), medium available N (225 Kg/ha), high in available P (38.2 kg/ha) and low available K (240.7 kg/ha). The experiment was laid out in RBD with 10 treatments each replicated thrice. The treatment combinations are T<sub>1</sub>: 2% Sulphur+0.5% zinc, T<sub>2</sub>: 2% Sulphur+ 1% Zinc, T<sub>3</sub>: 2% Sulphur + 1.5% Zinc, T<sub>4</sub>: 3% Sulphur + 0.5% Zinc, T<sub>5</sub>: 3% Sulphur +1% Zinc, T<sub>6</sub>: 3% Sulphur +1.5% Zinc, T<sub>7</sub>: 4% Sulphur + 0.5% Zinc, T<sub>8</sub>: 4% Sulphur+ 1% Zinc, T<sub>9</sub>: 4% Sulphur +1.5% Zinc T<sub>10</sub>: N:P: K 20:40:40 kg/ha.

The growth parameters and yield, production was recorded at harvest from randomly selected plants in each plot. The data was computed and analyzed by following statistical method of Gomez and Gomez (1984).

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters Plant Height (cm)

The Data revealed that that the 75 DAS, plant height (79.17 cm) was maximum T<sub>9</sub> 4% Sulphur + ZnSO<sub>4</sub> 1.5%. While T<sub>8</sub> 4% Sulphur + ZnSO<sub>4</sub> 1% (75.77 cm) and T<sub>7</sub> 4% Sulphur + ZnSO<sub>4</sub> 0.5% (74.17 cm) was found to be statistically at par with T<sub>9</sub>. "Increase in plant height might be the involvement of micronutrients in different physiological processes like enzyme activation, electron transport, chlorophyll formation, stomatal regulation, etc. Wytthe plant height gradually increased, which might be attributable to greater photosynthetic activity and chlorophyll synthesis due to Sulphur and zinc fertilization resulting into better vegetative growth" [11]. Similar results were reported by Barbara M Humtsoe et al. [12]; Lal Babu Singh et al. (2015).

#### 3.1.1 Number of leaves

Observed that the 75 DAS, plant leaf (28.67) was maximum T<sub>9</sub> 4% Sulphur + ZnSO<sub>4</sub> 1.5%. While T<sub>8</sub> 4% Sulphur + ZnSO<sub>4</sub> 1% (27.60), T<sub>7</sub> 4% Sulphur + ZnSO<sub>4</sub> 0.5% (26.40) T<sub>6</sub> Sulphur 3% + Zinc 1.5% (24.93) and T<sub>5</sub> Sulphur 3% + Zinc 1% (23.47) was found to be statistically at par with T<sub>9</sub>. "The best obtained results were

found with effect of Sulphur and zinc might be attributed to the favorable influence of them on plant metabolism and biological process activity and their stimulating effect on photosynthetic pigments and enzyme activity which in turn encourage vegetative growth. The reported positive effect of application of Zn on an enhanced branching impulses mainly attributed to promotion of bud and branch development by the auxins whereas Zn application ultimately increased the availability of other nutrients and accelerated the translocation of photo assimilates" Ravichandra et al. [13] and Hamouda et al. [14].

#### 3.1.2 Number of branches

Observed that the 75 DAS, plant branch (9.74) was maximum T<sub>9</sub> 4% Sulphur + ZnSO<sub>4</sub> 1.5%. While T<sub>8</sub> 4% Sulphur + ZnSO<sub>4</sub> 1% (9.59), T<sub>7</sub> 4% Sulphur + ZnSO<sub>4</sub> 0.5% (9.28), T<sub>6</sub> Sulphur 3% +Zinc 1.5% (9.02), T<sub>5</sub> Sulphur 3%+Zinc 1% (8.65), T<sub>4</sub> Sulphur3%+Zinc 0.5% (8.55) and T<sub>3</sub> Sulphur 2% +Zinc1.5% (8.28) was found to be statistically at par with T<sub>9</sub>. "The best obtained results were found with effect of Sulphur or zinc might be attributed to the favorable influence of them on plant metabolism and biological process activity and their stimulating effect on photosynthetic pigments and enzyme activity which in turn encourage vegetative growth. The reported positive effect of application of Zn on an enhanced branching in pulses mainly attributed to promotion of bud and branch development by the auxins whereas Zn application ultimately increased the availability of other nutrients and accelerated the translocation of photo assimilates" Ravichandra et al. [13].

#### 3.1.3 Number of nodules

Observed that the 75 DAS, plant nodule (30.93) was maximum T<sub>9</sub> 4% Sulphur + ZnSO<sub>4</sub> 1.5%. While T<sub>8</sub> 4% Sulphur + ZnSO<sub>4</sub> 1% (29.87) and T<sub>7</sub> 4% Sulphur + ZnSO<sub>4</sub> 0.5% (28.67), T<sub>6</sub> Sulphur 3% + ZnSO<sub>4</sub> 1.5% (27.20) and T<sub>5</sub> Sulphur 3% + ZnSO<sub>4</sub> 1% (25.73) was found to be statistically at par with T<sub>9</sub>. The best obtained results were found with effect of zinc or Sulphur might be attributed to the favorable influence of them on plant metabolism and biological process activity and their stimulating effect on photosynthetic pigments and enzyme activity which in turn encourage vegetative growth. The reported positive effect of application of Zn on an enhanced branching impulses mainly attributed

to promotion of bud and branch development by the auxins whereas Zn application ultimately increased the availability of other nutrients and accelerated the translocation of photo assimilates. Similar results were reported by Ravichandra et al. [13] and Hamouda et al. [14].

### 3.1.4 Plant dry weight

Observed that the 75 DAS, plant dry weight (22.83) was maximum T<sub>9</sub> 4% Sulphur + ZnSO<sub>4</sub> 1.5%. While T<sub>8</sub> 4% Sulphur + ZnSO<sub>4</sub> 1% (22.03 g), T<sub>7</sub> 4% Sulphur + ZnSO<sub>4</sub> 0.5% (21.02 g), T<sub>6</sub> Sulphur 3% + ZnSO<sub>4</sub> 1.5% (20.84 g), T<sub>5</sub> Sulphur 3% + ZnSO<sub>4</sub> 1 % (20.27 g), T<sub>4</sub> Sulphur 3%+ZnSO<sub>4</sub>0.5 % (19.45g) was found to be statistically at par with T<sub>9</sub>. Dry weight was increased significantly with increasing as spray of sulphur 4% and Zinc 1.5%. As Zinc generally influences cell division and nitrogen absorption from the soil might enhanced plant growth which reflects in terms of plant dry weight. These findings are in harmony with those obtained by Barbara M Humtsoe et al. [12] and El-Afifi et al. (2016).

## 3.2 Yield Parameters

### 3.2.1 Number of pods/plants

Maximum (12.14) number of pods/plants was obtained with application of T<sub>9</sub> (4% Sulphur + ZnSO<sub>4</sub> 1.5%). Whereas, T<sub>8</sub> (4% Sulphur + ZnSO<sub>4</sub> 1%) (11.53) were statistically at par T<sub>9</sub>. "Application of Sulphur along with Zinc increased the number of pods per plant might be due to the enhanced early vegetative growth in terms of higher leaf area, dry matter accumulation and

vigorous root system resulted in more branches which consequently increased the number of pods bearing branches significantly" Singh et al. 2020.

### 3.2.2 Seeds/pod (No.)

Maximum (11.03) number of seeds/pods was obtained with application of T<sub>9</sub> (4% Sulphur + ZnSO<sub>4</sub> 1.5%) Whereas, T<sub>8</sub> (4% Sulphur + ZnSO<sub>4</sub> 1%) (10.55) and T<sub>7</sub> (4% Sulphur + ZnSO<sub>4</sub> 0.5%) (9.88) were statistically at par T<sub>9</sub>. Application of Sulphur and Zinc to cowpea crop generally improves fruit growth by synthesizing tryptophan and auxin. The enhancement effect on seeds/pod and pods/plant attributed to the favorable influence of the Zn application to crops on nutrient metabolism, biological activity and growth parameters and hence, applied zinc resulted in taller and higher enzyme activity which in turn encourage more seeds/pod and pods/plant. Similar findings have been reported earlier by Hamouda et al. [14] and Nishant Srivastava et al. [15].

### 3.2.3 Test weight (g)

Maximum (100.23 g) test weight (g) was obtained with application of T<sub>9</sub> (4% Sulphur + ZnSO<sub>4</sub>1.5%) which was superior over rest of the treatments. Whereas, T<sub>8</sub> (4% Sulphur + ZnSO<sub>4</sub> 1%) (96.09 g) and T<sub>7</sub> (4% Sulphur + ZnSO<sub>4</sub> 0.5%) (94.50 g) were statistically at par T<sub>9</sub>. Zinc plays a very important role in the metabolism of the plant process by influencing the activity of growth enzymes as well as it is involved in carbohydrate metabolism, maintenance of the

**Table 1. Effect of sulphur and zinc on growth attributes of Cowpea**

S No	Treatments	plant height (cm)	Number of nodules/plant	Plant dry weight (g/plant)
1.	Sulphur 2% + ZnSO <sub>4</sub> 0.5%	61.91	21.40	17.36
2.	Sulphur 2% + ZnSO <sub>4</sub> 1%	68.40	24.13	18.07
3.	Sulphur 2% + ZnSO <sub>4</sub> 1.5%	69.37	24.33	18.76
4.	Sulphur 3% + ZnSO <sub>4</sub> 0.5%	70.17	24.93	19.45
5.	Sulphur 3% + ZnSO <sub>4</sub> 1%	71.17	25.73	20.27
6.	Sulphur 3% + ZnSO <sub>4</sub> 1.5%	72.60	27.20	20.84
7.	Sulphur 4% + ZnSO <sub>4</sub> 0.5%	74.17	28.67	21.02
8.	Sulphur 4% + ZnSO <sub>4</sub> 1%	75.77	29.87	22.03
9.	Sulphur 4% + ZnSO <sub>4</sub> 1.5%	79.17	30.93	22.83
10.	(Control) N: P: K 20:40:40 kg/ha	59.78	19.67	16.25
F-test		S	S	S
SEm(±)		2.16	1.89	1.30
CD at 5%		6.42	5.65	3.88

**Table 2. Effect of Sulphur and zinc on Yield of cowpea**

S. No.	Treatments	No. of pods/plant	No. of seed/pod	Test weight (g)	Seed yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
1.	Sulphur 2% + ZnSO <sub>4</sub> 0.5%	7.11	7.03	86.87	0.86	2.96	21.90
2.	Sulphur 2% + ZnSO <sub>4</sub> 1%	7.75	7.38	87.85	0.90	2.97	22.56
3.	Sulphur 2% + ZnSO <sub>4</sub> 1.5%	8.19	7.88	88.54	0.95	3.07	23.81
4.	Sulphur 3% + ZnSO <sub>4</sub> 0.5%	8.81	8.61	89.55	1.00	3.10	25.40
5.	Sulphur 3% + ZnSO <sub>4</sub> 1%	9.59	9.03	89.67	1.05	3.25	26.14
6.	Sulphur 3% + ZnSO <sub>4</sub> 1.5%	10.48	9.43	90.98	1.12	3.28	27.32
7.	Sulphur 4% + ZnSO <sub>4</sub> 0.5%	10.95	9.88	94.50	1.22	3.31	27.58
8.	Sulphur 4% + ZnSO <sub>4</sub> 1%	11.53	10.55	96.09	1.27	3.39	27.66
9.	Sulphur 4% + ZnSO <sub>4</sub> 1.5%	12.14	11.03	100.23	1.32	3.48	29.48
10.	(Control) N: P: K 20:40:40 kg/ha	6.44	6.50	87.36	0.80	2.77	24.71
F-test		S	S	S	S	S	S
SEm(±)		0.354	0.531	2.748	0.053	1.21	0.12
CD at 5%		1.05	1.57	8.16	0.15	0.36	3.62

**Table 3. Economics of sulphur and zinc of cowpea**

Sr. No.	Treatments	Cost of cultivation (INR/ha)	Gross return (INR/ha)	Net return (INR/ha)	B:C
1.	Sulphur 2% + ZnSO <sub>4</sub> 0.5%	23983.10	56069.50	32086.40	1.34
2.	Sulphur 2% + ZnSO <sub>4</sub> 1%	24133.10	58665.00	34531.90	1.43
3.	Sulphur 2% + ZnSO <sub>4</sub> 1.5%	24283.10	61711.50	37428.40	1.54
4.	Sulphur 3% + ZnSO <sub>4</sub> 0.5%	24030.10	64737.00	40706.90	1.69
5.	Sulphur 3% + ZnSO <sub>4</sub> 1%	24180.10	68393.00	44212.90	1.83
6.	Sulphur 3% + ZnSO <sub>4</sub> 1.5%	24330.10	72184.00	47853.90	1.97
7.	Sulphur 4% + ZnSO <sub>4</sub> 0.5%	24077.10	78231.00	54153.90	2.25
8.	Sulphur 4% + ZnSO <sub>4</sub> 1%	24227.10	81695.00	57467.90	2.37
9.	Sulphur 4% + ZnSO <sub>4</sub> 1.5%	24377.10	84971.50	60594.40	2.49
10.	(Control) N: P: K 20:40:40 kg/ha	23139.10	52662.60	29523.50	1.28

integrity of cellular membranes, protein synthesis, and regulation of auxin synthesis and pollen formation. The positive effect of Sulphur may be due to key role in plant metabolism and in the synthesis of nucleic acid. Similar findings were under the conformity of Ravichandra et al. [13].

### 3.2.4 Seed yield (t/ha)

Highest (1.32 t/ha) seed yield (t/ha) was obtained with application of T<sub>9</sub> (4% Sulphur + ZnSO<sub>4</sub> 1.5%) Whereas, T<sub>8</sub> (4% Sulphur + ZnSO<sub>4</sub> 1%) (1.27t/ha) and T<sub>7</sub> (4% Sulphur + ZnSO<sub>4</sub> 0.5%) (1.22t/ha) were statistically at par T<sub>9</sub>. Zinc plays a vital role in increasing seed yield because zinc takes place in many physiological processes of plant such as chlorophyll formation, stomatal regulation, starch utilization which enhance seed yield. Zinc also converts ammonia to nitrate in crops which contribute to yield. These results are in confirmatory with the work of Abid Khan et al. [16].

### 3.2.5 Stover yield

Stover yield (t/ha) was maximum (3.48 t/ha) in the application T<sub>9</sub> (4% Sulphur + ZnSO<sub>4</sub>1.5%). Whereas, T<sub>8</sub> (4% Sulphur + ZnSO<sub>4</sub>1%) (3.39 t/ha) and T<sub>7</sub> (4% Sulphur + ZnSO<sub>4</sub> 0.5%) (3.31 t/ha) T<sub>6</sub> Sulphur 3% + ZnSO<sub>4</sub> 1.5% (3.28t/ha) and T<sub>5</sub> Sulphur 3% + ZnSO<sub>4</sub> 1% (3.25t/ha) were statistically at par T<sub>9</sub>. This indicates that higher uptake of Zn resulted maximum yield under the same treatments. The present findings are well in agreement with that of Debnath et al. [17].

### 3.2.6 Harvest index

Maximum harvest index (%) was obtained with application of T<sub>9</sub> (4% Sulphur + ZnSO<sub>4</sub> 1.5%)

(29.48%). While T<sub>8</sub> 4% Sulphur + ZnSO<sub>4</sub> 1% (27.66%), T<sub>7</sub> 4% Sulphur + ZnSO<sub>4</sub> 0.5% (27.58%), T<sub>6</sub> Sulphur 3% + ZnSO<sub>4</sub> 1.5% (27.32%) harvest index (%) was found to be statistically at par with T<sub>9</sub>. This was mainly because of increase in seed yield with optimum stover yield which in turn resulted in higher harvest index. These results were in conformity with findings of Nishant Srivastava et al. [15].

### 3.3 Economics

The result showed that maximum gross returns (Rs.84971.50/ha), net return (Rs.60594.40/ha) and benefit cost ratio (1.28) was recorded in treatment 9 Sulphur 4% and ZnSO<sub>4</sub> 1.5%. Higher seed yield and stover yield positively correlates with economics. The increased gross returns, net returns and benefit: cost ratio was due to increased seed and stover yield under foliar application of Sulphur 4% with zinc at 1.5%. The findings are in close conformity with those of Durugude et al. [18-24].

### 4. CONCLUSION

It can be concluded that in Cowpea with the application of Sulphur 4% and ZnSO<sub>4</sub> 1.5% treatment 9 recorded higher Seed yield, higher gross returns, net returns and benefit cost ratio.

### ACKNOWLEDGEMENT

The authors are thankful to Department of Agronomy, Naini Agricultural Institute, Prayagraj, Sam Higginbottom University of Agriculture Technology and sciences, (U.P) India for providing necessary facilities to undertaken the studies.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Ahlawat IPS, Shivakumar BG. Kharif Pulses. In Textbook of Field Crops Production (R. Prasad, Ed.) ICAR, New Delhi, India; 2005.
2. Upadhyay RG, Anita Singh. Effect of nitrogen and zinc on nodulation, growth and yield of cowpea (*Vigna unguiculata*). Legume Research. 2016;39(1):149-151.
3. Anonymous. Annual report. Ministry of Agriculture and Farmers Welfare, Director of Pulse Development, Government of India. 2016-17a;9.
4. Vinodkumar HV, Channakeshava S, Basavaraja B, Ananathakumar. Effect of soil and foliar application of zinc on growth and yield of green gram (*Vigna radiata* L.). International Journal of Current Microbiology and Applied Sciences. 2020; 9(4):501512.
5. Najar GR, Singh SR, Akthar F, Hakeem SA. Influence of sulphur levels on yield, uptake and quality of soyabean (*Glycine max*) under temperate conditions of Kashmir valley. Ind. J. Agric. Sci. 2011;81(4):340-343.
6. Wani MA, Agha FA, Malik MA, Rather ZA. Response of sunflower to sulphur application under Kashmir conditions. Appl. Biol. Res. 2001;3:19-22.
7. Brady NC, Weil RR. The nature and properties of soils. Singapore: Pearson Educ; 2002.
8. Hitsuda K, Yamada M, Dirceu K. Soil and crop management, sulphur requirement of eight crops at early stages of growth. Agron. J. 2005;97:155159.
9. Chalak AL, Vaikar SL, Barangule Sanjivani. Effect of varying levels of potassium and zinc on yield, yield attributes, quality of pigeon pea (*Cajanus cajan* L. Millsp.). International Journal of Chemical Studies. 2018;6(5):1432-1435.
10. Shukla AK, Tiwari PK. Progress report of AICRP on micro and secondary nutrients and pollutant elements in soils and plants, IISS, Bhopal. Journal of Pharmacognosy and Phytochemistry. 2014;7(6):1-4.
11. Singh MV. Detrimental effect of zinc deficiency on crops productivity and human health. First Global Conference on Biofortification, Harvest Plus, Washington, USA; 2010.
12. Karimunnisa S, Umesha C, Khan SA. Effect of foliar application of Boron and soil application of Zinc levels on growth and yield of Cowpea (*Vigna unguiculata*). The Pharama Innovation Journal. 2021;10(8): 1341-4.
13. Barbara M Humtsoe., Joy Dawson and Praveena Rajana. 2018. Effect of nitrogen, boron and zinc as basal and foliar application on growth and yield of maize (*Zea mays* L.).
14. Ravichandra K, Naga Jyothi Ch, Jaipal Singh B, Joy Dawson, Krupakar A. Growth of groundnut (*Arachis hypogaea* L.) and its yield as influenced by foliar spray of boron along with rhizobium inoculation. Indian Journal of Dryland Agricultural Research and Development. 2015;30(1):60-63.
15. Hamouda HA, Anany TG, El-Bassyouni MSS. Growth and yield of dry bean (*Phaseolus vulgaris* L.) as affected by Zn and B foliar application. Middle East Journal of Agriculture Research. 2018;7(2): 639-649.
16. Nishant Srivastava, Joy Dawson, Ritesh Kumar Singh. Interaction effect of spacing, sources of nutrient and methods of zinc application on yield attributes and yields of green gram (*Vigna radiata* L.) in NEPZ. Journal of Pharmacognosy and Phytochemistry. 2017;6(4):1741-1743.
17. Abid Khan, Zafar Hayat, Asad Ali Khan, Junaid Ahmad, Muhammad Waseem Abbas, Haq Nawaz, Farhan Ahmad, Kaleem Ahmad. Effect of foliar application of zinc and boron on growth and yield components of wheat. Agricultural Research and Technology: Open Access Journal. 2019; 21(1): 3-6.
18. Debnath P, Pattanaaik SK, Sah D, Chandra G, Pandey AK. Effect of boron and zinc fertilization on growth and yield of cowpea (*Vigna unguiculata* L. Walp.) in Inceptisols of Arunachal Pradesh. Journal of the Indian Society of Soil Science. 2018;66(2):229-234.
19. Durgude AG, Kadam SR, Pharande AL. Response of hybrid maize to soil and foliar application of iron and zinc on entisols. An Asian Journal of Soil Science. 2014;9(1): 36-40.
20. Christopher Aboyeji, Oluwagbenga Dunsin, Aruna O Adekiya, Chinomnso Chinedum, Khadijat O Suleiman, Faith O Okunlola, et

- al. Zinc sulphate and boron-based foliar fertilizer effect on growth, yield, minerals, and heavy metal composition of groundnut (*Arachis hypogaea* L.) grown on an Alfisol. International Journal of Agronomy. 2019; 53(4):1-7.
20. Evangeline Marngar, Joy Dawson. Effect of biofertilizers, levels of nitrogen and zinc on growth and yield of hybrid maize (*Zea mays* L.). International Journal of Current Microbiology and Applied Sciences. 2017; 6(9):3614-3622.
21. Pabitra Adhikary, Soma Giri, Ananda Hansda, Sukanta Saren, Babulal Tudu. Effect of boron on growth and yield of lentil in alluvial soil. International Journal of Pure and Applied and Bioscience. 2018;6(5): 1171-1175.
22. Raj AB, Raj SK, Prathapan K, Radhakrishnan NV, Swadija OK. Effect of seed invigouration on yield enhancement in grain cowpea [*Vigna unguiculata* ( L.) walp]. Legume Research- An International Journal. 2021;44(9):1118-23
23. Ram Pratap. Integrated use of fertilizers and manures under varying levels of zinc in pearl millet (*Pennisetum glaucum* (L) R. Br. emend stuntz) and their residual effect on mustard, Thesis, Rajasthan Agricultural University, Bikaner; 2006.
24. Yadav BN, Umesha C, Sanadiya LK. Influence of Sulphur and Zinc on Growth and Yield of Cowpea (*Vigna unguiculata* L.). International Journal of Environment and Change. 2022;12(10):93-7.

© 2023 Kundu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/105637>