



Development of Phyto-cosmetic Herbal Sindoor from *Rosa indica* L. and Its Characterization

**Akanksha Singh^a, Meenu Verma^a, Vaishali Mishra^a,
K. J. Singh^b, S. K. Tewari^b and Mahesh Pal^{a*}**

^a Phytochemistry Division, CSIR-National Botanical Research Institute, Lucknow-226001, India.
^b Botanical Garden Division, CSIR-National Botanical Research Institute, Lucknow-226001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIMPS/2023/v12i4233

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106933>

Original Research Article

Received: 25/07/2023

Accepted: 30/09/2023

Published: 14/10/2023

ABSTRACT

Aim: The objective of this research study was to develop a liquid herbal sindoor containing flower petal extract of *Rosa indica* L., and to evaluate the stability, antioxidant and antibacterial activities of the developed formulation.

Place and Duration of Study: The formulation was prepared in the Phytochemistry Division of CSIR- National Botanical Research Institute. The duration of the study period was 90 days.

Methodology: The crude ethanolic extract of *R. indica* flowers was incorporated into a base for the preparation of liquid herbal sindoor. The antioxidant activity of this herbal product was evaluated according to the DPPH radical scavenging method, and the stability study of the formulation was evaluated for 90 days. The antibacterial activity was performed on those strains of bacteria, which are responsible for several skin infections using Disc Diffusion Assay. The HPLC- DAD analysis

*Corresponding author: E-mail: m.pal@nbri.res.in, drmpal.nbri@rediffmail.com;

identified the anthocyanins cyanidin 3,5-diglucoside and pelargonidin 3,5-diglucoside in *R. indica*, flower extract.

Results and Discussion: The formulation showed antioxidant activity and its stability was assessed in terms of pH values and color stability of the flower extract. It was observed that both these properties did not show any significant changes.

Conclusion: The prepared herbal liquid sindoor was found safe to use having a potential to substitute the toxic lead tetraoxide (Pb_3O_4) containing sindoor available in the market.

Keywords: *Rosa indica*; flower extract; formulation; antioxidant; antibacterial; pH value and color stability; lead tetraoxide.

1. INTRODUCTION

The use of phyto-cosmetics or herbal cosmetics has increased in personal care products. There is a high demand for cosmetics, having additional raw plant-derived ingredients as they are safe, non-toxic, and environment friendly. On the other hand, the overuse of products made of synthetic materials has put human health at risk and led to a number of health disorders. Sindoor is a traditional vermilion red or orange-red coloured cosmetic powder, usually worn by married women along the part of their hairline in the Indian subcontinent. Sindoor is a crucial cosmetics item in India for married ladies, worship, and other uses. Long-term usage of synthetic sindoor containing lead tetraoxide (Pb_3O_4) has shown symptoms of skin cancer, edema, hair loss, and greying of the hair [1]. Lead is one of the neurotoxins that has been the subject of the most recent and intensive research. Over time, it accumulates like a typical poison [2]. In light of the situation mentioned above, a process technique has been developed to produce an alternative, safe, non-toxic, ecofriendly, natural dye based Sindoor. The process provides an opportunity for the preparation of red color liquid herbal Sindoor using *Rosa indica* as a raw material.

Rosa indica L. is a perennial flower shrub under the Rosaceae family, which contains herbs, shrubs or trees that are thorny, rhizomatous or climbing [3]. Rose is used in a wide range of ways. It is traditionally been used in folk medicines, particularly in the field of skincare. It is appropriate for all skin types, although it is particularly beneficial for dry, sensitive, or aging skin. It helps to reduce the redness brought on by swollen capillaries since it has a tonic and astringent impact on the capillaries right below the skin's surface [4]. There are many important groups of compounds of rose that play an important role in the antibacterial activity of rose. Among them, the groups of compounds with

antibacterial activity are flavonoids, terpenes, anthocyanins. Flavonoids are classified as natural plant compounds, generally secondary metabolites with different phenolic structures [5]. Rose flower is one of the best sources for antibacterial activity against microorganisms. Strong antibacterial activity against test pathogens such as *E. coli*, *P. aeruginosa*, and *S. aureus* is shown by ethanolic extracts of rose petals, leaves, and stems [6]. In addition to their antibacterial and antifungal properties, flavonoids are also responsible for the aroma and color of flowers [5].

Due to the presence of antibacterial and antioxidant activities *R. indica* extract may be useful for the synthesis of herbal drugs in the future [7]. Natural sources with high anthocyanin contents are becoming more and more popular in the food and medical industries for making dietary supplements for therapeutic and nutritional purposes [8].

To find safe cosmetic alternatives in place of toxic traditional products, this work aims to develop an antioxidant formulation containing ethanolic extract of the petals of *R. indica* and, evaluate its stability and antibacterial activity.

2. MATERIALS AND METHODS

2.1 Plant Material and Preparation of Crude Ethanolic Extract

Flowers of *R. indica* were collected from the temples of Lucknow, Uttar Pradesh, India, in the month of July, 2021. The raw material was authenticated in the Herbarium of CSIR- NBRI, Lucknow (LWG). Petals were separated from the flowers and weighed. The petals were macerated with ethanol (100%) for 24 hours. The ethanolic extract obtained was filtered and concentrated under reduced pressure on rotary evaporator.

$$\text{Extract yield (\%)} = (W_1/W_2) \times 100$$

Where, W_1 is the weight of extracted plant residues in grams, and W_2 is the weight of rose petals in grams. The yield of the extract obtained was 15.6 %.

2.2 Anthocyanin Characterization Using HPLC- DAD Analysis

HPLC analysis of ethanolic extract of *R. indica* was performed on Shimadzu prominence system equipped with an autosampler (SIL-20 AC), LC Solution 1.0 software and DAD detector. The plant extract was analyzed in triplicate at the concentration of 1mg/ml as described by Huihua Wan et al. [9] under the following conditions: stationary phase: C_{18} column (dimensions 250mm x 4.6mm and particle size of 5 μ m); mobile phase: solution A: water + 0.5% (v/v) formic acid and solution B: Acetonitrile (ACN), considering the elution gradient as shown in Table 1. The injection volume and the flow rate were set at 10 μ l and 0.5ml/min respectively at the constant temperature of 25°C. Before, analysis the solutions were degassed and filtered through 0.22 μ m filter membrane. The wavelength was set at 254 nm for analysis. All the reagents used for chromatographic analysis were of HPLC grade.

2.3 Development of Formulation of Liquid Herbal Sindoor

For the development of the formulation the compatibility between the components and the flower extract has to be studied. The components used were as follows: bees wax, sunflower wax, castor oil, coconut oil and flower extract. These components were mixed together in petridish, kept on a hot plate at the temperature of 50°C. Then the herbal formulation was subjected to stirring on magnetic stirrer for 7 to 8 hours at 45°C.

2.4 Stability Studies

The stability of the colour of the formulation was analyzed on 0, 30, 60 and 90 days using a colorimeter (High- Quality Spectrophotometer

NS800). Using a colorimeter, the colours are expressed numerically in accordance with international standards. Anyone may comprehend what colour is being conveyed in this way. In order to more precisely note the changes in colour, the data from the colorimeter could enable the identification of different samples by any single quality (lightness, chroma, or hue). The informational value of the data is substantially enhanced by the capacity to appropriately define different colours [10]. The following parameters— L^* , a^*b^* , C^* , and h° —were monitored in order to accurately assess the formulation's colour.

A (previously calibrated) digital pH meter of LABMAN (Model no: LMPH- 12) was used for measuring pH of the sample. Before measurement, the samples were diluted to 10% (w/v) in distilled water [11].

2.5 Antioxidant Activity

The DPPH radical-scavenging activity of the samples was measured by the method described by Blois [12] with a slight modification using gallic acid and ascorbic acid as reference. The formulation prepared from ethanolic extract of petals of *R. indica* was analyzed for radical-scavenging activities. Briefly, the stock solution of formulation having concentration 1mg/ml was prepared. Various concentrations (20 μ l, 40 μ l, 60 μ l, 80 μ l, 100 μ l) of formulation were prepared by adding both methanol (80 μ l, 60 μ l, 40 μ l, 20 μ l) and 1mL of a DPPH ethanolic solutions. The mixtures were vortexed and incubated for 30 min at room temperature in the dark. The absorbance of mixtures was determined by using a spectrophotometer at 517 nm. UV-Vis absorption spectra were measured on a Perkin Elmer Lambda 35 UV/VIS spectrophotometer (WIN LAB V6 STD Software). The free radical scavenging activity was expressed in percentage using the following formula:

DPPH radical scavenging activity (%) = $\frac{\text{absorbance control} - \text{absorbance of sample}}{\text{absorbance control}} \times 100$

Table 1. Gradient elution method for HPLC analysis of anthocyanin in *Rosa indica*

Time (min)	Solution A: water+ 0.5% (v/v) formic acid (% v/v)	Solution B: Acetonitrile (ACN) (% v/v)
0	95	05
5	90	10
30	81	19
50	60	40
50.01- 60	95	05

Table.2 Ingredients with their composition (in percentage) in formulation

S.No.	Ingredients	Composition (in %)
1	Bees wax	25
2	Sunflower wax	20
3	Castor oil	05
4	Coconut oil	10
5	Flower extract	40

2.6 Antibacterial Activity

2.6.1 Preparation of inoculums of microorganism for testing

Three pathogens *Escherichia coli*, *Bacillus pumilis*, and *Staphylococcus aureus* were used as test subjects to determine the antibacterial activity of rose petals. These three pre-isolated bacterial cultures were obtained from KGMU, Lucknow, Microbiology Division. The cultures were subcultured on NA slants and stored at 4°C until needed. For testing of the sample, the inoculums were prepared from the stock culture and subcultured into the nutrient broth (30ml) using a sterilized wire loop. The inoculate was further incubated overnight at 37°C in a rotary shaker. These inoculates were further stored at 4°C until usage.

2.6.2 Activity analysis (Disc Diffusion Assay)

The antibacterial activity of formulation and extract was done by Disc Diffusion Assay [13]. The antibacterial activity of formulation prepared from *R. indica* petals, was evaluated against three pathogenic bacterial strains by using agar well diffusion method under sterilized conditions. For this three NA plates were prepared for formulation and petal extract. 400µl inoculum of each selected bacterium was uniformly spread over agar plates with the help of sterilized glass spreader.

After five minutes three wells, approximately 7mm in diameter were bored with the help of borer. The equal volume (50 µl) of antibiotic

Tetracycline (100 mg/ml), formulation and petal extract (200mg/ml) were poured into the wells. The plates were incubated at 37 °C for 24 hrs in incubator. Next day, the results were observed and the antibacterial potential was measured in terms of diameter of zone of inhibition.

3. RESULTS AND DISCUSSION

With the purpose of obtaining the crude ethanolic extract, the fresh petals of *R.indica* were subjected to maceration with ethanol. The final solution, after being concentrated on a rotary evaporator, was subjected to HPLC analysis thereafter, a formulation was developed for preparation of liquid herbal sindoor. The prepared formulation was evaluated for various physical parameters as described in Table 3.

The analysis, of the flower extract of *R. indica* using high-performance liquid chromatography coupled to a DAD detector (HPLC–DAD), enabled the identification and quantification of cyanidin 3,5-diglucoside and pelargonidin 3,5-diglucoside, the two analytical markers previously reported in this plant species [14]. The retention time (Rt) of the two anthocyanins, 3,5-diglucoside and pelargonidin 3,5-diglucoside were 6.72 and 8.80 min. respectively. The analysis of the peak area percentage quantified the content of these anthocyanins in the crude ethanolic extract, the amount of 3,5-diglucoside and pelargonidin 3,5-diglucoside was 0.0255 and 0.0047 mg/ml, respectively. Fig. 1 presents the chromatographic profile of the crude ethanolic extract indicating the identified markers.

Table 3. Evaluation of physical parameters of herbal formulation

S. No.	Parameters	Observation
1	Appearance	Thick liquid
2	Color	Dark red
3	Texture	Smooth
4	Odour	Pleasant
5	Water washability	Washable




Fig. 1. HPLC Chromatogram of ethanolic extract of *R. indica* at the wavelength of 254 nm. (A = cyanidin 3,5-diglucoside, B = pelargonidin 3,5-diglucoside)

On a fresh weight basis the total anthocyanin content ranged from 0.24 to 578.10 mg/100g. The environmental factors such as temperature and light strongly influence the anthocyanin content of the rose petals [15]. Lin et al. [16] reported 0.025 to 18.69 mg/g of total content of anthocyanin in rose petals. Anthocyanins give plants a variety of physiological health benefits and disease preventive actions in addition to giving natural beautiful colors [17].

The pH analysis of the extract, base (castor oil, coconut oil, sunflower and bees wax) and formulation was performed during the study period for 0, 30, 60 and 90 days and no significant variation was observed. The observed pH values were around 4.2 showing biocompatibility with the skin, being potentially non-irritating. The pH gradient for the skin surface ranges from 4 to 6 [17]. The pH value of the used base ranged from 7.4 to 7.6. Finally, the pH value of the formulation ranged from 4.0 to 4.3 over the period of study (90 days).




Fig. 2. pH values of base, extract and formulation during the study period (0, 30, 60 and 90 days)

Table 4. The values of color parameters L* (indicating lightness), a*(redness), b*(yellowness), C*(chroma), h°(hue) measured using colorimeter

ΔT^*	Standard					Formulation					Difference between standard and formulation					
	L*	a*	b*	C*	h°	L*	a*	b*	C*	h°	ΔL	Δa	Δb	ΔC	Δh	ΔE
0	0.24	0.08	0.07	0.11	42.42	0.23	0.07	-0.05	0.09	324.51	-0.01	-0.01	-0.12	-0.02	-0.12	6.12
30	0.60	0.27	0.32	0.42	50.34	0.52	0.12	0.54	0.55	77.82	-0.08	-0.15	0.22	0.14	0.23	0.28
60	0.33	0.39	-0.95	1.03	292.3	0.31	0.35	-0.44	0.56	308.07	-0.01	-0.04	0.51	-0.47	0.21	0.51
90	0.29	0.09	-0.03	0.03	268.9	0.60	0.99	-1.18	1.54	509.89	0.31	0.90	-1.15	1.15	0.15	1.55

ΔT^* = Time interval in days




Fig. 3. Analysis of Free Radical Scavenging Activity of herbal formulation at different concentration using Gallic acid as standard

The color of the formulation was evaluated at the time interval of 0, 30, 60 and 90 days to observe any significant change in color as described in Table 4. The color parameters such as L^* (indicating lightness), a^* (redness), b^* (yellowness), C^* (chroma), h° (hue) were measured using colorimeter. While the parameter b^* is positive for yellowish colors and negative for bluish hues, the parameter a^* is positive for reddish colors and negative for greenish colors. Each hue can be thought of as being equivalent to a member of the greyscale that is in between black and white according to L^* , an approximation of brightness. Thus, each scale's L value represents the degree of lightness or darkness. The delta values (ΔL , Δa , and Δb) reflect the degree of variation between a standard and sample for L , a , and b respectively, and ΔE is the total color difference. L^* , a^* , and b^* all have delta values i.e. ΔL , Δa , and Δb respectively, that can be either positive (+) or negative (-). The sign of the delta value indicates whether the sample is redder or greener than the standard. Delta E (ΔE), is always positive [18-20]. The value of a^* and ΔE were always found to be positive (+) in each measurement. No significant change in color of the formulation was observed during the study period.

The antioxidant potential, of the formulation was tested according to the DPPH radical scavenging assay. The concentration of the sample necessary to scavenge 50% of the DPPH free radicals, or the IC_{50} value, was computed from the graph. The term " IC_{50} " refers to the quantity or concentration of extracts required to neutralise 50% of free radicals [8]. Polyphenols present in the extract have a major role, in the antioxidant activity. The antioxidant potential of the formulation according to DPPH assay has an IC_{50} value of $27 \pm 05 \mu\text{g/ml}$ in case where gallic acid was used as reference standard as described in Fig. 3.

The antioxidant activity of formulation was also determined using ascorbic acid as reference standard (Fig. 4) and the IC_{50} value was $32.82 \pm 03 \mu\text{g/ml}$.

Results of the test for antibiotic susceptibility were expressed as the diameter of the inhibition zone (in mm) of ethanolic extract and formulation against each bacterial strain employed. The zone of inhibition for the standard antibiotic (Tetracycline) against each strain was also recorded. Data of antibacterial activity of ethanolic extract and formulation are demonstrated in Table 5:




Fig. 4. Analysis of Free Radical Scavenging Activity of herbal formulation at different concentration using Ascorbic acid as standard

Table 5. Antibacterial susceptibility assay of *R. indica* petals extracts and herbal formulation

Tested Bacteria	Zone of Inhibition (in mm)		
	Ethanolic Extract	Herbal formulation	Tetracycline
<i>E. coli</i>	22.0 ± 1.0	23.0 ± 1.0	31.0 ± 2.0
<i>S. aureus</i>	18 ± 1.0	17 ± 0.0	30.0 ± 1.0
<i>B. pumilis</i>	20 ± 0.0	21.0 ± 1.0	34.0 ± 1.0

4. CONCLUSION

In this research efforts were made to illustrate the compatibility between the base used and the crude ethanolic extract of petals of *R. indica*. The stability study of the formulation at room temperature and pressure showed no significant change in the pH and colour of the sample. The formulation also showed antioxidant activities carried out by DPPH radical scavenging method using gallic acid and ascorbic acid as standard. The anti-bacterial activity on various strains of bacteria (*S. aureus*, *B. pumilis*, and *E. coli*) indicated the efficacy of the formulation against such bacterial strains and demonstrated the antibacterial stability of the formulation. All these studies confirmed the anti-irritant, anti-allergic and antibacterial properties of the prepared formulation (liquid herbal sindoor). Thus this herbal product with no side effects may be used as substitute of lead tetraoxide (Pb_3O_4) containing sindoor available in the market. However, in future, further studies are still required to develop such herbal based products using *R.indica* and to discover their beneficial health effects.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENT

The authors are thankful to the Director, CSIR-National Botanical Research Institute, Lucknow, India for facilities and encouragement. The financial support received from the Council of Scientific and Industrial Research, New Delhi under the project 'Floriculture Mission (HCP-0037)' is duly acknowledged. This manuscript has an Institutional communication number: 'CSIR-NBRI_MS/2022/11/04'.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shivam Samariya, Sumeet Dwivedi, Shweta Patil, Debadash panigrahi and hemant joshi: Formulation and evaluation of herbal sindoor using different natural/herbal ingredients. International Journal of Pharmacy Teaching & Practices. 2013;3:752-754
2. Durgesh Wadhwa, Gopal Arora. Identification of lead from sindur samples. International Journal of Innovative Research in Engineering & Management (IJIREM). 2022;9:357-360
3. Crepet WL, Nixon KC, Gandolfo MA. Fossil evidence and phylogeny: The age of major angiosperm clades based on mesofossil and macrofossil evidence from cretaceous deposits. American Journal of Botany. 2004; 91: 1666–1682
4. Thakare Priya, Ashok Rao, Deshbhratar Kiran, Suryawanshi MN. A brief review on therapeutic effects of – “ornamental plant” rose. International Journal of Ayurveda and Pharma Research. 2017;5:46-52
5. Yusra Safdar, Taqdees Malik. Antibacterial activity of the rose extract. Open Access Journal of Complementary and Alternative Medicine. 2020;2:194-201
6. Mishra RP, Mohammad Arshad, Abdul Sami. Antibacterial Properties of *Rosa indica* (L.) Stem, Leaves and Flowers. Journal of Pharmaceutical and Biomedical Sciences (JPBMS). 2011;12:1-3.
7. Perna, Prince sharma, Sanjana Chaudhary, Jyoti Tyag. A review on antimicrobial and antioxidant activity of rosa damascene against different species of microbes. International Journal of Advance Research and Innovative Ideas in Education. 2021;7:3555-3568
8. Jin Hwan Lee, Hyeon-Jin Lee, Myoung-Gun Choung: Food Chemistry. 2011;129: 272-278.
9. Huihua Wan, Chao Yu, Yu Han, Xuelian Guo, Sagheer Ahmad, Aoying Tang, Jia Wang, Tangren Cheng, Huitang Pan, Qixiang Zhang. Flavonols and carotenoids in yellow petals of rose cultivar (*Rosa* 'Sun City'): A possible rich source of bioactive compounds. Journal of Agricultural and Food Chemistry. 2018;66:4171–4181
10. Mohammed Azhar Bintory, Seetharamu GK, Munikrishnappa PM, Ramegowda GK, Basavaraj G. Evaluation of the Colour of Dried Dutch Rose Flowers Using a Colorimeter. Journal of Horticulture. 2015;2.
11. Rafaela Santos de Melo, Silvio Alan Gonçalves Bomfim Reis, Amanda Leite Guimarães, Naiane Darklei dos Santos Silva, Joao Miguel Rocha, Noureddine El Aouad, Jackson Roberto Guedes da Silva Almeida. Phytocosmetic Emulsion Containing Extract of *Morusnigra* L. (*Moraceae*): Development, stability study,

- antioxidant and antibacterial activities. *Cosmetics*. 2022;9:1-12.
12. Blois MA. Antioxidant determination by the use of a stable free radical. *Nature*. 1958; 181: 1199–1200.
 13. Amit Alexander Charan, Prerak Gupta. Comparative analysis of antibacterial, antioxidant and photosynthetic activity of *Azadirachta indica*, *Rosa indica* and *Moringa oliefera* cultivars. *International Journal of Current Research*. 2013;5:556-561.
 14. Huihua Wan, Chao Yu, Yu Han, Xuelian Guo, Le Luo, Huitang Pan, Tangchun Zheng, Jia Wang, Tangren Cheng and qixiang zhang: Determination of flavonoids and carotenoids and their contributions to various colors of rose cultivars (*Rosa* spp.). *Frontiers in Plant Science*. 2019;10. Article 123.
 15. Poonam Kumari, Raju DVS, Prasad KV, Kanwar Pal Singh, Supradip Saha, Ajay Arora, Firoz Hossain. Quantification and correlation of anthocyanin pigments and their antioxidant activities in rose (*Rosa hybrida*) varieties. *Indian Journal of Agricultural Sciences*. 2017;87:76-82.
 16. Lin L, Hyeonmi H, JeehyeS, Younghwa K, Heon-Sang JL. Antioxidant activities of methanolic extracts from four different rose cultivars. *Journal of Food and Nutrition Research*. 2014;2:69–73.
 17. Serup J, Jemec GBE, Grove GL. (Eds.) *Handbook of Non-Invasive Methods and the Skin*, 2nd ed.; Taylor & Francis. Boca Raton, FL, USA. 2006;1056.
 18. Deepali Singhee, Adrija Sarkar. colorimetric measurement and functional analysis of selective natural colorants applicable for food and textile products. *IntechOpen*. 2022;1-33.
 19. Akshada Kakade, Divya Yadav, Pranoti Bhojnale, Pallavi Velapure, Akshay Vedpathak, Prajyoti Hiware. A new approach of formulation and evaluation of herbal sindoor powder and stick from herbal Ingredients. *European Journal of Biomedical and Pharmaceutical Sciences*. 2016;3:251-253.
 20. Sheema Bai, Leena Seasotiya, Anupma Malik, Pooja Bharti, Sunita Dalal: Bioactive compounds and pharmacological potential of *Rosa indica* L. and *Psidiumguajava* L. methanol extracts as antiurease and anticollagenase agents. *Der Pharmacia Lettre*. 2015;7:179-184.

© 2023 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106933>