



International Journal of Environment and Climate Change

Volume 13, Issue 11, Page 1936-1941, 2023; Article no.IJECC.107777

ISSN: 2581-8627

(Past name: British Journal of Environment & Climate Change, Past ISSN: 2231-4784)

Assessment on Endangered Plant Biodiversity of Shirui National Park, Ukhrul, Manipur, India

Yumlembam Lichingamba Meitei ^{a++*} and Abhishek James ^{a#}

^a Department Environment Science and NRM, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj- 211007, U.P, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI:10.9734/IJECC/2023/v13i113351

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107777>

Original Research Article

Received: 19/08/2023

Accepted: 25/10/2023

Published: 31/10/2023

ABSTRACT

Five different random sites were analyzed for the study of the plant species using quadrat methods from Shirui National Park. The plants were measured and recorded by random sampling with 20 quadrats of 10m x10m in each study site during the period between January 2023 and April 2023. Site 4 Showed the maximum parameter for IVI value of Plant Species which was represent by *Bidens Pilosa* with the value of 130.7 Site 5 Showed the minimum parameter for IVI value of Plant Species which was represent by *Liliummackliniae* with the value of 24. The IVI results show that the plant species importance values differs from site to site. From the study *Lilium Mackliniae* (Shirui Lily) is found decreasing in population, a species which is only endemic to Shirui National Park, Manipur.

Keywords: Conservation; biodiversity; shirui lily; endangered; Ukhrul.

⁺⁺M.Sc;

[#]Assistant Professor;

^{*}Corresponding author: E-mail: lichingyum@gmail.com;

1. INTRODUCTION

Manipur is a land of Blue Hills and green valley. It is in the extreme northeastern border of India. According to 2011 census the total population is 28,55,794 lakhs. Different communities having their own language and culture are residing in the state. But the main language in the state is Meitei language. Shirui lily is the state flower of Manipur and this flower is found only in Ukhrul district. The Government of Manipur organized a festival in the name of this state flower called "Shirui festival" as a state function from 2017 onwards. The main theme of the festival is to conserve the flower as well as unite together all the communities through cultural exchange programmed and enhances the tourism industry in the state [1].

Biological diversity refers to the variety and variability among living organisms and the ecological complexes in which they occur [2]. Anthropogenic activities are now-a-days prominent and causing extinction of many plant species of ecological and economic significance [3].

Understanding of the dynamics of plant resources is important for their sustainable management, utilization and biodiversity conservation [4]. Forest plants are gathered for food, herbal medicine, spices, fuel, etc, which have built up their socio economic and cultural life [5]. The forests are the main repository of biodiversity and play an important role in maintaining the ecological balance of nature. The status of species diversity reflects the health of the ecosystem. Waponungsang et al. [6].

The aim of the study is to understand and assess the population structure of the endangered plant biodiversity of Shirui National Park, Manipur. *Lilium mackliniae*, state flower of Manipur is famous for its unique habitat, endemic to Shirui National Park, Manipur. The Northeastern region is also endowed with several plant and animal species of immense ecological and economic importance. These sources are under tremendous pressure and facing threats towards extinction.

2. MATERIALS AND METHODS

For the study of the plant species five random different sites selected from Shirui National Park, quadrat methods were used. The plants were measured and recorded by random sampling

with 20 quadrats of 10m x10m in each study site. All plants species recorded within each quadrat were identified then counted and estimated their cover percentage. The Importance Value Index (IVI) for the plant species was determined as the sum of the relative frequency, relative density and relative dominance [7].

2.1 Sampling Sites

Shirui National Park, Ukhrul, Manipur within 24°13'51.0"N 94°13'51.0"E latitude and longitude at an elevation of 2,835 meters above sea level, with annual average rainfall of 1500 to 2000 millimeters. The climate of Shirui National Park is influenced by its Northeastern location and its elevation of Eastern Himalayas. In winter the climatic condition fall under 0°C to 10°C (32°F to 50°F) and 20°C to 30°C (68°F to 86°F) during summer.

3. RESULTS AND DISCUSSION

On the survey area of the Shirui National Park comprised of 10 species of plant species with a total number of 1607 individual belong to 7 families were found.

From the given Table 1 Plant species having highest Frequency from each sites includes *Lilium mackliniae* (90) from site 1, *Curcuma rubescens* (90) from site 2, *Bidens pilosa* (100) from site 3, *Bidens pilosa* (90) and *Artemisia Nilagirica*(95) from site 5. Plant species having highest density from each sites includes *Lilium mackliniae* (6.1) from site 1, *Curcuma rubescens* (5.1) from site 2, *Bidens pilosa* (7.9) from site 3, *Bidens pilosa* (4.6) from site 4 and *Artemisia Nilagirica* (6.1) from site 5. Plant species having highest abundance from each sites includes *Lilium mackliniae* (6.4) from site 1, *Lilium mackliniae*(6) from site 2, *Bidens pilosa* (7.9) from site 3, *Bidens pilosa*(5.1) from site 4 and *Artemisia Nilagirica*(6.4) Plant species having lowest Frequency from each sites includes *Dendrobium densiflorum*, *Dendrobium nobile* and *Rhododendron arboretum* having the same value (70) from site 1, *Dendrobium densiflorum* (70) from site 2, *Lilium mackliniae* (80) and *Plantago erosa* (80) from site 3 and *Lilium mackliniae* (45) from site 5. Plant species having lowest density from each sites includes *Rhododendron arboretum* (1.8) from site 1, *Dendrobium nobile* (2.4) from site 2, *Lilium mackliniae* (3.7) from site 3, *Lilium mackliniae*(2.3) from site 4 and *Lilium mackliniae* (45) from site 5. Plant species having lowest

Abundance from each sites includes *Rhododendron arboretum* (2.5) from site 1, *Dendrobium nobile* (3) and *Rhododendron arboretum* (3) from site 2, *Lilium mackliniae* (4.6) from site 3, *Plantago major* (3.3) from site 4 and *Lilium mackliniae* (1.8) from site 5

From the given Table 1 it can be concluded that the value for frequency, density and abundance of the plant species are low in the study sites. Aigbe and Omokhua, [8] reported that the population density and abundance of a tree species in a patch of rainforest is substantially affected by the atmospheric conditions inside the forest and the number of seeds capable of surviving generated by the tree in question. Christie and Armesto [9] gave an account of very low population densities and abundance of a vast

number of economically viable tree species triggered by death of viable seeds and poor micro-sites for rejuvenation.

The plant species differ are from site to site, however *Lilium mackliniae* occurred to be found is all the five sites of the study area but the value of each site are differ from one another, *Biden pilosa* and *Plantago major* and *Rhododendron arboretum* are occurred to be found in three different site, *Curcuma rubescens*, *Dendrobium densiflorum*, *Dendrobium nobile* are occurred to be found in two different sites and *Plantago erosa*, *Artemisia Nilagirica*, *Eupatorium Adenophorim* are occurred to be found in only one site. Ojo [10] revealed the occurrence of high number of species in these families may be as a result of their mechanism of seed dispersal such

Table 1. Quantitative analysis of plant species of Site 1,2,3,4 and 5 are as categorized below

Species	F	D	A	RF	RD	RDo	IVI
Site 1							
<i>Curcuma rubescens</i>	85	4.5	5.3	22	26.7	24.7	73.6
<i>Dendrobium densiflorum</i>	70	2.4	3.5	18.1	13.9	15.8	48.2
<i>Dendrobium nobile</i>	70	2.3	3.5	17.1	13.6	16.3	46.8
<i>Rhododendron arboretum</i>	70	1.8	2.5	18.1	10.7	11.8	40.7
<i>Lilium mackliniae</i>	90	6.1	6.4	23.3	35.5	31.2	90.5
Site 2							
<i>Curcuma rubescens</i>	90	5.1	5.6	22.2	28.7	26	77.1
<i>Dendrobium densiflorum</i>	70	2.8	4	17.2	16.10	18.7	52.1
<i>Dendrobium nobile</i>	80	2.4	3	19.7	13.4	13.8	47.1
<i>Rhododendron arboretum</i>	85	2.5	3	20.9	14	13.8	49.2
<i>Lilium mackliniae</i>	80	4.8	6	19.7	27	27.6	74.4
Site 3							
<i>Bidens pilosa</i>	100	8	7.9	28.9	40	35.3	104
<i>Plantago major</i>	85	4.2	5	24.6	21	22	68.04
<i>Lilium mackliniae</i>	80	3.7	4.6	23.1	18.7	20.7	62.7
<i>Plantago erosa</i>	80	3.9	4.9	23.1	19.7	21.8	64.8
Site 4							
<i>Bidens pilosa</i>	90	4.6	5.1	39.13	48.9	42.66	130.7
<i>Plantago major</i>	75	2.5	3.3	32.61	26.5	27.5	86.44
<i>Lilium mackliniae</i>	65	2.3	3.6	28.26	24.4	29.8	82.85
Site 5							
<i>Eupatorium Adenophorim</i>	90	3.8	4.2	18.5	23.9	22.8	65.34
<i>Lilium mackliniae</i>	45	0.8	1.8	9.2	5.2	10.06	24.6
<i>Plantago Major</i>	80	1.6	2.0	16.5	9.9	10.66	37.12
<i>Bidens Pilosa</i>	90	1.8	2.0	18.5	11	10.66	40.43
<i>Artemisia Nilagirica</i>	95	6.1	6.4	19.5	38	34.5	92.4
<i>Rhododendron Arboretum</i>	85	1.8	2.1	17.5	11	11	40.03

D- Density, F- Frequency, RF- Relative Frequency; RD- Relative Density; RDo- Relative Dominance; IVI- Importance Value Index




Fig. 1. Map of Ukhrul District and Shirui National Park

as explosive mechanism and wind dispersal, where they germinate when conditions are suitable. However, less number of species noticed in some families may be as a result of poor germinability as seeds may require scarification or changes in thermal or light conditions to break dormancy or quiescence for germination to occur. Pausas and Austin. [11] reported such need on species richness in connection to the environment.

From the given Table 1 the highest Important Value Index of the plant species from each site includes *Lilium mackliniae* (90.5) from site 1, *Curcuma rubescens* (77.1) from site 2, *Bidens pilosa* (104) from site 3, *Bidens pilosa* (130.7) from site 4 and *Artemisia Nilagirica* (92.4) from site 5. The lowest Important Value Index of the plant species from each site includes *Rhododendron arboretum* (40.7) from site 1, *Dendrobium nobile* (47.1) from site 2,

Liliummackliniae (62.7) from site 3, *Lilium mackliniae* (82.85) from site 4 and *Liliummackliniae* (24.6) from site 5.

Importance Value Index (IVI) is used as a quantitative parameter that can express the level of dominance (level of control) of species in a community. The plant species that have the highest IVI are due to their higher density, frequency, and dominance than other types. The higher the IVI of a species, the higher the level of control in the community of that species grows, Similar result was conducted by another Author Sosilawatyet al [12] on the other study [13] revealed that the IVI values show variation in ecologically dominant from site to site. Hence, the IVI has helped in understanding the ecological significance of the species in the tropical dry deciduous forest[14].

4. CONCLUSION

In this present study, it revealed that the plant species were mostly low in their distribution and scantily populates the study area; this may result from the unrestrained exploitation. For reason, annually during the month of January the villagers burn down the main core area of grassland in order to minimize the other competitive species of *Lilium Mackliniae*, this result to the declination of other plant species. To make it possible to prevent the continuous decline of various other plant species; there is need for more effective measures of the forest by the indigenous communities.

ACKNOWLEDGEMENTS

The author is highly grateful to official of Department of Forestry, Manipur, India for proving the support for completing the research activities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Devi C. Remembering Shirui lily festival in Manipur, An International Journal of Creative Research Thoughts (IJCRT), 2021;9(1). ISSN: 2320-2882.
2. Pullaiah T, Bahadur B, Krishnamurthy KV. Plant Biodiversity: Plant Diversity,

Organization, Function and Improvement.2015;1

DOI 10.1007/978-81-322-2286-6_6

3. Choudhury B, Khan M. Conservation and management of endangered plant species: A case study of Northeast India, Global Science Books. 2010;4(1):47 – 53.
4. Sarkar M, Devi A. Assessment of diversity, population structure and regeneration status of tree species in Hollongapar Gibbon Wildlife Sanctuary, Assam, Northeast India, Tropical Plant Research an International journal. 2014.ISSN (E): 2349 – 1183, ISSN (P): 2349 – 9265.
5. Wangcha A, Konyak W. Survey of ethnomedicinal plants and its uses by the konyak tribe in Mon district, Nagaland, India, International Journal of Ecology and Environmental Science.2021;3(1).
6. Wapongnungsang. Ovung E, Tripathy SK. Assessment of tree diversity in tropical moist deciduous forest of Mizoram University, Northeast India, Journal of Applied and Natural Science;2021.ISSN : 0974-9411.
7. Cottam G, Curtis JT. The use of distance measurement in phytosociological sampling. Ecology. 1956;37:451-460.
8. Aigbe HI, Omokhua GE. Tree species composition and diversity in Oban Forest Reserve, Nigeria. Journal of Agricultural Studies. 2015;3(1):10-24.
9. Christie DA, Armesto JJ. Regeneration at micro site and tree species co-existence in temperate rainforest of Chilue Island, Chile, Journal of Ecology. 2003;91:776-784.
10. Ojo LO. The fate of a tropical rainforest in Nigeria; Abeku sector of Omo Forest Reserve, Global Nest: The International Journal. 2004;6(2):116-130.
11. Pausas JG, Austin MP. Patterns of plant species richness in relation to different environments: An appraisal, Journal of Vegetation Science. 2001;12:153-166.
12. Sosilawaty. Hastari, B., and Siregar, R. Composition and diversity of plant species in the two types of land cover of natural laboratory of hampangen educational forest, International Journal Of Biology And Biomedical Engineering .2021;15 DOI:10.46300/91011.2021.15.16
13. Narayan C. Anshumali Diversity indices and importance values of a tropical deciduous forest of Chhotanagpur plateau, India Journal of Biodiversity and Environmental Sciences (JBES) ISSN:

- 2220-6663 (P) 2222-3045. 2015;7(1):358-367, species in ten communal forest of Northern cross river state, Nigeria, International Journal of Current Research. 2017;9(10):59581-59596.
14. Ihoro A, Aja E. Population Frequency, Density, Abundance and Diversity of Tree

© 2023 Meitei and James; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107777>