



# Nutritional Value and Amino Acid Profile of Dried Seeds of *Cucumeropsis mannii* (Cucurbitaceae)

Joelle Lasm Agnei <sup>a</sup>, Yolande Dogoré Digbeu <sup>a\*</sup>,  
Edmond Ahipo Dué <sup>b</sup> and Kouakou Brou <sup>a</sup>

<sup>a</sup> Laboratory of Nutrition and Food Security, Nangui Abrogoua University, 02 BP 801 Abidjan 02, Côte d'Ivoire.

<sup>b</sup> Laboratory of Biocatalysis and Bioprocessing, Nangui Abrogoua University, 02 BP 801 Abidjan 02, Côte d'Ivoire.

## Authors' contributions

This work was carried out in collaboration among all authors. Author EAD and KB designed the study, performed the statistical analysis, wrote the protocol and supervised the work. Author JLA managed the analyses of the study. Author YDD wrote the first draft of the manuscript, managed the literature searches and edited the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJBCRR/2023/v32i10842

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109439>

Original Research Article

Received: 17/09/2023

Accepted: 23/11/2023

Published: 12/12/2023

## ABSTRACT

**Aims:** To assess the biochemical, phytochemical and amino acid content from *Cucumeropsis mannii* dried seeds flours grown in Côte d'Ivoire (West Africa).

**Study Design:** Experimental.

**Place and duration of study:** Sample: Laboratory of Nutrition and Food Security, Nangui Abrogoua University

**Methodology:** *Cucumeropsis mannii* seeds were obtained for this study from an experimental field in Gagnoa (Côte d'Ivoire). In March 2022, seeds were planted and after 7 months, the mature fruits that lead to the complete drying of the plant were harvested.

\*Corresponding author: E-mail: digbeuyolande@gmail.com;

**Results:** The biochemical composition of the seeds revealed that they contain varying proportions of carbohydrates, fibers, ash, total sugars with contents of  $9.28 \pm 0.79$  %;  $2.72 \pm 0.30$  %;  $3.12 \pm 0.38$ ;  $2.34 \pm 0.05$ , respectively. High levels of lipids, proteins, dry matter and energy were observed in seed flour with values of  $47.46 \pm 1.55$  %;  $31.80 \pm 1.16$  %;  $94.38 \pm 0.6$  %;  $591.46 \pm 1.17$  kcal /100 g DM. The resulting flour reached a pH of 6.5. The study of phytochemicals showed varying levels of oxalate (0.32 g/100g DM), tannins (3.02 g TAE/ 100g DM), total polyphenols (8.12 mg GAE / g DM), total flavonoids (27.5 mg QE/g DM) and carotenoids (2.9  $\mu$ g/100g DM). Furthermore, they were potential sources of essential amino acids, including histidine (905.6 mg/100g), isoleucine (844.5 mg/100g), threonine (730.1mg/100g), phenylalanine (550.0 mg/100g), valine (469.9 mg/100g), leucine (456.6 mg/100g), and methionine (358.5 mg/100g).

**Conclusion:** The overall result implies that *Cucumeropsis mannii* seeds could be added to the food preparations of malnourished children.

**Keywords:** *Cucumeropsis mannii*; phytochemicals; amino acids; lipids;protein.

## 1. INTRODUCTION

Agriculture is the basis of human development and could be an appropriate solution to the worldwide food shortage particularly in the developing countries. Today, recommendations from institutions and organisations, including the FAO, to increase the productivity and quality of the agricultural sector in developing countries focus on underutilised crops [1]. Among these are the cucurbits, which are plants with a high content of balanced protein and are cheap compared to animal protein. Indeed, the Cucurbitaceae are a remarkable family of plants that deserve attention for their multiple economic, cultural, medicinal and botanical importance. In some cultures, the seeds, leaves and roots are consumed. According to some studies, the seeds are rich in lipids, proteins and carbohydrates [2, 3]. Cucurbits have also been overlooked; however, underutilised plants in Africa are suitable for a wide range of agro-ecosystems and cropping systems [1]. Research by Zoro et al [4] identified five species of cucurbits in Côte d'Ivoire, including *Cucumeropsis mannii*. Known as the white-seeded melon, commonly called "n'vièle" in Côte d'Ivoire [4], it originates from tropical Africa, where it is prized for its oilseeds, grown for food and as a source of oil. However, *C. manni* seeds are still underexploited due to scientific ignorance about their nutritional potential. This research aims to determine the nutritional characteristics of *C. mannii* seeds grown in Côte d'Ivoir.

## 2. MATERIAL AND METHODS

### 2.1 Plant Collection

The biological material for this study was *Cucumeropsis mannii* seed which come from an

experimental field in the locality of Gagnoa (Côte d'Ivoire).

### 2.2 Preparation of Samples

*Cucumeropsis mannii* was harvested at leaf wilt to indicate fruit maturity and ripening. The latter underwent a hemispherical cut and were left to decompose for one or two weeks, until the pulp softened to facilitate the extraction of the seeds. The seeds were then separated from the fruit pulp, washed with water and dried in the oven at 45°C for 3 days. They were manually peeled and the almonds were finely ground to obtain powders that will be sieved (250  $\mu$ m diameter) and put in plastic bags for storage before the analysis tests.

### 2.3 Chemical Analysis

The pH of the dried seed powder was determined according to the method described in AOAC [5]. The AOAC method [6] was used to determine the moisture content after drying at 105°C for 24 hours to constant weight and the total ash content was determined by incinerating in a 550°C furnace. The AOAC method [6] was again used to determine crude fibre, and fat extraction in a Soxhlet apparatus for 8 hours using hexane for the determination of lipid content; while Nitrogen was done by Kjeldahl method and crude protein content subsequently calculated by multiplying the nitrogen content by a factor of 6.25. Ethanol-soluble sugars from *Cucumeropsis mannii* powder were extracted using the technique described by Martinez-Herrera et al., [7]; then total sugars were determined by Dubois *et al.* [8] using phenol and sulphuric acid. Available carbohydrate levels were determined using the difference method described by FAO [9] as following:

Total carbohydrate (%) = 100 % - [(%)moisture + (%) protein + (%) fat + (%) fibre + (%) ash]

The theoretical energy value was determined using the method outlined by Atwater and Rosa [10] from the specific coefficients for proteins, lipids and carbohydrates as following:

Energy value (Kcal/100g DM) = 4 x (%) protein + 9 x (%) lipids + 4x (%) carbohydrates

## 2.4 Antinutrients Analysis

Tannin were determined as reported by Bainbridge *et al.* method [11]. The principle of the assay is based on the fact that tannic acid in the presence of vanillin (0.1 mg/mL) and 70% of sulphuric acid gives a red coloration when the absorbance was measured at 500 nm. The tannins content of *C. mannii* was expressed as tannic acids equivalent (TAE). while oxalate and carotenoids were defined as indicated by AOAC [6] and FAO [12] respectively. Total polyphenols were measured according to Makkar *et al.*, [13]; Cicco *et al.*, [14] using the Folin Ciocalteu reagent. The absorbance is measured at 725 nm in the spectrophotometer and the results of polyphenols content were subsequently calculated as Gallic acid equivalent (GAE). As for flavonoids, they were determined as described by Meda *et al.* [15] using Quercetin as standard. The absorbance was measured at 415 nm; then flavonoid content of *C. mannii* was expressed as Quercetin equivalent (QE).

## 2.5 Amino Acid Analysis

The amino acid composition of sample was determined using a high-performance liquid chromatograph (HPLC), Applied Biosystems Model 172 A (Applied Biosystems, Foster City, California, USA) equipped with a RP-18 PTC column (2.1mm 22cm). Prior to injection, the proteins in sample were previously hydrolyzed by HCL 6 M with phenol (1%) at 150°C for 60 minutes in the Pico-Tag system (Waters, Milford, Mass., U. S.A.) The amino acid phenylisothiocyanate derivatives obtained were eluted by HPLC. Sodium acetate (105Mm, PH 4.6; 30%) and acetonitrile (70%) were used as elution buffers. A calibration chromatogram was established for the standard available amino acids.

## 2.6 Statistical Analysis

All measurements were carried out in triplicate. The results were processed by software package STATISTICA 7.1 (Stat soft Inc, Tulsa-USA, Head quarters). This results were expressed as means ± standard deviation. The statistical differences

among the means of data were calculating using one-way analysis of variance(ANOVA) and Duncan's Multiple Range Test (DMRT). Difference at  $P < 0.05$  were considered significant.

## 3. RESULTS AND DISCUSSION

### 3.1 Proximate Composition

The proximate composition of *Cucumeropsis mannii* seed flour was shown in table 1. The percentage protein of  $31,80 \pm 1,16$  was comparable to the value reported for melon seeds (31.1 %) and squash seeds (33.0 %) [16] but lower than the results of *C. mannii* seeds (37.4- 37.9 %) in Nigeria [17, 18]. Studies proved that, proteins were essential macronutrients for humans, with a key role in building and repairing the body [19]. The crude fibre of *C. mannii* seed (2.72 %) in this study was similar than those reported by Ogunbusola *et al.*, [17] and peanut seeds [20]. Adequate intake of crude fiber might have a helpful physiological role such as low incidence of colon cancer [21]. However, this low fibre content of powders from *Cucumeropsis mannii* seeds suggests that it would not be sufficient to meet the estimated human fibre needs of 14.2-17.5 g/day [22]. As a result, it should be combined with other fiber-rich dietary sources to prevent various chronic diseases [23]. The moisture content of *C. mannii* seed ( $5.62 \pm 0,6$  %) was also similar to  $5.53 \pm 0,35$  % [17], for *C. mannii* in Nigeria and previously comparable to  $5.01 \pm 0,01$  reported by Mohaammed *et al.*, [24]) in *Cucurbita maxima*. This low moisture content showed that the quality of this powder could protected it against the development of microorganisms. Indeed, according to Aryee *et al.* [25], the moisture content is a very important parameter in the storage of flours, since a moisture content greater than 12 % would promote the growth of microorganisms. *Cucumeropsis manii* powder showed a slightly acidic pH of 6.50, which certifies that the powder could be stored for as long without risk of deterioration. With respect to the lipid content of *C. manni* seeds, studies indicated a relatively high level (47.46 %) comparable to those reported by Ogunbusola *et al.* [17] ( $45.06 \pm 1.25 - 48.57 \pm 0.76$  %). Thus, *C. mannii* seeds could be classified as oilseeds that can be used as a source of vegetable oil for industrial and domestic purposes. The ash content ( $3.12 \pm 0.38$  %) was lower for the values reported for *Benincasa hispida* seeds (5.02- 11.81 %) [26] but similar to *M. oleifera* seeds (3.5 %) [18]. Therefore, *Cucumeropsis mannii* seeds could

**Table 1. Physicochemical characterization of *Cucumeropsis mannii* seeds flour**

Parameter	values
Moisture (%)	5.62 ± 0.60
pH	6.50 ± 0.00
Ashes (%)	3.12 ± 0.38
Protein (%)	31.80 ± 1.16
Lipids (%)	47.46 ± 1.55
Crude Fiber (%)	2.72 ± 0.3
Carbohydrates (%)	9.28 ± 0.79
Total sugars (%)	2.34 ± 0.05
Energie value (Kcal/100g DM)	591.46 ± 1.17

Values represent the mean ± standard deviation of three measurements (n = 3)

provide minerals essential to the proper functioning and development of the body. Concerning available carbohydrates content (9.28 ± 0.79 %), the rates investigated in this study were higher than those of *C. manni* (5.7 %) [18]) of the Congo and *C. ecirrhosus* seeds (18.69 ± 0.82 %) from Nigeria [27]. The energy value of *Cucumeropsis mannii* powder (591.46 ± 1.17 Kcal/ 100 g DM) was comparable with *C. ecirrhosus* seeds (601.7 Kcal/100g) [27]; almost lower to that of peanut paste (626 Kcal/100g) [28]; but higher with those reported for *Cucurbita maxima* (438.84 ± 0.07 Kcal/100g DM) [24] in Nigeria. The low level of total sugars (2.34 ± 0.05 %) observed was similar to that of *Citrullus lanatus* (2.23 %) from N'guetta et al., [29]. This suggests that *Cucumeropsis mannii* powder is a poor source of total sugars.

### 3.2 Phytochemical Compound

Table 2 shows the result for the anti-nutrient composition of *Cucumeropsis mannii* flour seeds. The total polyphenols in *Cucumeropsis mannii* powder (8.12 mg/g) in this study were lower than those of *Cucumeropsis mannii* (0.39 %) [30] in Cameroon. High polyphenols could increase the chelation capacity of these phenols and make nutrients and minerals in particular unavailable. Tannins have antioxidant, anti-mutagenic and anti-cancer properties [31].

The tannin content of *Cucumeropsis mannii* powder (3.02 g/100g) was higher than those reported by Ogunbusola et al., [17] with *Cucumeropsis mannii* (15.4 mg/100g) and *Cucumeropsis edulis* (4.69-29.11 mg EC/100g DM) obtained by Yété et al., [32]. However, they remain well below the recommended limit values of 1 mg/100 g DM, which influence protein bioavailability [33]. According to Kataria et al., [34], heat treatments of cooking reduce the level of tannins and denature insoluble tannin-nutrient

complexes. A research by Gülçin et al., [35] opined that tannins provided neuroprotective, cardioprotective and even antitumoral activities. Levels of flavonoids in this study were also lower than *Cucurbita maxima* (46.05 %) and *Citrullus lanatus* (41.57 %) seeds [36]. This concentration of flavonoid powder from *Cucumeropsis mannii* could protect the body from oxidative stress-related diseases, coronary artery disease and cancer [37]. The carotenoid (2.9 µg/100g DM) content was lower than the value recorded by Okorie [38] in white melon seed flour (Nigeria). It implies that the flour of *Cucumeropsis mannii* can reduce the risk of cardiovascular disease [39].

### 3.3 Amino Acid Profiles

The amino acid content for *Cucumeropsis mannii* seeds flour (table 3) indicated that histidine, leucine, isoleucine, methionine; phenylalanine, threonine, valine, aspartate, alanine, glycine, acid glutamate, tyrosine, serine and arginine were presents. In addition, the amino acids profile of *Cucumeropsis mannii* seeds flour showed that it contained both essential and non-essential amino acids; which would confirm that these proteins were of high quality. However, lysine has not been identified. Results have shown that aspartic acid (968.1mg/100g) was the most abundant of non- essential amino acids in melon seeds flour; while histidine (905.6 mg/100g) was the most abundant essential amino acids. the findings were similar to those of some authors who suggest that aspartic acid and glutamic acid were the most abundant amino acids in legume and nuts [40]. Essential amino acids were known for their significant physiological role in childhood growth [41]. Indeed, the proportion of arginine and histidine and over amino acids indicated that the seeds of *Cucumeropsis mannii* may be useful in formulating weaning food [42]. Some amino acids, such as phenylalanine, leucine, aspartate, serine, arginine and tyrosine, have been said to

**Table 2. Anti-nutrients composition of *Cucumeropsis mannii* seeds flour**

Parameter	Values
Flavonoïdes totaux (mg QE/g DM)	27.51 ± 0.03
Polyphénols totaux (mg GAE/g DM)	8.12 ± 0.09
Tanins (g TAE/100g DM)	3.02 ± 0.80
Oxalates (g/100g DM)	0.32 ± 0.02
Caroténoïdes (µg/100g DM)	2.9 ± 0.11

Values represent the mean ± standard deviation of three measurements (n = 3); DM: dry matter; Quercetin equivalent (QE); Gallic acid equivalent (GAE); tannic acids equivalent (TAE)

**Table 3. Amino acids profile of *Cucumeropsis mannii* seeds flour**

Amino acid	Values (mg/100g)
Histidine	905.6 ± 0.03
Isoleucine	844.5 ± 0.11
Leucine	456.6 ± 0.01
Méthionine	358.5 ± 0.04
Phénylalanine	550.0 ± 0.10
Thréonine	730.1 ± 0.02
Valine	469.9 ± 0.05
Aspartic acid	968.1 ± 0.06
Alanine	840.3 ± 0.32
Glycine	787.3 ± 0.22
Glutamic acid	771.3 ± 1.12
Tyrosine	769.4 ± 0.14
Serine	617.7 ± 0.27
Arginine	254.2 ± 0.33
Total essential amino Acids (ΣEAA)	616.45 ± 0.05
Total non Essential amino Acids ( ΣNEAA)	715.47 ± 0.32

Values represent the mean ± standard deviation of three measurements (n = 3)

have anti-sickling properties [43]. Researchers further explained that the anti-sickling effect of certain amino acid such as phenylalanine may be as a result of its anti-oxidative effects and also its ability to remove free radicals [44]. Results from this study would confirm that *Cucumeropsis mannii* proteins includes a significant proportion of essential amino acids. They could address children's and adult's protein needs in both developed and developing countries.

#### 4. CONCLUSION

The *Cucumeropsis mannii* seeds investigated in this study contained a high percentage of nutrients such as proteins and lipids, making them a good source of energy for human nutrition. The result of this finding also showed the presence of appreciable amounts of tannins, flavonoids and polyphenols. In addition, the presence of affordable quantities of tannins and carotenoids would represent a major nutritional asset because of their antioxidant power. However, *Cucumeropsis manni* seeds also contain a valuable source of essential amino acids, which could provide important health

benefits. This could make them valuable for animal and human diets.

#### ACKNOWLEDGEMENTS

The authors would like to acknowledge the staff of Nutrition and food safety Laboratory of Nangui Abrogoua University and Nangui Abrogoua's Biocatalysis and Bioprocessing Laboratory for their collaboration and providing the platforms under which this study where conducted.

#### CONSENT

Every participant in the study gave their informed consent.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Williams JT and Haq N. Global research on underutilized crops. An assessment of

- current activities and proposals for enhanced cooperation. ICUC, Southampton, UK. 2002;54.
2. Badifu GIO. Effect of processing on proximate composition, antinutritional and toxic contents of kernels from cucurbitaceae species grown in Nigeria. *Journal of Food Composition and Analysis*. 2001;14:153-161.
  3. Grodji Albarin Gbogouri, Kouakou Brou, Michel Linder, Elmira Arab TEHRANY, Dago Gnakri & Irié Arsène Zoro BI. Comparative study of physicochemical and thermal properties of the seed oils of three cucurbit species. *International Journal of Biological and Chemical Sciences*. 2011;5(3):1165-1177
  4. Zoro Bi I, Koffi KK, Djè Y, Malice M., & Badoin JP. Indigenous cucurbits of Côte d'Ivoire: a review of their genetic resources. *Science & Nature*. 2006;3(1):1-9.
  5. AOAC. Official methods of analysis of association of official analytical chemists. 17th ed. Gaithersburg, Maryland, USA; 2000.
  6. AOAC (Association of Official Analytical Chemists)., Official methods of analysis association of official analytical chemists. K. Helrich (ed). Fifteenth edition. Virginia (USA). 1990:963-964.
  7. Martínez-Herrera J, Siddhuraju P, Francis G, Dávila-Ortiz G and Becker K. Chemical composition, toxic/antimetabolic constituents, and effects of different treatments on their levels, in four provenances of *Jatropha curcas*L. from Mexico. *Food Chemistry*. 2006;96:80-89.
  8. Dubois MN, Gruillies KA, Hamilton JK, Rogers PA and Smith F. Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*. 1956;28:350-356.
  9. FAO. Food energy-methods of analysis and conversion factors. *Food and nutrition*.2002;77:93.
  10. Atwater W and Rosa E. A new respiratory calorimeter and the conservation of energy in human body, II. *Physical Review*. 1899;9:214-251.
  11. Bainbridge Z, Tomlings K, Wellings K and Westby A. Methods for assessing quality characteristics of non-grain starch staples (Part 3, laboratory methods) Chatham, United Kingdom Natural Resources Institute. 1996;34-39.
  12. FAO/IAEA. Quantification of tannins in tree foliage. A Laboratory Manual; 2000. Available:<http://www.iaea.org/programmes/nafa/d3/index.html>
  13. Makkar HPS, Blümmel M, Borowy NK and Becker K. Gravimetric determination of tannins and their correlations with chemical and protein precipitation methods. *Journal. Science. Food Agriculture.*, 1993;61:161–165.
  14. Cicco N, Lanorte TM, Paraggio M, Viggiano M and Lattanzio V. A reproducible rapid and inexpensive Folin-Ciocalteu micro-method in determining phenolics of plant methanol extracts. *Microchemical Journal*. 2009 ;91(1) :107-110.
  15. Meda A, Lamien CE, Romito M, Millogo J and Nacoulma OG. Determination of the total phenolic, flavonoïd and proline contents in Burkina Faso honeys as well as their radical scavenging activity. *Food chemistry*. 2005;91:571-577.
  16. Olaofe O, Adeyemi FO, Adediran GO. Amino acid and mineral composition and functional properties of some oils seeds. *Journal. Agriculture. Food Chemistry*. 1994;42:878-884.
  17. Ogumbusola EM, Fagbemi NT and Osundahunsi OF. Chemical and functional properties of full fat and defatted white melon (*Cucumeropsis mannii*) seed flours. *Journal of Food Science and Engineering*. 2012;26:691-696.
  18. Ntalani H, Ngouanou RD, Makomo H, Elouma NAM, Boungou-TG, Loumouamou AN, Mouloungui Z and Ouamba J-M. Comparative studies of the chemical composition and coagulant activity of the seeds of *Cucumeropsis mannii* Naud., *Arachis hypogaea* L. and *Moringa oleifera* Lam. in the clarification of surface waters. *Journal of Applied Biosciences*. 2020;145:14974 – 14984
  19. Ponka R, Nankap ELT, Tambe ST and Fokou E. Nutritional composition of some artisanal infant flours from Cameroon. *International Journal of Innovation and Applied Studies*. 2016;16(2):280 –292.
  20. Knoden J, Dufour LC and Bindelle J. Manufacture of peanut butter. Draft. Technical Manuals.Collection. P3-4 ; 2011.
  21. Sanchez-Castillo CP, Dewey PJS, Lara JJ, Anderson DL, Solano ML & James WPT. The starch and sugar content of some mexican cereals products, pulses, snack

- food fruits and vegetables. J.Food Comp. Anal., 2000;13:157-170.
22. Saada AD, Chabane AO, Boudroua K and Selselet AG. Role of pulses in human nutrition – nutritional strengths and health impacts. Regional Forum for the Development of Food Legumes – University of Mostaganem. 2016;5p.
  23. Rémond D, Walrand S. Legume seeds: nutritional characteristics and health effects. Agronomic Innovations, INRA. 2017;60:133-144.
  24. Mohaammed SS, Paiko YB, Mann A, Ndamitso MM, Mathew JT, Maaji S. Proximate, Mineral and Anti-nutritional. Composition of *Cucurbita Maxima* Fruits Parts, Nigerian Journal of Chemical Research. 2014;19: 37-49.
  25. Aryee FNA, Oduro I, Ellis WO, & Afuakwa JJ. The physicochemical properties of flour samples from the roots of 31 varieties of cassava. Food control. 2006;17:916- 922.
  26. Anwar F, Mohammad NA, Othman F and Saarii N. Inter-varietal variation in the composition of seeds and seed oils from winter melon (*Benincasa hispida* (Thumb) Cogn.) fruit. Pakistan Journal of Botany. 2011;43:2029-2037.
  27. Umar KJ, Hassan LG, Usman H, Wasagu RSU. Nutritional composition of the seeds of Wild melon (*Citrullus ecirrhosus*). Pakistan journal of Biological science. 2013 ;16(11):536-540
  28. Ciqal. Food Nutrition Composition Table. Retrieved on 16-08-2022; 2017. Available:<https://www.ANSES.fr>
  29. N'guetta AMN. Agro-morphological and nutritional characteristics of the cultivars of *Citrullus lanatus* (Cucurbitaceae) grown in Côte d'Ivoire and use of their meal in broiler production (*Gallus gallus*). Thesis for the title of Doctor in Biochemistry and Food Technology of Nangui Abrogoua University (Côte d'Ivoire). 2016;260p.
  30. Achu MB, Fokou E, Kamcsi G and Fotso M. Chemical evaluation of protein quality and phenolic compound levels of some cucurbitaceae oil seeds from Cameroun. African journal of biotechnology. 2013;12(7):735-743.
  31. Maisetta G, Batoni G, Caboni P, Esin S, Rinaldi AC and Zucca P. Tannin profile, antioxidant properties, and antimicrobial activity of extracts from two Mediterranean species of parasitic plant *Cytinus*. BMC complementary and alternative medicine. 2019;19 (82):1-11
  32. Yété P, Togbé A, Yaya K, Agbangnan P, Vital N, Djènontin TS, D Wotto D, Azandégbé EC and Sohounhloue D. Comparative study of phenolic compounds and radical-scavenging activity of the extracts of seeds of *Garcinia kola* (Guttiféreae) and *Cucumeropsis edulis* (cucurbitacéae) of Benin. International Journal of Innovation and Scientific Research, 2015 ;15(1):217-227.
  33. Li H. Wang Z, Liu Y. Reviews in the studies on tannins activity of cancer prevention and anticancer. Zhong-Yao-Cai, 2003 ;26(6):444 448.
  34. Kataria A., Chauhan B.M., & Punia D. Antinutrients and protein digestibility (in vitro) of mungbean as affected by domestic processing and cooking, Food Chemistry. 1989;32:9-17 .
  35. Gülçin L, Huyut Z, Elmasta M, Aboul-Enein HY. Radical scavenging and antioxidant activity of tannic acid. Arab J Chem. 2010;3:43-53.
  36. Karaye UI, Alero AA, Muhammad S, Bilbis LS. Evaluation of nutrient and antinutrient contents of selected Nigerian cucurbits seeds. Research journal of pharmaceutical, Biological and Chemical Sciences. 2013;4(1):137-142.
  37. Knekt P, Kumpulainen J, Järvinen R, Rissanen H, Heliövaara M, Reunanen A, Hakulinen T, Aromaa A. Flavonoid Intake and Risk of Chronic Diseases. Am. J. Clin. Nutr., 2002 ;76 :560–568.
  38. Okorie Peter Anyigor. Determining the physiochemical and phytochemical properties of local nigerian white melon seed flour. International journal of research Granthaalayah, 2018;6(5):157-166
  39. Schunemann HJ, McCann S, Grant BJB, Trevisan M, Muti P and Freudenheim JL. Lung function in relation to intake of carotenoid and other antioxidant vitamins in a populationbased study. American Journal of Epidemiology. 2002;155(5):463–471
  40. Olagunju AI and Ifesan BOT. Changes in nutrient and antinutritional contents of sesame seeds during fermentation. Journal of Microbiology, Biotechnology and Food Sciences. 2012;2(6):2407-2410.
  41. Parikh P, Semba R, Manary M, Swaminathan S, Udomkesmalee E, Bos R, Poh KB, Rojroongwasinkul N, Geurts J, Sekartini R and Nga TT. Animal source foods, rich in essential amino acids, are

- important for linear growth and development of young children in low- and middle-income countries. Maternal and child Nutrition. 2021;18: 1-12.
42. Olagunju AI and Ifesan BOT. Changes in nutrient and antinutritional contents of sesame seeds during fermentation. journal of Microbiology, Biotechnology and food sciences, 2013;2(6):2407-2410.
43. Nwaoguikpe RN and Ejele EA. Amino acid profile of some antisickling plant extracts and their haemoglobin polymerization inhibition. Nig J Biochem Mol Biol. 2010;25(2):53-59.
44. Dean J and Schechter AN. Sicklec cell anemia: molecular and cellular basis of therapeutic approaches. N Engl J Med. 1978;229:753-755.

---

© 2023 Agnei at al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/109439>