



A Practice to Enhance Soil Physico-Chemical Properties and Viable Microbial Count as Effected by Organic Nutrient Sources in Iris under Mid Hill Zone of Himachal Pradesh, India

Divesh Thakur ^{a++*}, Puja Sharma ^{a#}, Shreedhar Beese ^{a++},
Neha Grace Angel Kisku ^{a++}, Reva Jaryal ^{b++},
Shabnam Pangtu ^{c†}, Nidhi Shrama ^{a++} and Aman Guleria ^{a‡}

^a Department of Floriculture and Landscape Architecture, Dr. Y.S. Parmar, UHF Nauni, Solan, Himachal Pradesh-173230, India.

^b Department of Seed Science and Technology, Dr. Y.S. Parmar, UHF Nauni, Solan, Himachal Pradesh-173230, India.

^c Department of Agriculture Science, DAV University, Jalandhar, Punjab- 144012, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i14353

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110794>

Original Research Article

Received: 17/10/2023
Accepted: 22/12/2023
Published: 20/01/2024

⁺⁺ PhD Research Scholar;

[#] Associate Professor;

[†] Assistant Professor;

[‡] MSc;

*Corresponding author: E-mail: thakurdivesh125@gmail.com;

ABSTRACT

The current study was conducted at Experimental Farm of Department of Floriculture and Landscape Architecture, Dr. YS Parmar UHF, Nauni, Solan, H.P. during the year 2019-2020 to determine the ideal Jeevamrit dosage for enhancing the physico-chemical and microbiological characteristics of soil for iris at intervals of 15 days, sixteen applications of jeevamrit were applied as a foliar spray and drench. Three replications of the field experiment were set up using Randomized Block Design (RBD). The study showed that treatment T₁₆, which involved both 7.5% Jeevamrit drenching and 15.0% Jeevamrit foliar application, had the most positive impact on soil quality. This treatment resulted in the highest organic carbon level (1.48), nitrogen content (362.59 kg/ha), phosphorus content (137.48 kg/ha), potassium content (267.85 kg/ha), and a high count of viable bacteria (75.63×10^6 cfu/g soil), viable fungi count (28.57×10^4 cfu/g soil), and viable actinomycetes count (45.90×10^3 cfu/g soil). Additionally, this treatment had the lowest pH and EC levels.

Keywords: Jeevamrit; microbial count; organic farming; soil properties.

1. INTRODUCTION

The 'Green Revolution' was a significant shift in agricultural practices that took place between the 1940s and 1960s. During this period, the application of chemical fertilizers, pesticides, herbicides, and fungicides became the industry standard worldwide. However, the excessive use of these techniques resulted in soil degradation, poor quality production, environmental pollution, and severe health hazards. Inorganic fertilizers can only provide one or two nutrient elements, which is why organic farming is considered an alternative practice to enhance microbial biomass and soil structure. (Naeem et al., [1]. Suresh et al., [2]. Agriculture is the primary source of livelihood for small and marginal farmers in states like Himachal Pradesh. Due to the topographic challenges, these farmers already practice traditional agriculture methods with minimal use of chemical inputs. However, with the steep increase in the cost of inorganic fertilizers, small and marginal farmers find it hard to afford them. To make technology accessible to economically poor farmers, we can adopt the practice of zero-budget natural farming (ZBNF), which is the process of farming with nature and without the use of chemicals. ZBNF replaces inorganic fertilizers with organic manure, making it a sustainable and cost-effective option for farmers. Naeem et al., [1]. Jeevamrit is a word that comes from two Hindi words: 'Jeevan' which means life and 'Amrit' which means medicinal potion. Earthworm activity and microbial population both significantly increase when Jeevamrit is applied to soil. This, in turn, increases nutrient availability in the soil and fortifies its resistance and resilience mechanisms. As a result, crop productivity is significantly increased

Vashishat et al., [3]. Lazarovits et al., [4] Sreenivasa et al., [5]. The soil ecosystem's resistance and stability are increased by the elevated microbial population diversity index. Palekar et al., [6] Devakumar et al., [7] Pathak et al., [8]. The rich microbial community of soil microorganisms plays a crucial role in transforming organic matter in soil and acting as a source and sink of nutrients that plants can use. This helps to improve soil fertility and crop yield [9]. Applying Jeevamrit to the soil is a great way to enhance the growth of beneficial microorganisms. These microorganisms convert nutrients from an unavailable form to an easily absorbable form, so that plants can absorb them more easily. You can use Jeevamrit by either spraying or sprinkling it on the crop field. It is essential to adopt sustainable agricultural practices based on scientific facts for growing different crops. In this context, a study was carried out in the mid-hill region of Himachal Pradesh to assess the effects of organic nutrient sources on the physico-chemical properties of the soil and the viable microbial count.

2. MATERIALS AND METHODS

The purpose of this study was to investigate the impact of organic nutrient sources on Iris (*Iris orientalis* Mill.) cv. Frigia, soil physical and chemical properties, and viable microbial count. The research was conducted from December 2019 to May 2020 at the Experimental Farm of Department of Floriculture and Landscape Architecture, Dr. YS Parmar UHF, Nauni, Solan, H.P. This experimental site is situated at a latitude of 30°51'0" North and a longitude of 77°11'30" East, at an elevation of 1276 m above mean sea level. The experimental site is in the

Himachal Pradesh mid-hill region, which is classified as an agro-climatic zone with moderate rainfall between 1000 and 1300 mm.

The experiment was set up on December 30, 2019, in a Randomized Block Design (RBD) using combination of sixteen treatments with three replications. The treatment combinations were assigned to different plots randomly, and Table 1 presents the details of the treatments.

For a networking area of 1.00 square meters, beds measuring 1.20 by 1.20 meters were made for the planting of rhizomatous plants. The rhizomatous plants are spaced 20 cm x 30 cm apart. The treatment combinations will be applied at 15-day intervals alternatively in treatments T₈ to T₁₆, as shown in Table 1.

2.1 Preparation of Jeevamrit

Jeevamrit was prepared by dissolving the suggested ingredients Sreenivasa et al, [10].

All the ingredients were mixed in a plastic drum, wrapped with a moist jute sack, and then kept in the shade before being used according to the treatment schedule.

3. RESULTS AND DISCUSSION

The viable microbial count along with various soil physico-chemical property data are shown in Table 3 and Table 4 respectively.

3.1 Organic Carbon (%)

A noteworthy maximum of 1.48 percent of organic carbon was observed when 7.5% of Jeevamrit was drenched and 15.0% of Jeevamrit foliar was applied (T₁₆). Minimum organic carbon (1.02 %) was noticed with treatment T₁ (Control). The application of liquid manure may have increased the incorporation of organic materials and stimulated root growth, which has resulted in an increase in soil organic carbon. Increased soil organic carbon content could have been the consequence of these compounds' breakdown [11] Applying Jeevamrit may have caused a drop in bulk density, which could be linked to higher levels of organic carbon content Chandrakala, [12] Jenny and Malliga, [13]. The findings of Singh et al. [14] and Rai et al. [15] corroborate the results.

3.2 pH and EC

The application of 7.5% Jeevamrit drenching + 15.0% Jeevamrit foliar (T₁₆) application resulted in the lowest soil pH (6.58). Soil EC was non significantly affected by the application of Jeevamrit. It is preferable to have lower soil pH and Electrical ConductivityOne possible explanation for the drop in soil pH and EC levels is the organic acids released during the organic manure's breakdown. This might be because applying these manures has increased the permeability of the soil. Rai et al. [16] have similarly noted a reduction in soil EC and pH when Jeevamrit is applied.

Table 1. Detail of treatments

Treatments	Treatment details
T ₁	: Control (Without Jeevamrit)
T ₂	: 2.5% Jeevamrit drenching
T ₃	: 5.0% Jeevamrit drenching
T ₄	: 7.5% Jeevamrit drenching
T ₅	: 5.0% Jeevamrit foliar application
T ₆	: 10.0% Jeevamrit foliar application
T ₇	: 15.0% Jeevamrit foliar application
T ₈	: 2.5% Jeevamrit drenching + 5.0% Jeevamrit foliar application
T ₉	: 2.5% Jeevamrit drenching + 10.0% Jeevamrit foliar application
T ₁₀	: 2.5% Jeevamrit drenching + 15.0% Jeevamrit foliar application
T ₁₁	: 5.0% Jeevamrit drenching + 5.0% Jeevamrit foliar application
T ₁₂	: 5.0% Jeevamrit drenching + 10.0% Jeevamrit foliar application
T ₁₃	: 5.0% Jeevamrit drenching + 15.0% Jeevamrit foliar application
T ₁₄	: 7.5% Jeevamrit drenching + 5.0% Jeevamrit foliar application
T ₁₅	: 7.5% Jeevamrit drenching + 10.0% Jeevamrit foliar application
T ₁₆	: 7.5% Jeevamrit drenching + 15.0% Jeevamrit foliar application

Table 2. Composition of jeevamrit

Ingredients	Quantity
Cow dung	10 kg
Cow urine	10 litres
Jaggery	2 kg
Pulse flour	1 kg
Live soil	A handful of soil
Water	200 litres

*Dilution of 5 litres of Jeevamrit in 100 litres of water for 5% foliar spray/drench.

Table 3. Effect of Jeevamrit on the soil physico-chemical properties in Iris

Treatments	Organic carbon (%)	pH	Electrical conductivity (dS/m)	N (kg/ha)	P (kg/ha)	K (kg/ha)
T ₁	1.02	6.82	0.41	240.89	93.33	204.19
T ₂	1.15	6.70	0.39	284.55	112.51	223.86
T ₃	1.18	6.65	0.32	301.91	123.29	242.03
T ₄	1.20	6.62	0.28	345.07	129.83	259.16
T ₅	1.06	6.77	0.40	258.67	97.96	211.53
T ₆	1.09	6.74	0.39	268.12	100.97	219.03
T ₇	1.12	6.70	0.38	276.73	104.62	228.26
T ₈	1.19	6.72	0.39	289.01	113.50	226.92
T ₉	1.21	6.71	0.39	294.69	114.40	229.15
T ₁₀	1.24	6.71	0.37	298.16	115.40	231.82
T ₁₁	1.27	6.68	0.34	312.26	125.10	243.82
T ₁₂	1.31	6.66	0.32	320.90	126.43	245.86
T ₁₃	1.35	6.64	0.30	326.08	128.22	248.76
T ₁₄	1.41	6.62	0.95	352.55	132.45	262.26
T ₁₅	1.45	6.60	0.27	358.33	134.61	264.89
T ₁₆	1.48	6.58	0.25	362.59	137.48	267.85
C.D. (0.05)	0.009	0.008	NA	1.837	0.568	1.845

Table 4. Effect of Jeevamrit on the soil's microbial count (cfu) in Iris

Treatments	Viable bacterial count ($\times 10^6$ cfu/g soil)	Viable fungi count ($\times 10^4$ cfu/g soil)	Viable actinomycete count ($\times 10^3$ cfu/g soil)
T ₁	32.40	9.57	30.67
T ₂	41.57	15.33	38.07
T ₃	51.37	21.33	42.70
T ₄	60.83	25.00	44.67
T ₅	30.69	9.07	30.70
T ₆	32.93	9.77	31.20
T ₇	34.83	9.77	32.30
T ₈	58.93	18.93	38.57
T ₉	59.03	19.17	38.70
T ₁₀	59.50	19.23	39.17
T ₁₁	67.33	24.17	43.37
T ₁₂	67.67	24.23	43.47
T ₁₃	68.87	24.93	43.50
T ₁₄	74.07	28.23	45.43
T ₁₅	73.30	28.03	45.33
T ₁₆	75.63	28.57	45.90
C.D. (0.05)	2.07	1.08	0.98

3.3 Available N (kg/ha)

It is clear from the information in Table 3 that the application of Jeevamrit influenced the available nitrogen. Maximum nitrogen (362.59 kg/ha) was found in T₁₆ with application of 7.5 % Jeevamrit drenching + 15.0 % Jeevamrit foliar. However, minimum available nitrogen (240.89 kg/ha) was registered with treatment T₁ (Control). The nitrogen content in soil becomes more available when supplemented with organic manures, which increases basal respiration by promoting the growth of microorganisms [17]. The gradual release of nutrients through FYM may have contributed to the rise in the pool of accessible nitrogen. Applying jeevamrit increased the rate at which soil microorganisms multiplied and nitrogen-fixing bacteria worked synergistically to convert organic nitrogen to inorganic form. The outcomes align with the findings of Rai et al. [16].

3.4 Available P (kg/ha)

Significantly maximum phosphorus (137.48 kg/ha) was recorded with the application of 7.5 % Jeevamrit drenching + 15.0 % Jeevamrit foliar application (T₁₆). Minimum available phosphorus (93.33 kg/ha) was recorded with treatment T₁ (Control). Due to soil microbial activity resulting in the release of organic acids during the decomposition of organic matter, phosphates were eventually solubilized, increasing the phosphorus content of the soil. This could be the reason for the increase in phosphorus availability in the soil as a result of the addition of these manures. Nitika et al. [18] reported similar results.

3.5 Available K (kg/ha)

Maximum potassium (267.85 kg/ha) was recorded with the application of 7.5 % Jeevamrit drenching + 15.0 % Jeevamrit foliar application (T₁₆). Whereas minimum available potassium (204.19) was noticed with treatment T₁ (Control). Potassium may have been added directly to the soil through organic manures, and organic matter may have interacted with clay to release K₂O ions that resulted in an increase in the amount of potassium that is available in the soil. Suklabaidya et al. [19] and Prativa and Bhattarai [20] reported identical results.

3.6 Viable Microbial Count

Treatment T₁₆ (5% Jeevamrit foliar application + 7.5% Jeevamrit drenching) produced the highest

viable bacterial count (75.63 × 10⁶ cfu/g soil), viable fungal count (28.57 × 10⁴ cfu/g soil), and viable actinomycetes count (45.90 × 10³ cfu/g soil). The microbial population may have increased as a result of organic inputs (Jeevamrit), which release carbon dioxide (CO₂) during the soil's breakdown process, giving bacteria food and a microenvironment in which to develop and multiply (Subba Rao, 2018). Similar results were recorded by Pathak and Ram [8] Devakumar et al. [21] and Vemaraju (2014) [22,23].

4. CONCLUSION

After conducting the experiment, it was determined that applying a mixture of 7.5% Jeevamrit drenching and 15.0% Jeevamrit foliar application (T₁₆) at 15-day intervals significantly improved the NPK content, organic carbon, and viable microbial count of Iris plants in the mid-hill zone of Himachal Pradesh. Moreover, the application also resulted in reduced pH and electrical conductivity levels.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Naeem M, Iqbal J, Bakhsh MAA. Comparative study of organic fertilizers and organic manures on yield components of Mungbean. *Journal of Agriculture and Social Sciences*. 2006;2(4):227-239.
2. Suresh KD, Sneh G, Krishn KK, Mool CM. Microbial carbon and microbial activities of soils receiving chemical fertilizers and organic amendments. *Archives Agronomy and Soil Science*. 2004;5(6):641-647.
3. Vashishat RK, Sharma S, Laishram C. Problems and factors affecting adoption of natural farming in Sirmaur district of Himachal Pradesh, *Indian Journal of Ecology*. 2021;48(3):944-949.
4. Lazarovits G, Nowak J. Rhizobacteria for improvement of plant growth and establishment. *Hortscience*. 1997;32(2):188-192.
5. Sreenivasa MN, Nagaraj MN, Bhat SN. Beejamrutha, A source for beneficial bacteria. *Karnataka Journal of Agricultural Sciences*. 2009;22(5):1038-1040.

6. Palekar S. Shoonya bando valadanai sargika krushi, in: Swamy Anand, Agri Prakashana, Bangalore, India; 2006.
7. Devakumar N, Rao GGE, Shubha S, Khan I, Nagaraj S, Gowda SB. Organic Farming and Activities of Organic Farming Research Centre. A Bulletin by University of Agricultural Sciences, Bangalore. 2008;18-27.
8. Pathak RK, Ram RA. Role of the cow for the evergreen revolution through the integrated organic farming system. In: Proceedings of National Conference on Glory of Gomatha, Dec. 1-3, Sri Venkateswara Veterinary University, Tirupati, Andhra Pradesh, India. 2007;170-177.
9. Gupta AP, Antil SR, Narwal PR. Effect of farm yard manure on organic carbon, available N and P contents of soil during different periods of wheat growth. Journal of the Indian Society of Soil Science. 1988;36:269-273.
10. Sreenivasa MN, Naik N, Bhat SN. Nutrient status and microbial load of different organic liquid manures. Karnataka Journal of Agricultural Sciences. 2011;24(4):583-584.
11. Kumari R, Thakur KS, Lalhruaitluangi N. Effect of different organic nutrient sources on soil properties in onion (*Allium cepa* L.). International Journal of Current Microbiology and Applied Sciences. 2019;8(4):1783-1792.
12. Chandrakala M. Effect of FYM and Fermented Liquid Manures on Yield and Economics of Chilli (*Capsicum annum* L.). Research Journal of Agricultural Science. 2008;2(3):722-724.
13. Jenny S, Malliga P. Assessment of different concentrations of organic manure on the growth and yield of *Solanum lycopersicum* L. (Tomato). International Journal of Innovation Research Science Engineering and Technology. 2016;5(3):3722-3731.
14. Singh SN, Rai JP, Singh SR, Goyal SK, Singh SP. Effect of integrated use of organic manures and fertilizers on yield, nutrient uptake and soil fertility in onion on red soils of Vindhyan region. Vegetable Science. 2014;41:150-154.
15. Rai S, Rani P, Kumar M, Rai AK, Shahi SK. Effect of integrated use of vermicompost, FYM, PSB and Azotobacter on physico-chemical properties of soil under onion crop. Environment and Ecology. 2014;32(4B):1797-1803.
16. Rai S, Rani P, Kumar M, Rai AK, Shahi SK. Effect of integrated use of vermicompost, FYM, PSB and Azotobacter on physico-chemical properties of soil under onion crop. Environment and Ecology. 2014;32(4B):1797-1803.
17. Rathore G, Kaushal R, Sharma V, Sharma G, Chaudhary S, Dhaliwal SS, Alsuhaibani AM, Gaber A, Hossain A. Evaluation of the usefulness of fermented liquid organic formulations and manures for improving the soil fertility and productivity of brinjal (*Solanum melongena* L.). Agriculture. 2023;13(2):1-14.
18. Nitika, Thakur KS, Mehta DK. Soil nutrient status, NPK uptake and viable bacterial count as influenced by different organic nutrient sources in European carrot. International Journal of Current Microbiology and Applied Sciences. 2018;7(8):1783-1789.
19. Suklabaidya A, Datta M, Kandpal BK. Response of organic sources and biofertilizer in soil fertility and yield of cauliflower in the foot hills of Tripura. International Journal of Agriculture Sciences. 2017;9(15):4874-4875.
20. Pratava KC, Bhattarai BP. Effect of integrated nutrient management on the growth, yield and soil nutrient status in tomato. Nepal Journal of Science and Technology. 2012;12(2011):23-28.
21. Devakumar N, Shubha S, Gouder SB, Rao GGE. Microbial analytical studies of traditional organic preparations beejamrutha and jeevamrutha. In: Proceedings of Building Organic Bridges. 4th ISOFAR Scientific Conference, Istanbul, Turkey. 2014;639.
22. Devakumar N, Rao GGE, Shuba SE. Valuation of locally available media for the growth and development of nitrogen-fixing micro-organisms, 28 September - 01 October, in: Proceedings of the 3rd Scientific Conference of ISOFAR Organic Are Life, knowledge for tomorrow, Korea. 2001;504-509.

23. Palekar S. Text book on Shoonya Anand, Agri Prakashana, Bangalore, India.
Bandovalada Naisargika Krushi. Swamy 2006;154.

© 2024 Thakur et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/110794>