



Knowledge of Safe Period and Utilization of Long-Lasting Reversible Contraceptives among Female Adolescents in an Urban City in Nigeria

Abade Ese ^a, Frank Abade ^a, Elechi Comfort ^a
and Blessing UKoro ^{b*}

^a Department of Human Kinetics, Health and Safety Education, Rumuolumeni, Port Harcourt, Rivers State, Nigeria.

^b Department of Human Physiology, University of Port Harcourt, Port Harcourt, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111557>

Original Research Article

Received: 12/11/2023
Accepted: 17/01/2024
Published: 22/01/2024

ABSTRACT

The study sought to evaluate the understanding of safe period and long-lasting reversible contraceptives among adolescent females in an urban city in Nigeria, employing a descriptive research approach. The study was directed by three research questions and one hypothesis. The study utilized a sample size of 900 teenagers residing in Port Harcourt city. The researchers utilized a methodical self-administered questionnaire to collect data from the participants. The research questions and hypothesis were analyzed using percentages and binary logistic regression with a significance level of 0.05 (alpha level). The study revealed that 65.6% of the respondents demonstrated a good understanding of safe period, while 62.1% had a good knowledge of long-lasting contraceptive methods. Only 14.7% of the respondents reported frequent use of

*Corresponding author: Email: blessing_ukoro@uniport.edu.ng;

contraception, with the majority (25.6%) of frequent users being in SS2. The findings revealed a notable disparity in the average rate of contraceptive utilization among teenagers residing in an urban area of Nigeria, depending on their social class [$F(2,887) = 854.499$; $p < 0.05$]. The study determined that adolescents possessed a strong understanding of safe periods and long-acting reversible contraceptives. Furthermore, a notable correlation was observed between class and contraceptive usage. Policy makers are advised to ensure that reproductive health services are easily available to teenagers by enacting regulations that promote the development of family planning clinics specifically designed for young people.

Keywords: family planning; contraceptives; follicular phases; ovulation.

1. INTRODUCTION

The menstrual cycle is a regular monthly process including physiological changes in the ovaries and uterus of adult women. The ovarian cycle and uterine cycle are two subdivisions of the menstrual cycle, which is an essential process for sexual reproduction [1]. The three phases of the ovarian cycle are the luteal, ovulation, and follicular phases. Ovulation is the process of releasing an egg from a Graafian follicle. Ovulation, the most important event for the possibility of pregnancy, takes place approximately halfway through the menstrual cycle [2].

In order to ascertain the likelihood of conception, it is necessary to possess a fundamental comprehension of the physiology of the menstrual cycle, specifically the period of ovulation. It is imperative to endorse or utilise family planning methods (FPMs), which are widely employed by a significant number of women worldwide [1]. As per the United Nations (2013), it is imperative for nations to limit their fertility rates because population size affects the standard of living for citizens. As a result, women are using hormonal contraceptives, which are linked to a number of health problems such as obesity, hypertension, and cancer. Globally, the rate at which people stop using contemporary contraception is rising at a pace that ranges from 19% to 36% [3].

Knowledge of fertile period is one of the natural family planning strategies that uses scientific ways to delay pregnancy. Those who prefer not to use hormonal, mechanical, or surgical methods of contraception are the predominant users [4]. This technique consists of the cervical mucus (or Billings) method, the symptom thermal approach, and the basal body temperature method. The successful application of the fertile period can be facilitated by the use of these three strategies. Women must accurately determine

when their menstrual cycle is fertile in order to practice periodic abstinence [5]. However, the majority of women sometimes cannot tell for sure when in their menstrual cycle they have the highest chance of becoming pregnant.

A single strategy to reduce unwanted pregnancies is to be aware of the fertile period. According to a World Health Organization (WHO) research, two-thirds of women who wanted to postpone or reduce having children discontinued using modern contraception due to health concerns, fear of adverse effects, and underestimating the likelihood of becoming pregnant [6]. An accurate understanding of one's ovulatory cycle might prevent unwanted and unintended pregnancies. Similarly, individuals who do not employ contraception and lack knowledge of their fertile period are at a higher risk of experiencing unintended pregnancies [7]. Moreover, the majority of couples who were actually engaging in the practice of identifying their fertile period were uninformed about their fertility window and did not depend on the concept of fertility period due to insufficient understanding.

Knowledge of fertile period is one of the main causes of the decline in high fertility, which has detrimental effects on social and economic growth [8]. Women would be better able to plan their pregnancies, identify their pregnancies early, and comprehend the risk of pregnancy if they had access to sufficient information about fertile times and relevant physiology. Natural family planning has many benefits, but a study revealed that some couples find it difficult to maintain their relationship because of issues with abstinence, a decline in frequency and spontaneity, and unequal sexual desires between partners. Additionally, some of the couple's disagreements and feelings of resentment stemmed from their unfulfilled sexual desires, which made their relationship worse [9].

According to a study done in the US, almost 40% of women were unaware of the ovulatory period [10]. According to a comparable study done in the US, 51% of women were unaware of the fertile period [11]. About 40% of individuals in an Australian study were unsure of when menstrual cycle design was most likely to occur [12]. In a similar vein, a Turkish study on youth revealed that just 40% of youth had adequate awareness of the fertile period [13]. Furthermore, a study on a specific adult population in Pakistan found that just 46% of the participants were aware of the female cycle's fertile phase [14].

In twenty-nine (29) sub-Saharan African nations, a research employing data from the Demographic and Health Survey (DHS) among women of reproductive age found that only 8.3% of them knew what ovulation was [15]. This is a relatively low percentage. According to a study on male teenagers in Ghana who were asked to identify the female fertile period, only 14.2% of them could accurately name the particular period that falls inside the female menstrual cycle [16]. Comparably, a Kenyan study discovered that fewer than 8% of participants accurately identified their years of fertility [17].

Long-acting reversible contraceptives (LARCs) are implant and intrauterine device contraceptive techniques that are very effective and convenient, as well as long-lasting and requiring little to no maintenance. These methods have become more widely available recently, giving women the option to adopt their use. It is also more affordable and has far higher compliance rates than other hormonal approaches. Compared to user-dependent and shorter-term methods, which both raise the risk of non-compliance associated method failure (18, 19), long-term contraceptive methods (LARCs) are the best options for many women to avoid pregnancy.

When used as prescribed, Long-acting reversible contraceptives (LARCs) offer women ongoing pregnancy prevention for a minimum of three years and a maximum of ten years. Unlike short-acting techniques, these devices do not suffer from user errors, which accounts for their 99% effectiveness [18]. Moreover, LARC techniques can narrow the difference in failure rates between "typical use" and "perfect use". Globally, the failure of contraceptives results in about 48% of unwanted pregnancies, primarily as a result of improper use, inadequate adherence, and/or technological malfunction [19,20]. By using LARC approaches, this can be prevented as they do not rely on taking prescribed medications as directed,

remembering to replace patches or rings, or scheduling doctor's appointments [20; 21].

The aim of this study was to assess the knowledge of safe period and long lasting reversible contraceptives among adolescent female population in an urban city in Nigeria.

2. METHODOLOGY

The study used a descriptive survey research design with a study population of 70535 female adolescents in Port Harcourt, Nigeria (National Population Commission of Nigeria, 2011). Using the Taro Yamane formula for a finite population, a sample size of 900 was calculated. The study used a multi-stage sampling approach. To collect data from respondents, a validated self-administered structured questionnaire was used. The researcher administered the questionnaire with the assistance of two trained research assistants. The research assistants were instructed on how to complete the questionnaire and the study's objectives. The questionnaire is divided into two pieces. Section A collected information on the respondents' socio-demographics, while Section B gathered information on adolescents' awareness of safe period and long-lasting contraception in Port Harcourt. The validated instrument was pre-tested on a part of the population homogenous to the research sample in a different city and was proven to have a good internal consistency. Data was analyzed using descriptive statistics of percentage while regression analysis was used to test the hypotheses.

3. RESULTS

Research question 1: What is the knowledge of safe period among adolescents in an urban city in Nigeria?

Table 1 shows the knowledge of safe period among adolescents in an urban city in Nigeria. According to the findings, 584 (65.6%) of respondents had high understanding of the safe period, whereas 306 (34.4%) had poor knowledge.

Research question 2: What is the knowledge of long lasting contraceptives among adolescents in an urban city in Nigeria?

Table 2 shows the knowledge of long lasting contraceptive methods among adolescents in an urban city in Nigeria. The result shows that 553(62.1%) of the respondents had good

knowledge of long lasting contraceptive methods and 337(37.9%) had poor knowledge.

The finding of the study showed that 131(14.7%) of the respondents frequently used contraceptive

while 254(28.5%) sometimes used contraceptives. The result further showed that 23(7.4%) of respondents who were in SS1, 100(25.6%) in SS2 and 8(4.2%) in SS3 frequently used contraceptive.

Table 1. Level of knowledge of safe period

S/N	Items	True Freq %	False Freq %
Knowledge of safe period			
1	Appropriate calculations of ovulation/periods can prevent pregnancy	781(87.8)	109(12.2)
2	Appropriate calculations of ovulation/periods can prevent pregnancy	779(87.5)	111(12.5)
3	A woman will not get pregnant during her safe period	709(79.7)	181(20.3)
4	Ovulation does not take place during safe period	383(43.0)	507(57.0)
5	I know how to calculate ovulation period	176(19.8)	714(80.2)
6	Safe period is effective	677(76.1)	213(23.9)
Grand total		584(65.6)	306(34.4)

**Decision: >50% Good, <50% Poor*

Table 2. Level of knowledge of long lasting contraceptives methods

S/N	Items	True Freq %	False Freq %
The following are long lasting contraceptives			
1	Intrauterine device	750(84.3)	140(15.7)
2	Contraceptive implant	756(84.9)	134(15.1)
3	Contraceptive injection	692(77.8)	198(22.2)
4	Contraceptive ring	392(44.0)	498(56.0)
5	Diaphragm	174(19.6)	716(80.4)
Grand total		553(62.1)	337(37.9)

**Decision: >50% Good, <50% Poor*

Table 3. Class and contraceptive use among adolescents

Variables	Practice of contraceptive use			Total Freq (%)
	Frequent Freq (%)	Sometimes Freq (%)	Rarely Freq (%)	
Class				
SS1	23(7.4)	30(9.7)	256(82.8)	309(100)
SS2	100(25.6)	214(54.7)	77(19.7)	391(100)
SS3	8(4.2)	10(5.2)	172(90.5)	190(100)
Total	131(14.7)	254(28.5)	505(56.7)	890(100)

Table 4. Regression analysis showing significant difference in the mean value of contraceptive use among adolescents in an urban city in Nigeria based on class

Model		Sum of Squares	Df	Mean Square	F	Sig.	Decision
1	Regression	141.503	2	70.751	854.499	0.00*	Rejected
	Residual	73.442	887	.083			
	Total	214.945	889				

**Statistical significant (p<0.05)*

4. HYPOTHESIS

Ho: There is no significant difference in the mean value of contraceptive use among adolescents in an urban city in Nigeria based on class.

The finding of this study shows that there is a significant difference in the mean value of contraceptive use among adolescents in an urban city in Nigeria based on class [F(2,887) = 854.499; p<0.05]. Therefore, the null hypothesis which states that there is no significant difference in the mean value of contraceptive use among adolescents in an urban city in Nigeria based on class was rejected.

5. DISCUSSION OF FINDINGS

The findings of the study show that more than half of the respondents had good knowledge of safe period. This was in variance with the findings of a cross-sectional study carried out in 29 African nations, which showed that the average level of ovulation awareness in those nations was 21.5% [15]. The results were higher than those of a Michigan state survey [1], which found that approximately 32.8% of women knew when to expect their period. In a similar vein, the findings of this study exceeded those of a study carried out in Ghana [22], which revealed that around 38% of women knew that the ovulation window occurs halfway through the menstrual cycle. Getahun and Nigatu [23] showed only 3,699 (23.6%) were found to be knowledgeable about a woman's ovulation period. Factors associated with increased knowledge of the ovulation period among reproductive women included living in an urban area (AOR=1.93; 95% CI=1.76, 2.12), higher educational status (AOR=4.39; 95% CI=3.77, 5.11), receiving family planning counseling within the last 12 months (AOR=1.33; 95% CI=1.20, 1.48), and having menstruation within the previous six months (AOR=1.30; 95% CI=1.18, 1.42).

However, the findings of this investigation were consistent with those of research done in Australia and India [24; 25]. This discrepancy may result from variations in the study participants' socioeconomic background (educational attainment), study environments, and study population. The general consensus from numerous research reviews was that women know very little to nothing about fertility, the fertile window, or the ovulatory phase.

The findings of this study showed majority of the respondents had good knowledge of long lasting

contraceptive methods. This finding did not agree with that of Bolarinwa and Olagunju [26] which showed only 14.8% of women were using long lasting contraceptive methods, but showed a significant relationship between knowledge and use of long lasting contraceptives which implied there was low knowledge of long lasting contraceptive among the women and this calls for an awareness campaign on the benefits of LARC. However the study's finding agreed with that of White et al. [27] which showed overall 51%, 23% and 47% of women reported they had heard of the copper IUD, LNG-IUS and implant, respectively. The main justifications given by women who were considering LARC were that it was more convenient, effective, and offered longer-term contraception protection than their current method.

The findings of the study showed that majority of the respondents in SS2 indicated good contraceptive utilization, while the less than a quarter respondents from the other classes indicated good contraceptive utilization. This implies that there is influence of class on contraceptive usage. The findings of the present study is not in line with the study of [28] which indicated that majority of the students of the senior secondary class three had better knowledge and use of contraceptives compared to other classes. Also, the present study deviates from the findings of Gbagbo [29] which reported that awareness and use of contraceptives are more prevalent among junior high school pupils. The discrepancies in the findings could be attributed to the pupils' exposure to and access to sex education information. However, students who got contraceptive information from their parents/guardians were more likely to use contemporary contraception regularly than their peers who did not. As a result, with young people in primary schools becoming sexually active, there is a need for formalized contraceptive education in primary schools to provide right information and teaching.

6. CONCLUSION

Based on the findings of this study, there was good knowledge of safe periods and long lasting contraceptives among adolescents, Also there was a strong association between class and use of contraceptives.

7. RECOMMENDATIONS

The study's findings led to the following recommendations:

- Policy makers should make reproductive health services accessible to adolescents by promulgating laws that encourages the establishment of youth friendly family planning clinics.
- It is crucial for the governments, non-governmental organisations (NGOs), donor agencies, and key stakeholders to collaborate in order to guarantee the availability, accessibility, and continuous promotion of long-lasting contraceptives for teenagers.
- School heads should encourage peer-led initiatives that empower students with accurate knowledge about safe periods and long lasting contraceptives to their peers.
- Ministry of health should develop and maintain informative websites or mobile applications that provide reliable and up-to-date information on use of contraceptives, leveraging on social media to disseminate educative and relatable content.

ETHICAL APPROVAL AND CONSENT

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ayoola AB, Zandee GL, Adams YJ. Women's knowledge of ovulation, the menstrual cycle, and its associated reproductive changes. *Birth*. 2016; 43(3):255–262.
2. Ben-Nun L. Awareness of Ovulation BN. Publication House. Ashkelon, Israel; 2015.
3. Belete N, Zemene A, Hagos H, Yekoye A. Prevalence and factors associated with modern contraceptive discontinuation among reproductive age group women, a community based cross-sectional study in humera town, northern Ethiopia. *BMC Women's Health*. 2018; 18(1):1–8.
4. Manhart MD, Fehring RJ. The state of the science of natural family planning fifty years after humane vitae: A report from NFP Scientists Meeting Held at the US Conference of Catholic Bishops. *Linacre Q*. 2018;85:339–347. DOI: 10.1177/0024363918809699.
5. Sheon AR, Stanton C. Use of periodic abstinence and knowledge of the fertile period in 12 developing countries. *International Family Planning Perspectives*. 1989;15:29–34. DOI: 10.2307/2133276
6. WHO. High rates of unintended pregnancies linked to gaps in family planning services: New WHO study. World Health Organization; 2019 October 25. Available: <https://www.who.int/news-room/detail/25-10-2019-high-rates-of-unintended-pregnancies-linked-to-gaps-in-family-planning-services-new-who-study>
7. Boah M, Bordotsiah S, Kuurdong S. Predictors of unsafe induced abortion among women in Ghana. *Journal of Pregnancy*; 2019. DOI: 10.1155/2019/9253650
8. Alene GD, Worku A. Differentials of fertility in north and south Gondar zones, northwest Ethiopia: A comparative cross-sectional study. *BMC Public Health*. 2008;8:1–12. DOI: 10.1186/1471-2458-8-397
9. VandeVusse L, Hanson L, Fehring RJ, Newman A, Fox J. Couples' views of the effects of natural family planning on marital dynamics. *Journal of Nursing Scholarship*. 2003;35:171–176. DOI: 10.1111/j.1547-5069.2003.00171.x
10. Lundsberg LS, Pal L, Garipey AM, Xu X, Chu MC, Illuzzi JL. Knowledge, attitudes, and practices regarding conception and fertility: A population-based survey among reproductive-age United States women. *Fertility and Sterility*. 2014;101:767–774.e2. DOI: 10.1016/j.fertnstert.2013.12.006
11. Potter RG, Sagi PC, Westoff CF. Knowledge of the ovulatory cycle and coital frequency as factors affecting conception and contraception. *Milbank Mem Fund Q*. 2020;40:46–58. DOI: 10.2307/3348610
12. Hammarberg K, Setter T, Norman RJ, Holden CA, Michelmores J, Johnson L. Knowledge about factors that influence fertility among Australians of reproductive age: A population-based survey. *Fertility and Sterility*. 2013;99:502-507. DOI: 10.1016/j.fertnstert.2012.10.031

13. Ozsoy S, Adana F, Colakoglu HS. University students' knowledge about fertile period. *HealthMED*. 2012; 6:3174–9.
14. Ali S, Sophie R, Imam AM, Khan FI, Ali SF, Shaikh A. Knowledge, perceptions and myths regarding infertility among selected adult population in Pakistan: A cross-sectional study. *BMC Public Health*. 2011;11:760.
DOI: 10.1186/1471-2458-11-760
15. Iyanda AE. Women's reproductive health cross-national variation in knowledge of ovulation timing in sub-saharanafrica cross-national variation in knowledge of ovulation timing in sub-saharan Africa. *Women's Reproductive Health*. 2020;7:127–143.
DOI: 10.1080/23293691.2020.1740484
16. Iyanda AE. Women's Reproductive health cross-national variation in knowledge of ovulation timing in sub-saharanafrica cross-national variation in knowledge of ovulation timing in Sub-Saharan Africa. *Women's Reproductive Health*. 2020;7:127–143.
DOI: 10.1080/23293691.2020.1740484
17. Ajayi AA, Marangu LT, Miller J, Paxman JM, Ajayi AA, Marangu LT. Practices adolescent sexuality and fertility in kenya: A survey of knowledge, perceptions, and practices. *Stud Fam Plann*. 2020;22:205–216.
DOI: 10.2307/1966477
18. Stoddard A, McNicholas C, Peipert JF. Efficacy and safety of long-acting reversible contraception. *Drugs*. 2011;71(8):969–980.
19. Moreau C, Bohet A, Hassoun D. Trends and determinants of use of long-acting reversible contraception use among young women in france: Results from three national surveys conducted between 2000 and 2010. *Fertility and Sterility*. 2013; 100(2):451–458.
20. Belayneh F, Abreha S, Meskele M. Knowledge, attitude and factors associated with the use of long-acting and permanent contraceptives methods among women of reproductive age in Gesuba town, southern Ethiopia. *Journal of Biology, Agriculture and Healthcare*. 2015;5(21).
21. Zenebe CB, Adefris M, Yenit MK. Factors associated with the utilization of long-acting and permanent contraceptives methods among women who have decided not to have more children in Gondar city. *BMC Womens Health*. 2017;17(1): 75.
DOI: 10.1186/s12905-017-0432-9
22. Juayire CA. Knowledge of Ovulatory Cycle and Current Fertility Among Women in Ghana. University of Ghana, Legon; 2016.
23. Getahun MB, Nigatu AG. Knowledge of the Ovulatory Period and Associated Factors Among Reproductive Women in Ethiopia: A Population-Based Study Using the 2016 Ethiopian Demographic Health Survey. *International Journal of Women's Health*. 2020;12:701–707.
Available: <https://doi.org/10.2147/IJWH.S267675>
24. Hampton K, Mazza D. Fertility-awareness knowledge, attitudes and practices of women attending general practice. *Australian Family Physician*. 2015;44(11):840.
25. Mahey R, Gupta M, Kandpal S. Fertility awareness and knowledge among Indian women attending an infertility clinic: A cross-sectional study. *BMC Women's Health*. 2018;18(1): 177.
26. Bolarinwa OA, Olagunju OS. Knowledge and factors influencing long-acting reversible contraceptives use among women of reproductive age in Nigeria. *Gates Open Research*. 2020; 3:7.
Available: <https://doi.org/10.12688/gatesopenres.12902.3>
27. White K, Hopkins K, Potter JE, Grossman D. Knowledge and attitudes about long-acting reversible contraception among latina women who desire sterilization. *Women's Health Issues: Official Publication of The Jacobs Institute Of Women's Health*. 2013;23(4):e257–e263.
Available: <https://doi.org/10.1016/j.whi.2013.05.001>
28. Baku PM, Nyarko SH. Ghanaian male adolescents' knowledge about female fertile period. *Ghana Journal of Geography*. 2018;10:23–35.

- DOI: 10.4314/gjg.v10i2.2
29. Boah M, Bordotsiah S, Kuurdong S. Predictors of unsafe induced abortion among women in Ghana. Journal of Pregnancy; 2019. DOI: 10.1155/2019/9253650

© 2024 Ese et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111557>