



International Journal of Environment and Climate Change

Volume 14, Issue 2, Page 149-160, 2024; Article no.IJECC.112492

ISSN: 2581-8627

(Past name: British Journal of Environment & Climate Change, Past ISSN: 2231-4784)

Global Climate Change and Its Effects on Medicinal and Aromatic Plants: A Review Article

**Saumya Shruti ^{a*}, Ashwin Trivedi ^b, K. B. Chaudhary ^a
and Jatinkumar Ghadiali ^a**

^a *Department of Plant Physiology, B.A.C.A, Anand Agriculture University, Anand-388 110,
Gujarat, India.*

^b *ICAR- Directorate of Medicinal and Aromatic Plants Research, Boriavi, Anand- 388 110,
Gujarat, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i23931

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112492>

Review Article

Received: 26/11/2023

Accepted: 29/01/2024

Published: 02/02/2024

ABSTRACT

Medicinal herbs have been used for a long time to cure health problems and prevent disease, including epidemics. Medicinal plants are used for different purposes and in diverse uses of human beings. People around them have a basic knowledge and using through their traditional medicine from the past several years. Natural and manmade calamities, such as rapid climate change, urban growth, industrial boom, overpopulation pollution, declining forest cover, habitat loss, over-harvesting, destructive harvesting, and floods, are the principal reasons of the reduction of wild medicinal plant species. The life cycles and distribution of the world's vegetation, especially wild medicinal plants, are being affected by climate change. The impact of global climate change on medicinal plants could have a huge influence, especially in terms of their utility in conventional medication systems and as economically useful plants. The current essay emphasises the

*Corresponding author: E-mail: shrutisaumya82@gmail.com;

importance of study to improve our understanding of climate influences on medicinal plants properties, phenology and metabolic responses. The purpose of this paper is to review the impact of abiotic variables on secondary metabolite formation and different adaptation measures as well as future research initiatives. Furthermore, such issues with global climate change will undoubtedly become more visible or immediate threats, putting further strain on medicinal plant species. The impact of global climate change on medicinal plants could negatively correlate with environment concerns parameters, especially in terms of their utility in conventional medication systems and as economically useful plants.

Keywords: Adaptation; climate change; medicinal plants; phenology; secondary metabolites.

1. INTRODUCTION

“Climate change has increasingly been considered as one of the most serious threats to humanity and all other forms of life on the planet. It is now beyond doubt that anthropogenic Greenhouse gases (GHG) emissions are the main cause for recently observed climate change. Global warming of 1.5°C relative to 1850–1900 would be exceeded during the 21st century under the intermediate, high and very high GHG emissions scenarios” [1]. “Warming at 1°C affects all continents and is generally larger over land than over the oceans in both observations and models” [2]. “Plant growth and yield will be positively and negatively affected by climate change. Diverging effects are caused by the rising CO₂ concentrations, higher temperature, changing precipitation patterns, changing water availability, increased frequency of weather extremes (i.e., floods, heavy storms, and droughts), climate-induced soil erosion, and sea-level rise. Medicinal plants have been used as a source of medicine in practically all cultures. Humans employ medicinal plants for a variety of goals and in a variety of ways. Medicinal herbs have been used for a long time to cure health problems and prevent disease, including epidemics” [3]. People developed basic knowledge of medicinal plant values and how to treat various illnesses from an ancient day [4]. “Medicinal plants are essential natural resource which constitutes one of the potential sources of new products and bioactive compounds for drug development” [5]. “Traditional medicine, largely plant medications, is used by 60 percent of the world's population and 80 percent of the population of developing countries for primary health care” [6]. “According to reports, traditional plant-based remedies are used by 70% of India's population” [7]. “About 7500 species of higher plants in India are known to have therapeutic properties, accounting for a significant part of the total flowering plants” [8]. “Anthropogenic climate change has already had a significant impact on

global species ranges and ecological communities, including medicinal plants” [9]. “Climate change is “Unequivocal,” according to the Intergovernmental Panel on Climate Change (IPCC). Temperatures are expected to rise much further, from 1.4°C to 5.8°C by 2100 and 2300, respectively” [10]. “There is a need to recognise the pattern of climate change, which is one of the most serious global environmental concerns, and to understand and assess the various sorts of impacts” [11]. As the climate changes and the globe heats, there is a greater chance of biodiversity mass extinction than species can adapt to [12]. On the other hand, agricultural production industries and techniques are adapted to local climate variability [13], particularly with medicinal plants such as *Plantago ovata*, *Sisymbrium irio*, and many others [14] in arid and semi-arid settings. Local groups in diverse places that have used medicinal plants for generations have said that there are constant threats and a rapid loss of species worldwide. Climate change causes stress in plants, which can impact the secondary metabolites and other substances they generate, which are the basis for their therapeutic activity. It is already causing bio diversity and ecosystems to adapt to altering habitats, life cycles, and the emergence of new physical characteristics [15]. Plant species that are endemic to climate-vulnerable geographic locations may face serious threats in the near future [16] As a result, research on the effects of climate change on medicinal plants is critically insufficient [17]. In the therapy of any disease, the constituents of plants that serve as active ingredients interact simultaneously, complementing or neutralising the damage. The study on medicinal plants with respect to global climate change is very infrequent and insignificant in comparison with other commercial crops [18]. These Medicinal and Aromatic Plants (MAP's) species are critical sources of important biomolecules and nutraceuticals, and it is imperative that they not be abandoned. People

living in isolated and far from metropolitan areas are fully reliant on plants and their products for their basic existence and the treatment of various maladies and diseases, therefore the conservation of threatened medicinal plants at higher altitudes is critical [19]. "The demand for medicinal plants is currently increasing in both developed and developing countries for various reasons. Medicinal plants have a promising future since there are around half a million plants on the planet, and most of their pharmaceutical potential has yet to be discovered, necessitating current and future research" [20]. In compared to other commercial crops, research on medicinal plants in relation to climate change is intermittent and modest. It is likewise imperative for these plants to be protected because they are potential sources of biomolecules and nutraceuticals. While the future implications of climate change are likely to be unpredictable, they are already having an influence on MAPs and may become a much greater concern in the future. The disappearance of some MAPs could have a significant impact on the lives of a huge number of people. The problem of global warming and altered seasonal occurrences is similarly difficult to comprehend, but prompt action can likely avoid biodiversity loss. The method for assessing the influence of climate change on medicinal plants is based on contemporary literature, journal articles, and some personal observation.

2. THERAPEUTIC IMPORTANCE OF MEDICINAL AND AROMATIC PLANTS

"Secondary metabolites are chemical compounds that plants have evolved to synthesise in a wide variety of ways" [21]. "These secondary metabolites don't appear to play a part in basic plant growth and development, thus they're only found in plants from a single species, and their levels rise at times of extreme stress, such as drought, high temperature, or bacterial infection" [22]. "The glycosides are secondary metabolic products that include a variety of secondary metabolites coupled to mono or oligosaccharide and uronic acid" [23]. "Medicinal plants that contain cardio glycosides are commonly used in the treatment of heartfailure and cardiac arrhythmia. Cardioactive glycosides are found in a diverse group of plant species including *Digitalis purpurea* and *D. lanata* (Scrophulariaceae), *Nerium oleander* (Apocynaceae) and *Convallaria majalis* (Convallariaceae)" [24]. "Plant pigments such as flavonoids and proanthocyanidins can be found in a wide variety of plant groups. Isoflavones are

generated by Fabaceae species (bean family) Flavonoids are the most widespread and widely dispersed group of plant phenolic chemicals, found in almost every part of the plant, especially the photosynthetic plant cells" [25]. "In vascular plants, condensed tannins are the most widely distributed tannins. Hydrolyzable tannins, on the other hand, are only found in dicotyledons" (Bernays *et al.*, 1989). Fagaceae (beech family) and Polygonaceae are two plant groups that have been linked to the occurrence of tannins (knotweed family). The biological actions of pyrrolizidine alkaloids, which include acute hepatotoxic, mutagenic, carcinogenic, teratogenic, anticancer, and neuroactive characteristics, make them important secondary metabolites [26].

3. CLIMATE CHANGE EFFECTS DUE TO ENVIRONMENTAL POLLUTION

Environmental challenges all over the world. However, there is need to become increasingly aware of the problems that surround it. Some are minor and only influence a few ecosystems, while others have a significant impact on the environment. We are in the midst of a global emergency, with environmental issues stacking up all around us. Current environmental issues demand immediate attention.

Environmental pollution has existed for millennia, but it was only during the industrial revolution in the 19th century that it became serious. When the natural environment is unable to degrade an element without causing injury or damage to itself, pollution occurs. The elements involved are not found in nature, and the process of destruction can take anything from a few days to thousands of years.

3.1 Major Global Pollution Factors

- a) Industries: "As previously stated, industries have been contaminating our environment since the beginning of the industrial revolution, particularly due to the increased usage of fossil fuels. Coal was used to help machines work quicker in the 19th century and throughout much of the 20th century, substituting human labour" [27].
- b) Agricultural Activities: "Agriculture is mostly responsible for water and soil contamination. The increased use of pesticides, as well as the intensive nature of its manufacturing, are to blame.

Pesticides are almost often made of chemical components and are used to keep illnesses and dangerous animals ie. deer, wild pigs, wild rabbits, moles., away from crops" [28].

- c) "The production and exchange of commodities and services are examples of trading activities. When it comes to commodities, pollution is primarily generated by packing (which frequently entails the use of fossil-fuel-derived plastic) and transportation" [29].

4. PHENOLOGICAL CHANGES

"Plant life cycles follow seasonal patterns, and global climate change is altering species and ecosystems. These phenological changes are putting endangered medicinal plant species in jeopardy. Bud burst and leaf unfolding, flowering and establishing fruit, autumn or dry season leaf drop, and the linked processes of winter hardening and breaking are all important phenological events for medicinal plants that have adapted to climatic change [30]. The advent of spring and the length of the growing season will be affected as global warming progresses" (Bidart *et al.*, 2008).

The alteration of plant phenology in *Trifolium repens* by 1–11 days has been noted as a result of increased concentrations of carbon dioxide (CO₂) and elevated temperatures compared to the normal conditions [31]. *Aconitum heterophyllum* exhibits early flowering when exposed to elevated CO₂ levels [32]. Elevated CO₂ and temperature lead to a significant reduction in the duration taken for flower initiation, 50% flowering, seed setting, and seed maturity in *Valeriana jatamansi* [33]. *Hypericum perforatum* experiences a decreased timeframe from bud formation to flower formation under elevated CO₂ and temperature conditions [34].

5. GLOBAL WARMING AND CLIMATE CHANGE

Another environmental issue that has surfaced in recent decades is climate change. The phrase "climate change" has become synonymous with modern global warming for the purposes of current and future discussions [35]. "The latter, in turn, refers to variations in global mean surface-air temperature after the Industrial Revolution that are thought to have been driven by rising atmospheric carbon dioxide (CO₂). An active greenhouse gas throughout the

eighteenth, nineteenth, and twentieth centuries. CO₂ levels in the atmosphere were around 280 parts per million by volume (ppmv) in 1765, and around 364 parts per million by volume (ppmv) in 2000, (ppmv). Other greenhouse gases (methane, nitrous oxide, chlorofluorocarbons) have also increased in the atmosphere as a result of increased economic activity and energy use during the last two centuries [36]. The cause-and-effect relationship between the increase in important greenhouse gases and the predicted rise in global mean surface-air temperature is yet unknown. This is owing to the earth's climate's complicated fluctuation and interconnectedness with a variety of terrestrial and interplanetary processes" [37].

Medicinal plants, particularly endemic medicinal plants, are extremely vital to human survival [38]. Anthropogenic climate change has already had an impact on medicinal plants all across the planet. For example, heavier monsoon rainfall, more frequent and hotter summer days, less frequent and lower dry season rainfall, stronger and more frequent storms with high winds, and so on. Climate change, habitat damage, overharvesting, and a decline in natural pollinators as a result of overharvesting are all major factors influencing endemic populations [39].

6. METABOLIC RESPONSES OF PLANTS

Change in Flowering Time

Crop blossoming and maturation alter as the temperature rises. A 1°C increase in mean seasonal temperature reduces wheat output by 5–8% [40]. The time of flowering is a vital period of development in the life cycle of most plants when seed quantity is determined. When temperatures (>32–36 °C) coincide with a small crucial period of only 1–3 days around the time of blooming, it has been discovered that seed set and, ultimately, crop output are considerably reduced [41].

7. EFFECT OF ELEVATED CO₂ ON PLANT DEVELOPMENT AND MORPHOLOGY

CO₂ has a direct influence on plants because of photosynthetic gas exchange, but it also has an indirect effect as a potent greenhouse gas that contributes to global warming [42]. CO₂ increases photosynthetic carbon absorption rates by 31 percent on average across 40 species [43].

Heat has detrimental effects on plant leaves, such as reduced leaf water potential, reduced leaf area, and premature leaf senescence, all of which have a negative impact on the plant's overall photosynthetic performance. The lower activity of sucrose phosphate synthase, ADP-glucose pyro phosphorylase, and invertase affects starch and sucrose synthesis [44].

8. HEAT STRESS AND SECONDARY METABOLITE PRODUCTION

Plant metabolism aids heat stress tolerance by supplying energy and producing secondary metabolites that are necessary for cellular homeostasis. Temperature has a significant impact on plant secondary metabolite synthesis [45]. Carotenoids are required for photoprotection during photosynthesis and serve as a precursor in signalling during plant development under stressful situations. They improve plant output and nutritional quality. According to many findings, plants exposed to high-temperature stress have reduced chlorophyll production [46], which is the first step in plastids to be damaged or destroyed by the high temperature.

9. HIGH TEMPERATURE AND OSMOLYTE ACCUMULATION

Many plants use the buildup of low molecular weight water-soluble molecules known as "osmolytes" to combat environmental stressors.

The most prevalent defensive mechanism used by plants in response to high temperatures is the production and accumulation of suitable solutes. Amino acids (asparagine, proline, serine), amines (polyamines and glycine betaine), and γ -amino-N-butyric acid are among the compounds (GABA) [47]. Fructose, sucrose, trehalose, raffinose, and proline are all total sugars [48,49] as well as pools of anti-oxidants such as glutathione (GSH) and ascorbate [50] accumulate in response to heat stress.

10. PHYSIOLOGICAL ASPECTS OF HEAT STRESS

Heat stress causes plants to adapt their physiological status and metabolic level, resulting in both permanent and temporary metabolic changes. After drought, heat, and combination stress treatments, Jin *et al.*, (2015) detected changes in physiological and metabolic parameters in *Portulaca oleracea* L. Furthermore, under heat conditions, the content of the amino acid asparagine was enhanced. This is due to the connection of amino acids with protein synthesis substrates, which may aid in the rapid recovery of plant metabolism following stress. Glycine betaine is crucial in plants that are subjected to high temperatures [51]. Glycine betaine is generated in chloroplasts and keeps Rubisco active by sequestering Rubisco activase near thylakoids, preventing it from being thermally inactivated [52].




Fig. 1. Molecular responses in medicinal plants to heat stress

(source: [53])

11. EFFECTS OF OZONE ON PLANT SECONDARY METABOLITES

The bioprotective properties of ozone against ultra violet radiation is well established. Global agricultural yields are already being reduced by ground-level ozone pollution (from 2.2–5.5 percent for maize to 3.9–15% and 8.5–14% for wheat and soybean, respectively), to varying degrees depending on genotype and climatic conditions [54]. These negative consequences can also be recognised in medicinal herbs. However, because there has been so little attention paid to O₃'s effects on medicinal plants and their PSM content, it has become necessary to assess its potential repercussions (Basaran *et al.*, 2020).

Melissa officinalis, a traditional medicinal plant used to treat dementia, anxiety, and CNS-related disorders, was found to have a significant increase in total anthocyanins, as well as phenolics and tannins, when exposed to elevated ozone concentrations [55]. A similar investigation on the Eco physiological and antioxidant features of *Salvia officinalis* under ozone stress (120 ppb for 90 days) revealed an increase in phenolic content, particularly Gallic acid (2-fold increase)

and Catechic acid (rise only once over the 90-day fumigation period) [56].

12. EFFECTS OF TOXIC GASES ON PLANT SECONDARY METABOLITES

Sulfur dioxide is a prominent atmospheric pollutant that can enter the plant system through the roots as well as through stomatal opening during photosynthesis and respiration. Damage to the photosystem [57], changes in stomatal density, and variations in carbon fixation efficiency are some of the reactions [58]. The atmospheric SO₂ along with H₂S also acts as sulfur source which can be up-taken through stomata of the plants apart from the sulphate uptake from the roots. When *Nicotiana tabacum* plants are forced to grow at a high temperature, hydrogen sulphide has been found to mediate nicotine production [59].

13. EFFECT OF ULTRAVIOLET RADIATION

According to the International Commission on Illumination, Ultraviolet (UV) wavelength (400–200 nm) is a small part of the solar radiation reaching the Earth's surface, which is divided into UV-A (315–400 nm), UV-B (280–315 nm)

Table 1: Effects of ozone and carbon dioxide on plant secondary metabolite

Component	Treatment	Affected medicinal plant	Affected secondary metabolite
O ₃	Elevated O ₃	<i>Salvia officinalis</i>	Gallic acid↑, Catechic acid↑, Caffeic acid↑, Rosmarinic acid↑
CO ₂	Elevated CO ₂ and Light intensity	<i>Labisia pumila</i>	Flavonoids↑, Phenolics↑
CO ₂	Elevated CO ₂	<i>Catharanthus roseus</i>	Phenolics↑, Flavonoids↑, Tannins↑, Alkaloids↑
O ₃	Elevated O ₃	<i>Melissa officinalis</i>	Phenolics↑, Anthocyanins↑, Tannins↑
O ₃	Elevated O ₃	<i>Pueraria thomsnii</i>	Puerarin↑, ABA↑
O ₃	Elevated O ₃	<i>Hypericum perforatum</i>	Hypericin↑
CO ₂	Elevated CO ₂	<i>Zingiber officinale</i>	Flavonoids↑, Phenolics↑
O ₃ and CO ₂	Elevated O ₃ and CO ₂	<i>Ginkgo biloba</i>	Tannins↓, Quercetinaglycon↑, Keampferolaglycon↓, Isorhamnetin↓, Bilobalide↓
CO ₂	Elevated CO ₂	<i>Digitalis lanata</i>	Digoxin↑, Cardenolide↑
O ₃	Elevated O ₃	<i>Betula pendula</i>	Dehydrosalidroside ↓, hyperoside↓, Betuloside↓, Platyfyloside↓, Salidroside↓, papyriferic acid↓, hyperoside↑
CO ₂	Elevated CO ₂	<i>Papaver setigerum</i>	Morphine↑, Codeine↑, Papaverine↑ and Noscapine↑

(Source: V.Kumar *et al.*, 2010)

and UV-C (200–280 nm), and it affects negatively all living organisms. The harmful effect of UV radiation increases towards the shorter wavelengths [60]. These rays can harm molecular and cellular structures, including proteins, DNA, and other biopolymers. This sort of radiation can also alter plant growth and development, causing changes in vegetative or reproductive biomass, height, leaf features, and flowering time [61].

14. IMPACT OF UV RADIATION ON THE SYNTHESIS OF SECONDARY METABOLITES IN MEDICINAL PLANTS

In vitro cultures of *Phyllanthus tenellus* grown under white light plus UV-A radiation produced more antioxidants like phenolics (geraniin and ellagic acid) and carotenoids than plants cultivated under white light alone (Victorio *et al.*, 2011). Seedlings of *Laurus nobilis* cultivated under low doses of supplementary UV-A had lower leaf concentrations of particular quercetin and kaempferol derivatives than plants grown under ambient UV light (Bernal *et al.*, 2015). After 3 and 5 days of continuous UV-A exposure, the content of total phenolics and total flavonoids in sowthistle (*Ixeris dentata* Nakai) was dramatically increased, while plant development was not impeded (Lee *et al.*, 2013).

15. EFFECT ON THREATS TO MEDICINAL PLANTS SPECIES

Species that rely on one other (co-existence) may be pushed to extinction if they no longer coexist in the same time or space. Pests, illnesses, and invasive species may expand their ranges, putting additional stress on vulnerable populations, including medicinal plants. Conservation-minded species with rigorous habitat requirements or prolonged incubation times are more vulnerable to extinction [62].

16. EFFECT OF SALINITY STRESS ON THE GROWTH AND PHYSIOLOGY OF MEDICINAL PLANTS

Approximately 20% of the world's irrigated soil has a high salt level, making it harmful to agriculture [63]. As much as 25% of the total drugs in developed countries like USA constitutes of plant drugs, whereas it is close to 80% in developing countries like China and India. The cultivation and maintenance of medicinal plants thus forms a profitable business throughout the world and must be protected.

Cultivating medicinal plants on salt-contaminated soils has resulted in significant physiology and growth problems [64].

17. CORRELATION BETWEEN SALT STRESS AND DEVELOPMENTAL MORPHOLOGY IN MEDICINAL PLANTS

Salt stress inhibits germination by either killing the embryo or significantly lowering the soil potential, causing water intake to be impeded. When *Ocimum basilicum*, *Eruca sativa*, *Petroselinum hortense*, *Matricaria chamomila*, *Origanum marjoram*, *Thymus maroccanus*, and *Achillea fragratissima* seeds were placed in salt polluted soil, germination was delayed [65-67]. Rastogi *et al.*, 2022. *Mentha piperita* var. *officinalis* and *Lipia citriodora* var. *verbena* subjected to salt stress had considerably less leaves, leaf area, and leaf biomass [68].

18. CORRELATION BETWEEN SALT STRESS AND PHOTOSYNTHESIS IN MEDICINAL PLANTS

The most significant and crucial physiological mechanism for plant development and survival is photosynthesis. This mechanism necessitates a high level of energy expenditure for maintenance, making it vulnerable to stress circumstances that alter the cell's metabolic equilibrium (Banerjee and Roychoudhury, 2017). In *Teucrium polium*, *Thymus vulgaris*, *Zataria multiflora*, *Ziziphora clinopodioides*, and *Satureja hortensis*, both chlorophyll a and b, as well as total chlorophyll content, were reduced [69]. *Hypericum perforatum* exhibited a 124% increase compared to the controlled conditions when exposed to 1000 $\mu\text{mol mol}^{-1}$ of carbon dioxide (Walia *et al.*, 2022). In *Podophyllum hexandrum*, there was an initial increase in response to elevated CO₂ during the first 30 days, followed by a subsequent decrease [70]. Additionally, *Trifolium repens* displayed a down-regulation of photosynthesis at elevated CO₂, indicating a reduction in the photosynthetic process under these conditions [71].

19. ADAPTATION MEASURES FOR CLIMATE CHANGE AND GLOBAL WARMING

Conservation of endangered flora and fauna can reduce the vulnerabilities of medicinal plants in the world. These will involve medicinal plant cultivation. Maintaining genetic diversity in

natural ecosystems is critical for medicinal plant adaptability [72]. Traditional indigenous knowledge that can help mitigate the effects of climate change should be investigated and documented, while supporting traditional art and craft-based livelihoods through local community involvement and eco-tourism as a source of revenue [73-79].

20. CONCLUSION

Over the centuries, the use of medicinal and aromatic plants has become an important part of daily life despite the progress in modern medical and pharmaceutical industry. Endangered plant species are thought to be particularly vulnerable to climate change and may face extinction. As a result, a better understanding of the causes that cause such change need a comprehensive and ongoing field study. As the Earth's environment is constantly in battle with it, several things have an impact on it. Plants exposed to severe temperatures generally exhibit a similar response known as oxidative stress, which damages lipids, proteins, and nucleic acids. As a nutshell, for significant medicinal plants, the reaction of the plant and the mechanisms underlying its adaptation to elevated temperatures must be better understood. In addition, metabolic changes in response to stress are critical for developing stress tolerance. In analysis to define if there is an induction of hazardous metabolites, it is necessary to analyse not only the chemicals of interest for a certain plant, but also a large number of secondary metabolites. In view of the steadily rising demands on these important natural resources, attention should be paid to the sustainable forms of production and utilization. Furthermore, more research on threatened medicinal plants in a climate-change scenario is necessary for developing conservation strategies, as is cultivating medicinal plants through the involvement of local communities with traditional indigenous knowledge, which can help to mitigate the effects of climate change.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. IPCC (2021). Contribution of working group II to the Fourth assessment report of the intergovernmental panel on climate

change. Parry OF, Canziani JP, Palutikof, PJ, Van der Linden, Hanson (Eds) Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA; 2007.

2. Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC BF, De Siqueira MF, Grainger A, Hannah L, Williams SE. Extinction risk from climate change. *Nature*. 2004;427(6970):145-148.
3. Mishra T. Climate change and production of secondary metabolites in medicinal plants: A review. *International Journal of Herbal Medicine*. 2016;4(4):27-30.
4. Bhat SG. Medicinal plants and its pharmacological values. *Natural Medicinal Plants*. 2021;12:217-228.
5. Gangwar KK, Deepali GR, Gangwar RS. Ethnomedicinal plant diversity in Kumaun himalaya of Uttarakhand. *Indian National Science Academy*. 2010;8(5):66-78.
6. Shrestha PM, Dhillion SS. Medicinal plant diversity and use in the highlands of Dolakha district, Nepal. *Journal of Ethnopharmacology*. 2003;86(1):81-96.
7. Gadgil M, Seshagiri Rao PR. Nurturing biodiversity. *Centre for Environment Education*;1998.
8. Gowthami R, Sharma N, Pandey R, Agrawal A. Status and consolidated list of threatened medicinal plants of India. *Genetic Resources and Crop Evolution*. 2021;68(6):2235-2263.
9. Magnan AK, Portner HO, Duvat VK, Garschagen M, Guinder VA, Zommers Z, Gattuso JP. Estimating the global risk of anthropogenic climate change. *Nature Climate Change*. 2021;11(10):879-885.
10. Nazarenko L, Schmidt GA, Miller RL, Tausnev N, Kelley M, Ruedy R, Zhang J. Future climate change under RCP emission scenarios with GISS ModelE2. *Journal of Advances in Modeling Earth Systems*. 2015;7(1), 244-267.
11. Cavaliere C. The effects of climate change on medicinal and aromatic plants. *Herbal Gram*. 2009;81:44-57.
12. Muluneh MG. Impact of climate change on biodiversity and food security: A global perspective- a review article. *Agriculture & Food Security*. 2021;10(1):1-25.
13. Marshall E, Aillery M, Malcolm S, Williams R. Agricultural production under climate change: The potential impacts of shifting regional water balances in the United

- States. American Journal of Agricultural Economics. 2015;97(2):568-588.
14. Datta P, Behera B. Climate change and Indian agriculture: A systematic review of farmers perception, adaptation, and transformation. Environmental Challenges. 2022;8:100543.
 15. Srivastava AK, Mishra P, Mishra AK. Effect of climate change on plant secondary metabolism: An ecological perspective. In Evolutionary Diversity as a Source for Anticancer Molecules. Academic Press. 2021;47-76.
 16. Manes S, Costello MJ, Beckett H, Debnath A, Devenish-Nelson E, Grey KA, Vale MM. Endemism increases species' climate change risk in areas of global biodiversity importance. Biological Conservation. 2021;257:109070.
 17. Parmesan C. Ecological and evolutionary responses to recent climate change. Annuals Review Ecological System. 2006;37:637-639.
 18. Applequist WL, Brinckmann JA, Cunningham AB, Hart RE, Heinrich M, Katerere DR, Van Andel T. Scientists' warning on climate change and medicinal plants. Planta Medica. 2020;86(01):10-18.
 19. Ullah S, Ullah A, Rashid A. Medicinal diversity of weeds in the historical valley of Landikotal, Khyber Agency, Pakistan, Pakistan Journal of Weed Science Research. 2014;20(4):531-539.
 20. Ghorbanpour M, Hatami M, Khavazi K. Role of plant growth promoting rhizobacteria on antioxidant enzyme activities and tropane alkaloid production of *Hyoscyamus niger* under water deficit stress. Turkish Journal of Biology. 2013;37(3):350-360.
 21. Yeshi K, Crayn D, Ritmejeriyè E, Wangchuk P. Plant secondary metabolites produced in response to abiotic stresses has potential application in pharmaceutical product development. Molecules. 2022;27(1):313.
 22. Taiz L, Zeiger E. Plant physiology. Sinauer Associates Inc., Sunderland, Massachusetts, USA. [A general text of plant physiology]; 2006.
 23. Wang J, Hou B. Glycosyltransferases: Key players involved in the modification of plant secondary metabolites. Frontiers of Biology in China, 2009;4:39-46.
 24. Bernhoft A, Siem H, Bjertness E, Meltzer M, Flaten T, Holmsen E. Bioactive compounds in plants—benefits and risks for man and animals. The Norwegian Academy of Science and Letters, Oslo. 2010;13–14.
 25. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. The scientific World Journal; 2013.
 26. Asres K, Sporer F, Wink M. Patterns of pyrrolizidine alkaloids in 12 Ethiopian *Crotalaria* species. Biochemical Systematics and Ecology. 2004;32(10): 915-930.
 27. Malik A, Sharma S, Batra I, Sharma C, Kaswan MS, Garza-Reyes JA. Industrial revolution and environmental sustainability: an analytical interpretation of research constituents in Industry 4.0. International Journal of Lean Six Sigma. 2024;15(1):22-49.
 28. Sun S, Sidhu V, Rong Y, Zheng Y. Pesticide pollution in agricultural soils and sustainable remediation methods: A review. Current Pollution Reports, 2018;4: 240-250.
 29. Andrady AL. (Ed.). Plastics and the Environment. John Wiley & Sons; 2003.
 30. Rana SK, Rana HK, Ranjitkar S, Ghimire SK, Gurmachhan CM, O'Neill AR, Sun H. Climate-change threats to distribution, habitats, sustainability and conservation of highly traded medicinal and aromatic plants in Nepal. Ecological Indicators. 2020;115:106435.
 31. Kumar R, Kaundal M, Jandrotia R, Vats SK. Free-air CO₂ enrichment (FACE) and free-air temperature increase (FATI) effects on *Trifolium repens* in temperate Himalayas. International Journal of Plant Production. 2020;14:43-56.
 32. Chandra S, Chandola V, Nautiyal MC, Purohit VK. Elevated CO₂ causes earlier flowering in an alpine medicinal herb *Aconitum heterophyllum* Wall. Current Science. 2020;118(11):1650-1651.
 33. Kaundal M, Kumar R. Effect of elevated CO₂ and elevated temperature on growth and biomass accumulation in *Valeriana jatamansi* Jones under different nutrient status in the Western Himalaya. Journal of Agrometeorology. 2020;22(4):419-428.
 34. Sharma S, Walia S, Rathore S, Kumar P, Kumar R. Combined effect of elevated CO₂ and temperature on growth, biomass and secondary metabolite of *Hypericum perforatum* L. in a western Himalayan region. Journal of Applied Research on

- Medicinal and Aromatic Plants. 2020B;16: 100239.
35. Demeritt D. The construction of global warming and the politics of science. *Annals of the Association of American Geographers*. 2001;91(2):307-337.
 36. Rowntree L, Lewis M, Price M, Wyckoff W. *Globalization and diversity: Geography of a changing world*. Pearson; 2016).
 37. Loaiciga HA. Climate change and ground water. *Annals of the American Association of Geographers*. 2003;93(1):30–41
 38. Anderson DM, Salick J, Moseley RK, Xiaokun O. Conserving the sacred medicine mountains: A vegetation analysis of Tibetan sacred sites in Northwest Yunnan. *Biodiversity & Conservation*. 2005;14(13):3065-3091.
 39. Byg A, Salick J. Local perspectives on a global phenomenon—climate change in Eastern Tibetan villages. *Global Environmental Change*. 2009;19(2):156-166.
 40. Wheeler TR, Batts GR, Ellis RH, Hadley P, Morison JIL. Growth and yield of winter wheat (*Triticum aestivum*) crops in response to CO₂ and temperature. *The Journal of Agricultural Science*. 1996;127(1):37-48.
 41. Amasino RM, Michaels SD. The timing of flowering. *Plant Physiology*. 2010;154(2): 516-520.
 42. Cassia R, Nocioni M, Correa-Aragunde N, Lamattina L. Climate change and the impact of greenhouse gasses: CO₂ and NO, friends and foes of plant oxidative stress. *Frontiers in Plant Science*. 2018;9: 273.
 43. Reddy AR, Rasineni GK, Raghavendra AS. The impact of global elevated CO₂ concentration on photosynthesis and plant productivity. *Current Science*. 2010;46-57.
 44. Baroja-Fernández E, Muñoz FJ, Montero M, Etxeberria E, Sesma MT, Ovecka M, Bahaji A, Ezqer I, Li J, Prat S, Pozueta-Romero J. Enhancing sucrose synthase activity in transgenic potato (*Solanum tuberosum* L.) tubers results in increased levels of starch, ADPglucose and UDP glucose and total yield. *Plant and Cell Physiology*. 2009;50(9):1651-1662.
 45. Kumar S, Korra T, Thakur R, Arutselvan R, Kashyap AS, Nehela Y, Chaplygin V, Minkina T, Keswani C, Keswani C. Role of Plant Secondary Metabolites in Defence and Transcriptional Regulation in Response to Biotic Stress. *Plant Stress*. 2023;100154.
 46. Reda F, Mandoura HMH. Response of enzymes activities, photosynthetic pigments, proline to low or high temperature stressed wheat plant exogenous proline or cysteine. *International Journal of Academic Research*. 2011;3:108–116.
 47. Rezaei-Chiyaneh E, Seyyedi SM, Ebrahimian E, Moghaddam SS, Damalas CA. Exogenous application of gamma-aminobutyric acid (GABA) alleviates the effect of water deficit stress in black cumin (*Nigella sativa* L.). *Industrial Crops and Products*. 2018;112:741-748.
 48. Wani SH, Gosal SS. Genetic engineering for osmotic stress tolerance in plants-role of proline. *IUP Journal of Genetics & Evolution*. 2010;3(4).
 49. Harsh A, Sharma YK, Joshi U, Rampuria S, Singh G, Kumar S, Sharma R. Effect of short-term heat stress on total sugars, proline and some antioxidant enzymes in moth bean (*Vigna aconitifolia*). *Annals of Agricultural Sciences*. 2016;61(1):57-64.
 50. Ding X, Jiang Y, He L, Zhou Q, Yu J, Hui D, Huang D. Exogenous glutathione improves high root-zone temperature tolerance by modulating photosynthesis, antioxidant and osmolytes systems in cucumber seedlings. *Scientific Reports*. 2016;6:35424
 51. Oddy J, Raffan S, Wilkinson MD, Elmore JS, Halford NG. Stress, nutrients and genotype: Understanding and managing asparagine accumulation in wheat grain. *CABI Agriculture and Bioscience*. 2020;1: 1-14.
 52. Allakhverdiev SI, Kreslavski VD, Klimov VV, Los DA, Carpentier R, Mohanty P. Heat stress: an overview of molecular responses in photosynthesis. *Photosynthesis Research*. 2008;98(1):541-550.
 53. Kumar V, Shriram V, Kavi Kishor PB, Jawali N, Shitole MG. Enhanced proline accumulation and salt stress tolerance of transgenic indica rice by over-expressing P5CSF129A gene. *Plant Biotechnology Reports*. 2010;4(1):37-48.
 54. de Gruijl F, Leun J. Environment and health. Ozone depletion and ultraviolet radiation. *Canadian Medical Association Journal*. 2000;163(7):851-855.
 55. Shakeri A, Sahebkar A, Javadi B. *Melissa officinalis* L.- a review of its traditional

- uses, phytochemistry and pharmacology. Journal of Ethnopharmacology; 2016
DOI: 10.1016/j.jep.2016.05.010
56. Pellegrini E, Francini A, Lorenzini G, Nali C. Ecophysiological and antioxidant traits of *Salvia officinalis* under ozone stress. Environmental Science and Pollution Research. 2015;22(17):13083-13093.
 57. Lu Z, Zhang Q, Streets DG. Sulfur dioxide and primary carbonaceous aerosol emissions in China and India;;1996–2010. Atmospheric Chemistry and Physics. 2011;11(18):9839-9864.
 58. Harrison EL, Arce Cubas L, Gray JE, Hepworth C. The influence of stomatal morphology and distribution on photosynthetic gas exchange. The Plant Journal. 2020;101(4):768-779.
 59. Chen X, Chen Q, Zhang X, Li R, Jia Y, Ef A, Jia A, Hu L, Hu X. Hydrogen sulfide mediates nicotine biosynthesis in tobacco (*Nicotiana tabacum*) under high temperature conditions. Plant Physiology and Biochemistry. 2016;104:174-179.
 60. Kumari R, Prasad MNV. Medicinal plant active compounds produced by UV-B exposure. Sustainable Agriculture Reviews. 2013;12:225-254.
 61. Lei Y, Hannoufa A, Yu P. The use of gene modification and advanced molecular structure analyses towards improving alfalfa forage. International Journal of Molecular Sciences. 2017;18(2):298.
 62. Benning TL, La-Pointe D, Atkinson CT, Vitousek PM. Interactions of climate change with biological invasions and land use in the Hawaiian Islands: modelling the fate of endemic birds using a geographic information system. The Proceedings of the National Academy of Sciences (PANS). 2002;99(22):14246-14249.
 63. Zaman M, Shahid SA, Heng L, Shahid SA, Zaman M, Heng L. Soil salinity: Historical perspectives and a world overview of the problem. Guideline for salinity assessment, mitigation and adaptation using nuclear and related techniques. 2018;43-53.
 64. Munir N, Hasnain M, Roessner U, Abideen Z. Strategies in improving plant salinity resistance and use of salinity resistant plants for economic sustainability. Critical Reviews in Environmental Science and Technology. 2022;52(12):2150-2196.
 65. Ali RM, Abbas HM, Kamal RK. The effects of treatment with polyamines on dry matter, oil and flavonoid contents in salinity-stressed chamomile and sweet marjoram. Plant Soil and Environment. 2007;53(12):529.
 66. Abd EL-Azim WM, Ahmed ST. Effect of salinity and cutting date on growth and chemical constituents of *Achillea fragratissima* Forssk, under Ras Sudr conditions. Research Journal of Agriculture and Biological Sciences. 2009; 5(6):1121-1129.
 67. Mir TA, Jan M, Khare RK, Dhyani S, Saini N. Influence of Salinity on the Growth, Development, and Primary Metabolism of Medicinal Plants. In Environmental Challenges and Medicinal Plants: Sustainable Production Solutions under Adverse Conditions. Cham: Springer International Publishing. 2022;339-353.
 68. Chrysargyris A, Solomou M, Petropoulos SA, Tzortzakis N. Physiological and biochemical attributes of *Mentha spicata* when subjected to saline conditions and cation foliar application. Journal of Plant Physiology. 2019;232:27-38.
 69. Chauhan J, Singh P, Choyal P, Mishra UN, Saha D, Kumar R, Anuragi H, Pandey S, Bose B, Mehta B, Singhal RK. Plant Photosynthesis Under Abiotic Stresses: Damages, Adaptive, and Signaling Mechanisms. Plant Stress. 2023; 100296.
 70. Ghasemzadeh A, Jaafar HZ, Karimi E, Ibrahim MH. The combined effect of CO₂ enrichment and foliar application of salicylic acid on the production and antioxidant activities of anthocyanin, flavonoids and isoflavonoids from ginger. BMC complementary and Alternative Medicine. 2012;12:1-10.
 71. von Caemmerer S, Ghannoum O, Conroy JP, Clark H, Newton PC. Photosynthetic responses of temperate species to free air CO₂ enrichment (FACE) in a grazed New Zealand pasture. Functional Plant Biology. 2001;28(6):439-450.
 72. Kadam ST, Pawar AD. Conservation of medicinal plants: A review. International Ayurvedic Medical Journal, 2020;8:3890-3895.
 73. Mooney HA, Winner WE, Pell EJ. Response of plants to multiple stresses. Academic Press, San Diego, California, USA; 1991.

74. Basaran N, Pasli D, Basaran AA. Unpredictable adverse effects of herbal products. *Food and Chemical Toxicology*, 2022;159:112762.
75. IPCC. Intergovernmental Panel on Climate Change. Working group, I report “The Physical Science Basis”, Working group II report “Impacts, Adaptation and Vulnerability”, Working group III report “Mitigation of Climate Change”. [Online]; 2007.
76. Jin R, Wang Y, Liu R, Gou J, Chan Z. Physiological and metabolic changes of purslane (*Portulaca oleracea* L.) in response to drought, heat, and combined stresses. *Frontiers in Plant Science*. 2016;6:1123.
77. WHO. Traditional Medicine Strategy. 2002;2014–2023. Available:<http://apps.who.int/iris/bitstream/10665/92455/1/9789241506090>
78. Zobayed SS, PK, Saxena PK. Production of St. John's wort plants under controlled environment for maximizing biomass and secondary metabolites. *In vitro Cellular & Developmental Biology-Plant*. 2004;40; 108-114.
79. Rastogi S, Shah S, Kumar R, Vashisth D, Akhtar MQ, Kumar A, Dwivedi UN, Shasany AK. *Ocimum* metabolomics in response to abiotic stresses: Cold, flood, drought and salinity. *PLoS One*. 2019;14(2): e0210903.

© 2024 Shruti et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112492>