



# Compatibility and Toxicity of *Lecanicillium lecanii* with Insecticides against Cotton Aphids *In vitro*

P. B. Nikhade <sup>a\*</sup>, S. K. Bhalkare <sup>a</sup>, Y. B. Patil <sup>a</sup>, Y. V. Ingle <sup>b</sup>,  
N. S. Satpute <sup>a</sup> and D. B. Undirwade <sup>a</sup>

<sup>a</sup> Department of Entomology, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola – 444104, Maharashtra, India.

<sup>b</sup> All India Coordinated Research Project on Fruits, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola – 444104, Maharashtra, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors PN, YP, SB and YI supervised the study, designed the work, reviewed and edited the manuscript. Authors DU and NS wrote and prepared the original draft. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i101428>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123472>

Original Research Article

Received: 11/07/2024

Accepted: 13/09/2024

Published: 18/09/2024

## ABSTRACT

**Background:** *Lecanicillium lecanii* is the most effective entomopathogenic fungus against all stages of sucking pests like aphids, whiteflies, scale insects, thrips and mealybugs. The compatibility of fungi with commonly used insecticides was investigated in this study.

**Aim:** To study the *In vitro* compatibility and toxicity of *Lecanicillium lecanii* with insecticides against cotton aphids.

**Study Design:** Completely Randomized Design.

\*Corresponding author: E-mail: [pragatinikh123@gmail.com](mailto:pragatinikh123@gmail.com);

**Cite as:** Nikhade, P. B., S. K. Bhalkare, Y. B. Patil, Y. V. Ingle, N. S. Satpute, and D. B. Undirwade. 2024. "Compatibility and Toxicity of *Lecanicillium lecanii* With Insecticides Against Cotton Aphids *In Vitro*". *Journal of Advances in Biology & Biotechnology* 27 (10):47-54. <https://doi.org/10.9734/jabb/2024/v27i101428>.

**Methodology:** Six different insecticides were evaluated under *in vitro* conditions against *L. lecanii* by the poison food technique. The results showed that insecticides, viz., Dinotefuran 20 SG (0.006%), Afidopyropen 50 g/L DC (0.1%), Diafenthiuron 50 WP (0.06%) and Buprofezin 25 SC (0.05%), were rated as harmless and proved compatible with *L. lecanii*. Pyriproxyfen 10 EC (0.016%) and Tolfenpyrad 15 EC (0.03%) were moderately harmful to *L. lecanii*. Furthermore, the toxicity of insecticides alone and in combination with the entomopathogenic fungus *L. lecanii* against the cotton aphid, *Aphis gossypii*, was assessed by topical application bioassay under laboratory conditions.

**Results:** Cent percent mortality of *A. gossypii* was recorded with Diafenthiuron 50 WP (0.06%), Dinotefuran 20 SG (0.006%) and Afidopyropen 50 g/L DC (0.1%) at the recommended dose and their combination at half of the recommended dose with *L. lecanii* ( $1 \times 10^7$  conidia/ml) at 72 h after treatment. While minimum mortality was observed with the individual treatment of *L. lecanii* ( $1 \times 10^7$  conidia/ml).

**Conclusion:** This study suggests that the most suitable insecticides at half of the recommended dose, when combined with *L. lecanii*, can be effectively integrated into pest management programs.

**Keywords:** *Aphis gossypii*; compatibility; insecticides; *Lecanicillium lecanii*; toxicity.

## ABBREVIATIONS

DC: Dispersible Concentrate;  
EC: Emulsifiable Concentrate;  
SC: Soluble Concentrate;  
SG: Soluble Granules;  
SDA: Sabouraud Dextrose Agar;  
WP: Wettable Powders.

## 1. INTRODUCTION

The cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae), is a versatile pest known for inducing substantial harm, such as leaf curling and deformation. Additionally, it acts as a carrier for over 76 viral diseases, including potyvirus, cucumber mosaic virus, and zucchini yellow virus, impacting various crops adversely [1]. Aphids alone have the potential to inflict yield losses of up to 82% in the case of cruciferous crops when insecticides are not applied [2]. The intensive use of insecticides to control aphids has led to populations that are now resistant to several classes of insecticides [3]. Moreover, the use of pesticides can lead to significant issues related to environmental contamination and can adversely affect beneficial insects like bee populations [4,5]. Biopesticides offer a route to protecting the crop while reducing the reliance on synthetic insecticides [6]. Entomopathogenic fungi (EPF) have been found to be effective as a biopesticide [7] and have the potential to minimize the target pest populations on multiple crops [8,9]. Another important fact to be considered in favor of these EPF is that, to date, there has been no report of developing resistance [10].

Entomopathogenic fungi are typically acknowledged for their slower action, requiring

more time than traditional methods to achieve adequate insect mortality. Incorporating these fungi into a management strategy along with faster-acting materials could offer a potential solution to this issue. The synergistic action of mycoinsecticides with chemical insecticides can increase mortality and reduce the time until death in insects [11,12]. In this study, we gauged the compatibility of different insecticides with *L. lecanii* and assessed their toxicity to a prominent aphid pest. The compatibility of EPFs with pesticides could simplify the process of choosing suitable products within IPM programs.

## 2. MATERIALS AND METHODS

### 2.1 Fungus Culture

*Lecanicillium lecanii* culture available with Centre for Organic Agriculture Research and Training, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola was used during the present study. This entomopathogenic fungus was mass multiplied on Sabouraud's dextrose medium and used for further studies.

### 2.2 *In vitro* Compatibility of *L. lecanii* with Insecticides

To assess the compatibility, the effect of different treatments on the radial growth of *L. lecanii* was evaluated. The recommended dose of insecticides was added to SDA (Sabouraud Dextrose Agar) in a 100 ml conical flask before solidification. Following thorough mixing, the media was transferred to Petri dishes and allowed to solidify with gentle shaking. The plates, after solidification, were inoculated centrally with a 6 mm disc of a young sporulating

culture of *L. lecanii* with the help of a sterilized cork borer and a fungal inoculating needle. The experiment on the compatibility of *L. lecanii* with insecticides consisted of seven treatments, each replicated three times. Petri dishes were sealed and placed in an incubator maintained at  $27 \pm 1$  °C and  $80 \pm 5\%$  relative humidity. The medium without pesticide was used as a control treatment. The observations on fungal diameter in each plate were recorded after 10 days of inoculation. Percentage inhibition of *L. lecanii* was calculated on the basis of the growth diameter of the colony using the formula of Hokkanen and Kutluoto [13].

The pesticides were further classified into evaluation categories based on a 1–4 scoring index, i.e., 1-harmless (<50% reduction), 2-slightly harmful (50–79%), 3-moderately harmful (80–90%), and 4-harmful (>90%) according to Hassan's classification (Hassan, 1989).

### 2.3 Toxicity of *L. lecanii* Alone and in Combination with Insecticides

A population of aphids was collected from the untreated cotton field of the Integrated Farming Unit, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola. Serial dilutions of the *L. lecanii* isolate, pesticides alone, and mixtures (*L. lecanii* + pesticides) were prepared for each treatment. Once sterilized using sodium hypochlorite (0.5% v/v), detached cotton leaves underwent three washes with distilled water, followed by air drying, before being positioned on 1.5% agar (non-nutritive) in plastic Petri dishes measuring  $90 \times 20$  mm<sup>2</sup>. This agar concentration provided moisture essential for sustaining relative humidity during the test. Approximately 30 aphids, comprising a mixture of adult and nymph populations, were allowed to settle on the leaves. A topical spray method was used to treat the aphids with individual and combined applications of pesticides with *L. lecanii*. Mortality data were recorded at 24, 48, and 72 hours post-treatment [14].

### 2.4 Statistical Analysis

A completely randomized design (CRD) was used in all experiments. The data obtained were converted to appropriate transformations, subjected to ANOVA, and means were compared by critical difference ( $p = 0.01$ ) [15].

## 3. RESULTS

### 3.1 *In Vitro* Compatibility of *L. lecanii* with Insecticides

The results on compatibility of *L. lecanii* with insecticides (Table 1) revealed that Dinotefuran 20 SG (0.006%) significantly supported the maximum radial mycelial growth (50.33 mm) of *L. lecanii* over the rest of the insecticides, and growth inhibition was 5.63 percent. This was followed by Afidopyropen 50 g/L DC (0.1%), Diafenthiuron 50 WP (0.06%), and Buprofezin 25 SC (0.05%), which recorded radial mycelial growth of 47, 45.66, and 31.83 mm and 11.87, 14.38, and 40.32 percent mycelial growth inhibition, respectively. The remaining treatments, Pyriproxifen 10 EC (0.016%) and Tolfenpyrad 15 EC (0.03%), with 10.50 and 9.33 mm radial mycelial growth and 80.31 and 82.50 percent mycelial growth inhibition, respectively, were found least compatible with *L. lecanii* in comparison to prior treatments. The insecticides Dinotefuran 20 SG (0.006%), Afidopyropen 50 g/L DC (0.1%), Diafenthiuron 50 WP (0.06%), and Buprofezin 25 SC (0.05%) showed less than 50 percent mycelial growth inhibition and are categorized under Grade 1, i.e., harmless, whereas treatments Tolfenpyrad 15 EC (0.03%) and Pyriproxifen 10 EC (0.016%) showed growth inhibition between 80 and 90 percent and were categorized under Grade 3, i.e., moderately harmful [16].

### 3.2 Toxicity of *L. lecanii* Alone and in Combination with Insecticides Against Cotton Aphids

The insecticidal treatments showed the same trend of efficacy against aphids on different days of observation (Table 2). The insecticides like Diafenthiuron 50 WP (0.06%), Dinotefuran 20 SG (0.006%), and Afidopyropen 50 g/L DC (0.1%) at the recommended dose and their combination at half of the recommended dose with *L. lecanii* showed a cent percent mortality in cotton aphid at 72 hours after treatment, followed by Buprofezin 25 SC (0.025%) + *L. lecanii*, Buprofezin 25 SC (0.05%), Pyriproxifen 10 EC (0.008%) + *L. lecanii*, Pyriproxifen 10 EC (0.016%), Tolfenpyrad 15 EC (0.015%) + *L. lecanii*, and Tolfenpyrad 15 EC (0.03%) recorded 97.78, 96.67, 94.44, 90.00, 87.78, and 85.56 percent aphid mortality. Comparatively less aphid mortality was recorded in the treatment of *L. lecanii* alone (40.00%).

**Table 1. Compatibility of *L. lecanii* with insecticides by Poison Food Technique**

Sr. No	Treatment	Concentration (%)	10 days after inoculation of <i>L. lecanii</i>		Grade*
			Mean of radial mycelial growth (mm)	% mycelial growth inhibition	
1	Buprofezin 25 SC	0.05	31.83 d	40.32	1
2	Diafenthuron 50 WP	0.06	45.66 bc	14.38	1
3	Dinotefuran 20 SG	0.006	50.33 ab	5.63	1
4	Pyriproxifen 10 EC	0.016	10.50 e	80.31	3
5	Tolfenpyrad 15 EC	0.03	9.33 e	82.50	3
6	Afidopyropen 50 g/L DC	0.1	47.00 bc	11.87	1
7	<i>L. lecanii</i> (Control)	-	53.33 a	-	-
'F' test		Sig.			
SE(m) ±		1.30			
CD (p = 0.01)		5.47			

Note: Different letter within the treatments denote significant differences in the same column.  
 \*Grade: 1 = harmless (<50% reduction in beneficial capacity), 2 = slightly harmful (50–79%),  
 3 = moderately harmful (80–90%), 4 = harmful (>90%) in in vitro toxicity tests (Hassan, 1989).

**Table 2. Toxicity of *L. lecanii* alone and in combination with insecticides against cotton aphids**

Sr. No.	Treatment	Concentration (%)	Per cent aphid mortality		
			24 hrs	48 hrs	72 hrs
1	Buprofezin 25 SC	0.05	26.67 f (31.06)*	63.33 e (52.75)	96.67gh (81.32)
2	Diafenthuron 50 WP	0.06	47.78 e (43.73)	77.78 cd (61.89)	100.00abcde (89.43)
3	Dinotefuran 20 SG	0.006	76.67 b (61.15)	100.00 a (89.43)	100.00 ab (89.43)
4	Pyriproxifen 10 EC	0.016	8.89gh (17.28)	40.00fg (39.22)	90.00ij (71.73)
5	Tolfenpyrad 15EC	0.03	2.22i (7.19)	23.33 h (28.85)	85.56jk (67.69)
6	Afidopyropen 50 g/L DC	0.1	63.33 cd (52.75)	93.33 b (75.36)	100.00abcd (89.43)
7	Buprofezin 25 SC + <i>L. lecanii</i>	0.025 + 1×10 <sup>7</sup> (conidia/ml)	31.11 f (33.90)	67.78 de (55.42)	97.78abcde (82.80)
8	Diafenthuron 50 WP + <i>L. lecanii</i>	0.03 + 1×10 <sup>7</sup> (conidia/ml)	56.67 de (48.84)	84.44 c (66.80)	100.00abcde (89.43)
9	Dinotefuran 20 SG + <i>L. lecanii</i>	0.003 + 1×10 <sup>7</sup> (conidia/ml)	85.56 a (67.69)	100.00 a (89.43)	100.00 a (89.43)
10	Pyriproxifen 10 EC + <i>L. lecanii</i>	0.008 + 1×10 <sup>7</sup> (conidia/ml)	14.44 g (22.31)	45.56 f (42.45)	94.44 hi (76.52)
11	Tolfenpyrad 15EC + <i>L. lecanii</i>	0.015 + 1×10 <sup>7</sup> (conidia/ml)	6.67 h (14.64)	34.44fgh (35.93)	87.78ijk (69.58)
12	Afidopyropen 50 g/L DC + <i>L. lecanii</i>	0.05 + 1×10 <sup>7</sup> (conidia/ml)	70.00bc (56.81)	95.56 b (80.16)	100.00abc (89.43)
13	<i>L. lecanii</i>	1×10 <sup>7</sup> (conidia/ml)	0.00 j (0.52)	6.67i (14.64)	40.00 l (39.19)
14	Untreated Control	-	0.00 k (0.52)	2.22 j (7.19)	5.56 m (13.48)
'F' test			Sig.	Sig	Sig
SE(m)±			1.40	1.99	1.77
CD (p = 0.01)			5.47	7.77	6.92

Note: Different letter within the treatments denote significant differences in the same column.

\*Figures in parentheses are corresponding arcsine transformed values

The value of 0% is substituted by (1/4n) and the value of 100% by (100-1/4n), where n is the number of units upon which the percentage data is based (i.e., the denominator used in computing the percentage).

## 4. DISCUSSION

### 4.1 *In vitro* Compatibility of *L. lecanii* with Insecticides

The insecticides resulted in varying degrees of inhibition of germination, vegetative growth, and sporulation in *L. lecanii*. This variation depends on the specific compounds that disrupt conidial metabolic functions and the concentrations of the active ingredients [17,18]. The findings regarding compatibility of *L. lecanii* with insecticides align with previous research by Kim [1], who observed no impact on the mycelial growth of *L. attenuatum* when exposed to neonicotinoid (Imidachloprid). Similarly, in this study, dinotefuran (also a neonicotinoid) demonstrated no harm to *L. lecanii*. Patel et al. [19] reported that dinotefuran and difenthiuron showed heightened compatibility with entomopathogenic fungi at recommended doses. Reddy et al. [20] corroborated that buprofezin posed no harm to *L. lecanii* at both recommended and half-recommended doses. Yadav et al. [21] revealed that tolfenpyrad was highly incompatible with tested entomopathogens, a result consistent with the decreased compatibility of *L. lecanii* observed with Tolfenpyrad, a pyrazole insecticide, in the current study. Tolfenpyrad can kill fungi by targeting complex I NADH oxidoreductase within the respiratory process [22]. Additionally, Barari et al. [23] demonstrated a significant reduction in conidial germination, vegetative growth, and spore production of *B. bassiana* when exposed to Pyriproxifen 10 EC.

### 4.2 Toxicity of *L. lecanii* Alone and in Combination with Insecticides Against Cotton Aphids

The findings from experiments testing toxicity of *L. lecanii* alone and in combination with insecticides against cotton aphids are supported by the research of Nawaz et al. [14] discovered that combining *M. anisopliae* with Dinotefuran and Pyriproxifen had a synergistic effect, resulting in increased mortality of cotton aphids. They concluded that the insecticides exhibited consistent toxicity aligned with their compatibility with *M. anisopliae*, ranking in efficacy exactly as they did for compatibility. Buprofezin is an insect growth regulator that primarily inhibits chitin synthesis, thereby impacting the development of aphids. Reddy et al. [20] found Buprofezin 25 SC, at recommended and varied concentrations, to be compatible with *L. lecanii*. Combining Buprofezin with entomopathogenic fungi

increased Brown Plant Hopper (BPH) mortality compared to using Buprofezin alone, with similar outcomes reported by Ijaz et al. [24] under laboratory and field conditions against *Sogatella furcifera*, *Akanthomyces lecanii*, and buprofezin. Halder et al. [25] noted that combining *L. lecanii* with neonicotinoid insecticides resulted in co-toxicity coefficient values (CTC) >1 and lower LT<sub>50</sub> values than using each independently, indicating compatibility and synergistic action. Combining *L. lecanii* with appropriate sub-lethal concentrations of neonicotinoids in a two-in-one tank mixture against sucking insect pests alleviates pressure on insecticide selection and curbs simultaneous resistance development in targeted pests. Goto et al. [26] and Horikoshi et al. [27] reported the highest insecticidal activity of the pyripyropene derivative, Afdopyropen, against aphids. Afdopyropen causes hyperactivation followed by eventual silencing of TRPV (Transient Receptor Potential Vanilloid) channels, disrupting the function of chordotonal organs. This impairment leads to a loss of coordination, which in turn results in feeding difficulties, dehydration, and ultimately death. On a different note, Shreekanth and Reddy [28], Zala et al. [29], and Bajya et al. [30] highlighted high effectiveness of diafenthiuron in suppressing sucking pests without adverse effects on natural enemies. Conversely, Kumar et al. [31] ranked Diafenthiuron>Pyriproxyfen>Tolfenpyrad in terms of toxicity against cotton's sucking pests, while Wang et al. [32] observed Tolfenpyrad's highest toxicity against sucking pests in laboratory conditions.

## 5. CONCLUSION

The findings suggest that insecticides such as Dinotefuran 20 SG, Diafenthiuron 50 WP, and Afdopyropen 50 g/L DC, when used in combination with *L. lecanii* at half the recommended dose, exhibit equal efficacy to their solo application at the recommended dose. These combinations resulted in a cent percent mortality rate and were deemed compatible with *L. lecanii*. Utilizing these combined applications can enhance control efficacy by decreasing the quantities applied, lowering the risk of environmental pollution, and mitigating the development of pest resistance.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image

generators have been used during the writing or editing of this manuscript.

## CONSENT FOR PUBLICATION

All authors have agreed to publish this paper.

## AVAILABILITY OF DATA AND MATERIALS

All data of the study have been presented in the manuscript, and high quality and grade materials were used in this study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Kim JJ. Influence of *Lecanicillium attenuatum* on the development and reproduction of the cotton aphid, *Aphis gossypii*. *BioControl*. 2007;52:789-799. Available: <https://doi.org/10.1007/s10526-006-9050-4>
2. Razaq M, Mehmood A, Aslam M, Ismail M, Afzal M, Shad SA. Losses in yield and yield components caused by aphids to late sown Brassica napus L, Brassica juncea L. and Brassica carinata A. Braun at Multan, Punjab (Pakistan). *Pak J Bot*. 2011; 43(1):319-324.
3. Tabacian H, Ravan S, Bandani AR. Susceptibilities of two populations of *Aphis gossypii* Glover to selected insecticides. *Afr J Biotechnol*. 2011;10(4):670-674.
4. Nawaz A, Razpotnik A, Rouimi P, De Sousa G, Cravedi, JP, Rahmani R. Cellular impact of combinations of endosulfan, atrazine, and chlorpyrifos on human primary hepatocytes and HepaRG cells after short and chronic exposures. *Cell Biol Toxicol*. 2014;30:17-29. Available: <https://doi.org/10.1007/s10565-013-9266-x>
5. Van der Sluijs JP, Simon-Delso N, Goulson D, Maxim L, Bonmatin JM, Belzunces LP. Neonicotinoids, bee disorders and the sustainability of pollinator services. *Curr Opin Environ Sustain*. 2013;5(3-4):293-305. Available: <https://doi.org/10.1016/j.cosust.2013.05.007>
6. Bailey A, Grant WP, Chandler D, Greaves J, Tatchell M, Prince G. Biopesticides: Pest management and regulation. CABI; 2010.
7. Faria MR, Wraight SP. Mycoinsecticides and mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types. *J Bio Control*. 2007; 43(3):237–256.
8. Jaber LR, Enkerli J. Fungal entomopathogens as endophytes: Can they promote plant growth. *Biocontrol Sci Technol*. 2017;27(1):28–41. Available: <https://doi.org/10.1080/09583157.2016.1243227>
9. Mukherjee A, Debnath P, Ghosh SK, Medda PK. Biological control of papaya aphid (*Aphis gossypii* Glover) using entomopathogenic fungi. *Vegetos*. 2020;33:1–10. Available: <https://doi.org/10.1007/s42535-019-00072-x>
10. Halder J, Divekar PA, Rani AT. Compatibility of entomopathogenic fungi and botanicals against sucking pests of okra: an ecofriendly approach. *Egypt J Biol Pest Control*. 2021;31(1):1-7. Available: <https://doi.org/10.1186/s41938-021-00378-6>
11. Bitsadze N, Jaronski S, Khasdan V, Abashidze E, Abashidze M, Latchininsky A, Samadashvili D, Sokhadze I, Rippha M, Ishaaya I, Horowitz AR. Joint action of *Beauveria bassiana* and the insect growth regulators diflubenzuron and novaluron on the migratory locust, *Locusta migratoria*. *J Pest Sci.* 2013;86:293-300. Available: <https://doi.org/10.1007/s10340-012-0476-4>
12. Sharififard, Mona, Mossadegh MS, Vazirianzadeh B, Zarei-Mahmoudabadi A. Interactions between entomopathogenic fungus, *Metarhizium anisopliae* and sublethal doses of spinosad for control of house fly, *Musca domestica*. *Iranian J Arthropod-Borne Dis.* 2011; 5(1):28.
13. Hokkanen HM, Kotiluoto R. Bioassay of the side effects of pesticides on *Beauveria bassiana* and *Metarhizium anisopliae*: standardized sequential testing procedure. *Bulletin OILB SROP (France)*. 1992;15(3).
14. Nawaz A, Razzaq F, Razzaq A, Gogi MD, Fernández-Grandon GM, Tayib M, Ayub MA, Sufyan M, Shahid MR, Qayyum MA, Naveed M. Compatibility and synergistic

- interactions of fungi, *Metarhizium anisopliae*, and insecticide combinations against the cotton aphid, *Aphis gossypii* Glover (Hemiptera: Aphididae). *Sci Rep.* 2022;12(1):4843.  
Available:<https://doi.org/10.1038/s41598-022-08841-6>
15. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & Sons; 1984.
  16. Hassan SA. Testing methodology and the concept of the OBC/WPRS working group In Pesticides and non target invertebrates (P.C. Jepson, ed.), Intercept, Wimborne Dorset. 1989;1-8.
  17. Antonio BF, Almeida JEM, Clovis L. Effect of Tiamethoxam on entomopathogenic microorganisms. *Neotrop. Entomol.* 2001;30(3):437–447.
  18. Kumar A. et al. Amphoterization of colloidal gold particles by capping with valine molecules and their phase transfer from water to toluene by electrostatic coordination with fatty amine molecules. *Langmuir.* 2000;16(25):9775–9783.
  19. Patel DS, Jethva DM, Kchot AV. Compatibility studies of different insecticides with entomopathogenic fungus *Beauveria bassiana* (Balsamo) Vuillemin. *J Entomol Zool.* 2020;8(6):2101-2104.
  20. Reddy BN, Lakshmi VJ, Laha GS, Maheswari TU. Compatibility of entomopathogenic fungi with buprofezin for management of brown planthopper, *Nilaparvata lugens* Stal (Delphacidae: Hemiptera) in rice. *J Plant Dev Sci.* 2020; 12(1):35-38.
  21. Yadav DS, Ranade Y, Mhaske S, Ghule S. Compatibility of insecticides with *Metarhizium brunneum* (Petch) and *Beauveria bassiana* (Bals.) for bio-intensive management of pink mealybug, *Maconellicoccus hirsutus* (Green) in grapes. *J Biol Control.* 2019;33(3): 253-263.  
DOI: 10.18311/jbc/2019/22728
  22. FRAC. FRAC Code List 2018: Fungicides sorted by mode of action (including FRAC Code numbering). Fungicide Resistance Action Committee (FRAC); 2018.  
Available:<http://www.phi-base.org/images/fracCodeList.pdf>
  23. Barari H, Ghazavi M, Naseri B, Sheikhi Garjan A, Rashid M. Compatibility of the entomopathogenic fungus *Beauveria bassiana* with the insecticides fipronil, pyriproxyfen and hexaflumuron. *J Entomol Soc Iran.* 2012;31(2):29-37.
  24. Ijaz H, Muhammad S, Muhammad A, Jalal AM, Sahi ST. Efficacy of Entomopathogenic fungi alone and in combination with Buprofezin against *Sogatella furcifera* (Horváth) on rice. *Gesunde Pflanz.* 2021;73(1):85-94.  
Available:<https://doi.org/10.1007/s10343-020-00531-5>
  25. Halder J, Kushwaha D, Rai AB, Singh B. Interaction effects between entomopathogenic fungi and neonicotinoid insecticides against *Lipaphis erysimi* in vegetable ecosystem. *Indian J Agric Sci.* 2019;89(8):1256-1261.
  26. Goto K, Horikoshi R, Mitomi M, Oyama K, Hirose T, Sunazuka T, Omura S. Synthesis and insecticidal efficacy of pyripyropene derivatives. Part II—Invention of afidopyropen. *J Antibiot.* 2019;72(9):661-681.  
Available:<https://doi.org/10.1038/s41429-019-0193-9>
  27. Horikoshi R, Goto K, Mitomi M, Oyama K, Hirose T, Sunazuka T, Omura S. Afidopyropen, a novel insecticide originating from microbial secondary extracts. *Sci Rep.* 2022;12(1):28-27.  
Available:<https://doi.org/10.1038/s41598-022-06729-z>
  28. Sreekanth PN, Reddy KS. Efficacy of different insecticides against sucking pests of cotton. *J Environ Ecol.* 2011;29(4A): 2035-2039.
  29. Zala MB, Bhut GD, Patel CC, Ghetiya LV, Bharpoda TM, Borad PK. Bioefficacy of diafenthiuron 50 WP from new source against sucking insect pests in Bt cotton. *Indian J Plant Prot.* 2014;42(4):383-388.
  30. Bajya DR, Ranjith M, Lakharan MC, Raza SK. Efficacy of diafenthiuron 47.8 SC against sucking pests of cotton and its safety to natural enemies. *Indian J Entomol.* 2016;78(1):15-23.
  31. Kumar V, Grewal GK, Dhawan AK. Relative toxicity of newly introduced insecticides against sucking insect pests of transgenic Bt cotton. *Pestic Res J.* 2016;28(1):108-112.

32. Wang Y, Mu W, Zhang L, Xu Y, Lin J, Zhang Z. Toxicity assessment of insecticides to three common piercing-sucking insects in tea plantations. J Tea Sci.. 2017;37(4):392-398.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/123472>