



The Influence of Coral Reef Benthic Conditions on Associated Fish Assemblages Around Puruf Island Water, Cenderawasih Bay, Papua, Indonesia

Natalia Mansoben ^a, Roni Bawole ^{b,c*}, Jonni Marwa ^{b,d},
Christovel Alfarani Bawole ^b, Eliezer V. Sirami ^d
and Mudjirahayu ^c

^a Master of Environmental Science Program, Postgraduate, University of Papua, Manokwari, Indonesia.

^b Environmental Science Department, Postgraduate, University of Papua, Manokwari, Indonesia.

^c Faculty of Fisheries and Marine Sciences, University of Papua, Manokwari, Indonesia.

^d Faculty of Forestry, University of Papua, Manokwari, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajee/2024/v23i11618>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124976>

Original Research Article

Received: 01/10/2024
Accepted: 11/10/2024
Published: 19/10/2024

*Corresponding author: E-mail: r.bawole@unipa.ac.id;

ABSTRACT

Coral substrate in lifeforms is an environmental aspect significantly affecting fish in the ecosystem. The conditions of these coral lifeforms greatly determine fish associations and distribution. Therefore, this study aimed to explore the effect of coral reef benthic conditions on associated with fish assemblages. This research used monitoring data in 2018. Coral data was taken using transect intercept points for corals, visual census data for coral fish, which were classified into functional fish (herbivores) and key fish species for coral fisheries (carnivores). The data had been analyzed by using principal component analysis and hierarchical classification. The results showed that there was a strong interaction between fish and coral lifeforms. The interactions occurring were more varied in space and time, which affected the composition of fish community, and the spatial pattern in coral reef ecosystem. Changes in coral lifeforms and live cover had the potential to limit reef fish populations. Furthermore, the presence of herbivore and carnivore fish was proportional in coral reef ecosystem. The future research challenge is to formulate the policies needed to maintain the condition of coral reefs as a whole.

Keywords: Coral reef; benthic; reef fish; cenderawasih bay; herbivore; carnivore.

1. INTRODUCTION

Indonesia is characterized by coral reef covering 39,538 km², equivalent to 16% of the total global reef location, and is recognized as the most diverse ecosystem (Burke and Reytar, 2012). Despite the relative abundance, this ecosystem has been severely damaged by anthropogenic factors, including local stressors such as pollution, eutrophication, overfishing, destructive fishing practices, and mass bleaching associated with climate change (Razak, et al, 2022, Razak, et al 2024). Approximately a quarter of Indonesia's 270 million population lives on the coast in 30 km of coral reef, constituting the largest human population associated with coral reefs. Due to the high concentration, more than 95% of coral reef are susceptible to threats due to overfishing and destructive fishing (Burke and Reytar, 2012, Razak, et al, 2022).

Among the concentration locations in Indonesia, Cenderawasih Bay National Park (TNTC) is the largest national park, comprising 460 coral species consisting of 67 genera and subgenera, with 260 species of Scleractinia along the coast. Other species consisted of 201 mollusks, 17 mangrove vegetation, 9 coastal forest vegetation, 35 terrestrial forest vegetation, 7 seagrass, 184 birds, 14 mammals, and 17 reptiles (BB-TNTC, 2017). Coral reef ecosystem is mostly found in Cenderawasih Bay National Park, consisting of fish which are important components. However, population growth and the rising economic needs of the community have caused excessive fishing efforts (Bawole et al, 2015, Bawole et al, 2019, Bawole et al, 2018, Sala et al, 2018). This has led to the disruption of marine resources sustainability including coral reef (Bawole, 2024).

The ability of ecosystem to recover from degradation is hampered by increasing frequency, intensity, and variety of disturbances (Nyström and Folke, 2001, Buddemeier et al, 2011, Wally and Dunning, 2023). Coral reef degradation caused by increasing anthropogenic pressures (fisheries exploitation, pollution) and climate change is higher compared to natural disturbances (cyclones) (Dixon et al, 2022, Puotinen et al, 2020, Carter et al, 2022, Sully et al, 2022). This causes a decrease in coral cover and structural complexity (López-Victoria and Zapata, 2018, Bruno and Selig, 2007), associated changes in coral and fish community composition (López-Victoria and Zapata, 2018, Emslie et al, 2011) as well as shifts in the dominant benthic biota (Dudgeon et al, 2010, Bruno et al, 2009, Mumby, 2009, Norström et al, 2009, Kimura and Chou, 2018).

The shifts recorded on coral reef include changes in coralline, sponge, or most commonly benthic macroalgae dominance (Mumby, 2009, Norström et al, 2009). However, other benthic lifeforms are a typical component of most scleractinian coral. This is due to their primary function as providers of complex structural habitats and responsible for the high diversity of reef-associated organisms, providing various important ecosystem services such as food resources (Cabral and Geronimo, 2018, Littler et al, 2009, Cruz-Trinidad et al, 2014).

All major coral reef locations globally have experienced a decrease in coral cover (Bruno and Selig, 2007, Chanda, 2022, Ateweberhan et al, 2011). In this context, reduced coral cover is indicative of an increasing shift in dominant

benthic biota (Norström et al, 2009), and changes in benthic community composition (Bawole et al, 2014). The shift from corals to macroalgae can lead to excessive herbivore fish populations and stimulate fisheries. However, the impact of benthic community on coral lifeforms, including the entire ecosystem, and assemblage of reef fish (herbivore and carnivore) is not fully understood.

The complex interconnections between organisms and their benthic substrates (coral lifeforms) imply that changes in one aspect of ecosystem can lead to a cascade of subsequent which are often unanticipated (Möllmann et al, 2011, Van den Elsen, 2020, Connell et al, 2011). A previous study has shown a strong relationship between coral reef fish and benthic habitats (Sym and Jones, 2000, 2004). There is variability in the specific responses of different fish, causing significant changes in coral cover (McCormick et al, 2017, Jones et al, 2006, Wilson et al, 2006). For example, the loss of live coral can cause shifts in entire fish assemblages (McCormick et al, 2017, Bellwood et al, 2012, Cheal et al, 2010, Bellwood et al, 2006), as well as a rapid decrease in fish abundance and diversity (Dulvy

et al, 2006, Williamson et al, 2014, Jones et al, 2004). Although the potential of benthic habitats to provide the environments required for reef fish had not been extensively explored, (Syms and Jones, 2001) showed that soft coral was unfavorable a substitute for hard types. From ecosystem perspective, some fish species can associate with the same or different habitats based on community by providing the environments required for associated reef fish assemblages.

Based on the description, this study aimed to explore the impact of coral reef benthic conditions on associated with fish assemblages.

2. MATERIALS AND METHODS

2.1 Location and Time

This study was conducted from July to September 2024, at Rumberpon and Wamesa Districts, Wondama Regency, as shown in Fig. 1. The location was part of Cenderawasih Bay National Park (BB-TNTC, 2017), which consists of three locations (Puruf, Nuana and Tapapai Island). Puruf Island is larger than the other two islands, and is an island for tourist destinations.




Fig. 1. Study location map

2.2 Data Collection Method

The data used were obtained from Reef Health Monitoring database conducted by Bird Head Seascape (BHS) Papua Consortium, stored at the Research and Community Service Institute, University of Papua, Manowari, Indonesia. The data used is observation data in 2018. Coral data was collected using Point Intercept Transect (PIT) method (Wilson and Green, 2009, Ahmadi et al, 2013) to determine the basic biota constituting coral reef ecosystem. Observations were made using lifeforms category or growth providing a morphological description of coral reef ecosystem (English et al, 1997). PIT method only monitored coral lifeforms by collecting reef data per point along the transect.

2.3 Data Analysis

The total percentage of coral cover and benthic growth forms are presented in graphs and tables using Excel. The calculation of the percentage of cover for each component was carried out using the formula expressed in studies (Manuputty Djuwariah, 2009). Subsequently, the condition of coral reef cover was divided into five categories, according to a previous study (English et al, 1997), comprising very poor (0 - 10%), poor (11 - 30%), moderate (31 - 50%), good (51 - 75%), and very good (76 - 100%). Analysis of location grouping based on coral lifeforms types was carried out using cluster analysis.

Fish species abundance and estimated length were assessed using bottom visual sensors (Ahmadi et al, 2013). Based on the results, species recorded were the families/subfamilies Acanthuridae, Carangidae, Carcharinidae, Dasyatidae, Haemulidae, Lethrinidae, Lutjanidae, Scaridae, Scrombidae, Serranidae, Siganidae, and *Cheilinus undulatus*. The total fish biomass recorded included high-value fish species (families Serranidae, Lutjanidae, Lethrinidae, and Haemulidae). These species were classified as carnivore, and the biomass of herbivore fish (families/subfamilies Acanthuridae, Scaridae, and Siganidae) was calculated to determine the status of coral reef.

3. RESULTS AND DISCUSSION

3.1 Habitat Characteristics Based on Coral Lifeforms

Habitat characteristics search of study locations based on coral lifeforms types showed significant grouping, as presented in Fig. 2. Group 1 was characterized by *Acropora* branching, coral

massive, coral mushroom, dead coral, sponge, coral branching, coral foliose, and sand. Meanwhile, group II had coral lifeforms *Acropora* encrusting, coral submassive, coralline algae, turf algae, and soft coral, as well as coral encrusting. Group III consisted of macroalgae, rock, OT, and Ruble. Regarding the proportion of coral lifeforms, further observation showed that group III had similar characteristics to Nuana Island location. The characteristics of group II were identical to Tapapai location and group I had Puruf Island group habitat (Fig. 2). In general, the research location was characterized by the type of coral lifeform from massive, foliose, branching and soft corals. Coral fragments (rubble) occupies a high portion as an abiotic factor (Fig. 3).

The influence of coral habitat characteristics in lifeforms on the interaction and distribution patterns of fish (herbivore and carnivore) showed a strong interaction (French et al, 1989). The interaction varied in space and time due to the significant influence on composition of fish community and the spatial pattern of target fish in coral reef ecosystem. Changes in coral lifeforms and live cover had the potential to limit reef fish populations (Kane and Tissot, 2017, Muller-Karanassos et al, 2021, Shi et al, 2022, Rummer and Illing, 2022) including Indonesia (Bawole et al, 2014, Runtuboi et al, 2019). The condition of coral reef cover was divided into five categories, according to English et al. (1994). These consisted of very poor (0-10%), poor (11-30%), moderate (31-50%), good (51-75%), and very good (76-100%). Analysis of location grouping based on coral lifeforms was carried out using cluster analysis. The results showed that coral reef condition of Puruf, Tapapai, and Nuana Island belonged to the good, poor, and moderate categories, respectively (Table 1).

The results showed that Puruf Island had a higher percentage of live coral cover compared to the study by (Sala et al, 2021), although Cenderwasih Bay obtained values of 47% and 44%, respectively. This study was approximately similar to the Tapapai coral cover which was categorized as moderate. The results indicated that the quality of coral reef was in poor and good conditions, but could be considered good. Some of coral was damaged (15%) due to pressure from human activities using environmentally unfriendly fishing gear and natural phenomena such as earthquakes and waves.




Fig. 2. Grouping of coral lifeforms characteristics based on study location




Fig. 3. Proportion of coral lifeforms at the study location

Table 1. Coral conditions on selected sites

Location Name	Hard coral (%)	Categories according to English et al. (1994)
Tapapai Island	48	Moderate
Nuana Island	27	Poor
Puruf Island	67.33	Good

3.2 Associated Fish and Coral Lifeforms

The analysis results showed that there was association between coral lifeforms as well as the presence of herbivore and carnivore fish, as presented in Fig. 4. Generally, carnivore is more associated with coral encrusting, branching, and rock, while herbivore is related to soft coral, macroalgae, and turf algae. This association supported previous analysis conducted in Cenderawasih Bay location (Bawole et al, 2014), where coral reef conditions had a close relationship with fish (Indrawati et al, 2020, Suwarni and Nurlina, 2022, Adrian et al, 2020, Erdana et al, 2022). In addition to the cover of live coral, the presence of reef fish was also related to coral lifeforms (Kamarumtham et al, 2016, Wanma et al, 2022, Harsindhi et al, 2020). Target fish groups often experience a decrease in population after major disturbances (Magel et al, 2020, Walpole and Hadwen, 2022). Moreover, several studies have shown the importance of coral habitat in supporting reef fish groups and promoting their distribution (Bawole et al, 2014, Shi et al, 2022).

The presence of herbivore fish in coral reef ecosystem plays a significant role as a cleaner of algae that sticks to living and dead coral. By eating algae and turf, herbivore fish regulate the

stability of coral ecosystem (Wilson and Green, 2009), ranking second at the tropical level as first-level consumers. Meanwhile, carnivore fish describe the condition of coral reef ecosystem. This group of fish is the target of fishing due to the high selling price, leading to elevated intensity of fishing activities.

The analysis results showed that the biomass of herbivore fish at the three observation locations was very different, as shown in Table 2. Herbivore fish at Purup Island location were lower compared to others (Tapapai and Nuana Island). This was related to the level of fishing at the existing location, where the intensity was greater at Purup Island and two other locations. The identified groups were from Achanturidae (sand skin), Scaridae (old parrotfish), Siganidae (baronang), and Labridae (napoleon/maming) families. This type of fish was often caught by local fishermen, particularly from Yomakan community.

Similar to herbivore group, carnivore fish shows the same biomass, which is lower at Purup Island compared to others (Tapapai and Nuana Island). Carnivore group consists of fish from Lutjanidae (snapper), Letrinidae (lencam), and Seranidae (grouper) families, serving as target of local fishermen.




Fig. 4. Grouping of associated herbivore and carnivore fish with coral lifeform types.

Table 2. Herbivore and carnivore reef fish of selected sites

Location Name	Herbivore Fish (kg/ha)	Carnivore Fish (kg/ha)
Tapapai	316.86	38.24
Nuana Island	128.21	90.67
Purup Island	64.84	13.18

Based on the existing biomass, both herbivore and carnivore groups, describe their presence proportionally in the three study locations. This shows that the predominance presence of herbivore group a location needs to be followed by the corresponding amount of carnivore and vice versa. Therefore, the condition of coral reef ecosystem and the oceanographic aspects of water will support fish stability. This finding was not too surprising because the same research results were also found by other researchers (Bawole et al, 2024).

Although habitat preferences may initially determine the spatial distribution of many reef fish, the patterns that are formed can be altered by several ecological processes, including mortality, competition, and fish behavior (Robert, 2000, Huijbers et al, 2011, Nagelkerken et al, 2000, Fulton et al, 2017). This is particularly relevant for target fish due to the lack of further movement after being established. Many fish species have very small home ranges, along with high fidelity to specific fishing sites (Welsh et al, 2013, Farmer and Ault, 2014, Moffitt et al, 2009). Moreover, the regulation of target fish in coral reef ecosystem is strongly influenced by trophic level structure, feeding patterns, the degree of habitat disturbance, and fishing activity (Kane and Tissot, 2017, Muller-Karanassos et al, 2021, Graham et al, 2007, Kittinger and Kittinger, 2013).

4. CONCLUSION

The association of fish and coral is largely determined by the characteristics of the benthic substrate (lifefom). This character is important in influencing the spatial distribution of coral fish based on coral lifeform. Herbivorous and carnivorous fish are proportional in number and type based on the characteristics of the coral lifeform.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

Informed consent was obtained from all participants prior to data collection, ensuring that they were aware of the purpose of the study and their rights as participants. Confidentiality was

maintained throughout the study, and participants were assured that their responses would be anonymized in any reports or publications

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Adrian, D., Kurniawan, D., & Putra, R. D. (2020). Hubungan persentaseutupan karang hidup dengan kelimpahan ikan indikator Chaetodontidae di perairan Pengudang, Kabupaten Bintan. *Jurnal Akuatiklestari*, 3(2), 53–59.
- Ahmadia, G. N., Wilson, J. R., & Green, A. L. (2013). Coral reef monitoring protocol for assessing marine protected areas in the Coral Triangle. *Coral Triangle Support Partnership*.
- Ateweberhan, M., McClanahan, T. R., Graham, N. A. J., & Sheppard, C. R. C. (2011). Episodic heterogeneous decline and recovery of coral cover in the Indian Ocean. *Coral Reefs*, 30(3), 729–741. <https://doi.org/10.1007/s00338-011-0747-z>
- Bawole, C. A., Kaber, Y., Sirami, E. V., Coloay, C. G., Schaduw, J. N. W., & Mudjirahayu. (2024). Spatial variation of reef fish based on coral lifeform on Lihaga Island, Likupang, North Sulawesi. *International Journal of Recent Advances in Multidisciplinary Research*, 11(01), 9539–944.
- Bawole, R. (2024). Permasalahan dan ancaman wilayah pesisir dan laut. In A. Yanto (Ed.), *Pengantar ilmu perikanan dan perairan* (1st ed., pp. 229–262). Padang: Get Press Indonesia.
- Bawole, R., Pattiasina, T. F., & Kawulur, E. I. J. J. (2014). Coral-fish association and its spatial distribution in Cenderawasih Bay National Park, Papua, Indonesia. *AACL Bioflux*, 7(4), 233–240.
- Bawole, R., Yulianda, F., Bengen, D. G., Fahrudin, A., & Mudjirahayu, M. (2015). Socio-ecological system within governance of marine protected area: Case from Cenderawasih Bay National Park, Indonesia. *Jurnal Manajemen Hutan Tropika (Journal of Tropical Forest Management)*, 21(1), 19–24. <http://journal.ipb.ac.id/index.php/jmht/article/view/9493>

- Bawole, R., Yulianda, F., Bengen, D. G., Fahrudin, A., & Mudjirahayu. (2019). Governability of marine protected areas: The effect of trade-offs, hybrid patterns, and policy implications in Cenderawasih Bay National Park, Indonesia. *IOP Conference Series: Earth and Environmental Science*, 278(3), 1–14. <https://doi.org/10.1088/1755-1315/278/3/032029>
- Bawole, R., Yulianda, F., Bengen, D. G., Fahrudin, A., Mudjirahayu, Rembet, U. N. (2018). *Tata kelola kawasan konservasi perairan: Sistem tata kelola, sistem yang dikelola, dan tata kelola interaktif*. Bandung: CV. Alfabeta Bandung.
- BB-TNTC. (2017). *Rencana pengelolaan jangka panjang taman nasional teluk cenderawasih provinsi papua dan papua barat periode 2018 – 2027*. Manokwari.
- Bellwood, D. R., Baird, A. H., Depczynski, M., González-Cabello, A., Hoey, A. S., Lefèvre, C. D., et al. (2012). Coral recovery may not herald the return of fishes on damaged coral reefs. *Oecologia*, 170(2), 501–503. <https://doi.org/10.1007/s00442-012-2290-3>
- Bellwood, D. R., Hoey, A. S., Ackerman, J. L., & Depczynski, M. (2006). Coral bleaching, reef fish community phase shifts and the resilience of coral reefs. *Global Change Biology*, 12(9), 1587–1594. <https://doi.org/10.1111/j.1365-2486.2006.01206.x>
- Bruno, J. F., & Selig, E. R. (2007). Regional decline of coral cover in the Indo-Pacific: Timing, extent, and subregional comparisons. *PLoS ONE*, 2(8), e711. <https://doi.org/10.1371/journal.pone.0000711>
- Bruno, J. F., Sweatman, H., Precht, W. F., Selig, E. R., & Schutte, V. G. W. (2009). Assessing evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology*, 90(6), 1478–1484. <https://doi.org/10.1890/08-0457.1>
- Buddemeier, R. W., Kleypas, J. A., & Aronson, R. B. (2011). *Coral reef and global climate change*. Prepared for the Pew Center on Global Climate Change.
- Burke, L., Reytar, K., Spalding, M., & Perry, A. (2012). *Reefs at risk revisited in the Coral Triangle*. World Resources Institute.
- Cabral, R. B., & Geronimo, R. C. (2018). How important are coral reefs to food security in the Philippines? Diving deeper than national aggregates and averages. *Marine Policy*, 91, 227–234. <https://doi.org/10.1016/j.marpol.2018.02.002>
- Carter, A. L., Gilchrist, H., Dexter, K. G., Gardner, C. J., Gough, C., Roccliffe, S., et al. (2022). Cyclone impacts on coral reef communities in southwest Madagascar. *Frontiers in Marine Science*, 9, 860675. <https://doi.org/10.3389/fmars.2022.860675>
- Chanda, A. (2022). An overview of the Indian Ocean coral ecosystems. In *Blue carbon dynamics of the Indian Ocean: The present state of the art*.
- Cheal, A. J., MacNeil, M. A., Cripps, E., Emslie, M. J., Jonker, M., Schaffelke, B., et al. (2010). Coral-macroalgal phase shifts or reef resilience: Links with diversity and functional roles of herbivorous fishes on the Great Barrier Reef. *Coral Reefs*, 29(4), 859–868. <https://doi.org/10.1007/s00338-010-0645-5>
- Connell, S. D., Russell, B. D., & Irving, A. D. (2011). Can strong consumer and producer effects be reconciled to better forecast “catastrophic” phase shifts in marine ecosystems? *Journal of Experimental Marine Biology and Ecology*, 400(1), 1–9. <https://doi.org/10.1016/j.jembe.2010.12.016>
- Cruz-Trinidad, A., Aliño, P. M., Geronimo, R. C., & Cabral, R. B. (2014). Linking food security with coral reefs and fisheries in the Coral Triangle. *Coastal Management*, 42(2), 160–182. <https://doi.org/10.1080/08920753.2014.886840>
- Dixon, A. M., Puotinen, M., Ramsay, H. A., & Beger, M. (2022). Coral reef exposure to damaging tropical cyclone waves in a warming climate. *Earth's Future*, 10(8), e2022EF002775. <https://doi.org/10.1029/2022EF002775>
- Dudgeon, S. R., Aronson, R. B., Bruno, J. F., & Precht, W. F. (2010). Phase shifts and stable states on coral reefs. *Marine Ecology Progress Series*, 413, 1–12. <https://doi.org/10.3354/meps08764>
- Dulvy, N. K., Jennings, S., Rogers, S. I., & Maxwell, D. L. (2006). Threat and decline in fishes: An indicator of marine biodiversity. *Canadian Journal of Fisheries and Aquatic Sciences*, 63(6), 1267–1275. <https://doi.org/10.1139/f06-007>
- Emslie, M. J., Pratchett, M. S., & Cheal, A. J. (2011). Effects of different disturbance types on butterflyfish communities of Australia’s Great Barrier Reef. *Coral*

- Reefs, 30(2), 351–363.
<https://doi.org/10.1007/s00338-011-0724-6>
- English, S., Wilkinson, C., & Baker, V. (1997). *Survey manual for tropical marine resources*.
- Erdana, R., Pratikto, I., & Suryono, C. A. (2022). Hubungan persentase tutupan karang hidup dan kelimpahan ikan di kawasan konservasi perairan Pulau Koon, Kabupaten Seram bagian timur, Provinsi Maluku. *J Mar Res*, 11(2), 87–93.
- Farmer, N. A., & Ault, J. S. (2014). Modeling coral reef fish home range movements in dry Tortugas, Florida. *The Scientific World Journal*, 2014, 1–13.
<https://doi.org/10.1155/2014/275185>
- French, D. D., Ludwig, J. A., & Reynolds, J. F. (1989). Statistical ecology: A primer on methods and computing. *Journal of Applied Ecology*, 26(3), 487–494.
<https://doi.org/10.2307/2403427>
- Fulton, C. J., Wainwright, P. C., Hoey, A. S., & Bellwood, D. R. (2017). Global ecological success of *Thalassoma* fishes in extreme coral reef habitats. *Ecology and Evolution*, 7(1), 194–200.
<https://doi.org/10.1002/ece3.2575>
- Graham, N. A. J., Wilson, S. K., Jennings, S., Polunin, N. V. C., Robinson, J., Bijoux, J. P., et al. (2007). Lag effects in the impacts of mass coral bleaching on coral reef fish, fisheries and ecosystems. *Conservation Biology*, 21(5), 1291–1300.
<https://doi.org/10.1111/j.1523-1739.2007.00751.x>
- Harsindhi, C. J., Bengen, D. G., Zamani, N. P., & Kurniawan, F. (2020). Abundance and spatial distribution of reef fish based on coral lifeforms at Tidung Island, Seribu Islands, Jakarta Bay. *AAFL Bioflux*, 13(2), 637–644.
- Huijbers, C. M., Nagelkerken, I., Govers, L. L., Van De Kerk, M., Oldenburger, J. J., & De Brouwer, J. H. F. (2011). Habitat type and schooling interactively determine refuge-seeking behavior in a coral reef fish throughout ontogeny. *Marine Ecology Progress Series*, 437(2), 241–251.
<https://doi.org/10.3354/meps09286>
- Indrawati, A., Edrus, I. N., & Hadi, T. A. (2020). Karakteristik struktur komunitas ikan karang target dan indikator di perairan taman nasional Komodo. *Jurnal Penelitian Perikanan Indonesia*, 26(2), 153–162.
- Jones, G. P., McCormick, M. I., Srinivasan, M., & Eagle, J. V. (2004). Coral decline threatens fish biodiversity in marine reserves. *Proceedings of the National Academy of Sciences of the United States of America*, 101(21), 8251–8253.
<https://doi.org/10.1073/pnas.0401689101>
- Jones, G. P., Polunin, N. V. C., Graham, N. A. J., Wilson, S. K., & Pratchett, M. S. (2006). Multiple disturbances and the global degradation of coral reefs: Are reef fishes at risk or resilient? *Global Change Biology*, 12(11), 1294–1305.
<https://doi.org/10.1111/j.1365-2486.2006.01213.x>
- Kamarumtham, K., Ahmad, Z., Hidayah Halid, N., Saad, S., Fikri Akmal Khodzori, M., Hamizan Yusof, M., et al. (2016). Diversity and distribution of coral lifeforms in Tioman Island. *Transactions on Science and Technology*, 3(3), 207–214.
- Kane, C. N., & Tissot, B. N. (2017). Trophic designation and live coral cover predict changes in reef-fish community structure along a shallow to mesophotic gradient in Hawaii. *Coral Reefs*, 36(3), 999–1012.
<https://doi.org/10.1007/s00338-017-1604-8>
- Kimura, T. T., & Chou, L. M. (2018). Status of coral reefs in East Asian Seas Region: 2018. *Status of Coral Reefs in East Asian Seas Region*.
- Kittinger, J. N. (2013). Participatory fishing community assessments to support coral reef fisheries comanagement. *Ocean & Coastal Management*, 67, 361–381.
<https://doi.org/10.1016/j.ocecoaman.2012.12.012>
- Littler, M. M., Littler, D. S., & Brooks, B. L. (2009). Herbivory, nutrients, stochastic events, and relative dominances of benthic indicator groups on coral reefs: A review and recommendations. *Smithsonian Contributions to Marine Science*, 38, 1–32.
- López-Victoria, M., & Zapata, F. A. (2018). Coral cover decline on La Chola reef (Eastern Tropical Pacific). *Boletín de Investigaciones Marinas y Costeras*, 47, 1–10.
- Magel, J. M. T., Dimoff, S. A., & Baum, J. K. (2020). Direct and indirect effects of climate change-amplified pulse heat stress events on coral reef fish communities. *Ecological Applications*, 30(6), e02054.
<https://doi.org/10.1002/eap.2054>
- Manuputty Djuwariah, A. E. W. (2009). *Method guide: Point Intercept transect (PIT) for community, baseline study and coral health monitoring at marine no take zone area (DPL)*. Coral Reef Rehabilitation and

- Management Program, Indonesian Institute of Sciences Jakarta, Indonesia.
- McCormick, M. I., Barry, R. P., & Allan, B. J. M. (2017). Algae associated with coral degradation affects risk assessment in coral reef fishes. *Scientific Reports*, 7, 2208. <https://doi.org/10.1038/s41598-017-02405-7>
- Moffitt, E. A., Botsford, L. W., Kaplan, D. M., & O'Farrell, M. R. (2009). Marine reserve networks for species that move within a home range. *Ecology Applications*, 19(7), 1835–1847. <https://doi.org/10.1890/08-0545.1>
- Möllmann, C., Casini, M., Daskalov, G. M., & De Young, B. (2011). Regime shifts in marine ecosystems: How overfishing can provoke sudden ecosystem changes. *Directorate General for Internal Policies Policy Department B: Structural and Cohesion Policies Fisheries European Parliament*.
- Muller-Karanassos, C., Filous, A., Friedlander, A. M., Cuetos-Bueno, J., Gouezo, M., Lindfield, S. J., et al. (2021). Effects of habitat, fishing, and fisheries management on reef fish populations in Palau. *Fisheries Research*, 241, 106048. <https://doi.org/10.1016/j.fishres.2021.106048>
- Mumby, P. J. (2009). Phase shifts and the stability of macroalgal communities on Caribbean coral reefs. *Coral Reefs*, 28(3), 581–589. <https://doi.org/10.1007/s00338-009-0491-5>
- Nagelkerken, I., Dorenbosch, M., Verberk, W. C. E. P., Cocheret de la Moriniere, E., & Van der Velde, G. (2000). Importance of shallow-water biotopes of a Caribbean bay for juvenile coral reef fishes: Patterns in biotope association, community structure and spatial distribution. *Marine Ecology Progress Series*, 202, 175–192. <https://doi.org/10.3354/meps202175>
- Norström, A. V., Nyström, M., Lokrantz, J., & Folke, C. (2009). Alternative states on coral reefs: Beyond coral-macroalgal phase shifts. *Marine Ecology Progress Series*, 376, 295–306. <https://doi.org/10.3354/meps07865>
- Nyström, M., & Folke, C. (2001). Spatial resilience of coral reefs. *Ecosystems*, 4, 406–417. <https://doi.org/10.1007/s10021-001-0002-7>
- Puotinen, M., Drost, E., Lowe, R., Depczynski, M., Radford, B., Heyward, A., et al. (2020). Towards modelling the future risk of cyclone wave damage to the world's coral reefs. *Global Change Biology*, 26(8), 4532–4546. <https://doi.org/10.1111/gcb.15273>
- Razak, T. B., Boström-Einarsson, L., Alisa, C. A. G., Vida, R. T., & Lamont, T. A. C. (2022). Coral reef restoration in Indonesia: A review of policies and projects. *Marine Policy*, 137, 104979. <https://doi.org/10.1016/j.marpol.2021.104979>
- Razak, T. B., Lamont, T. A. C., Hukom, F. D., Alisa, C. A. G., Asri, A. R., & Ferse, S. C. A. (2024). A review of the legal framework for coral reef restoration in Indonesia. *Ocean & Coastal Management*, 248, 105683. <https://doi.org/10.1016/j.ocecoaman.2023.105683>
- Robert, S. D. (2000). Grouper and napoleon wrasse ecology in Laamu Atoll, Republic of Maldives: Part 1. Habitat, Behavior, and Movement Patterns. *AACL Bioflux*.
- Rummer, J. L., & Illing, B. (2022). Coral reef fishes in a multi-stressor world. In *Fish Physiology*.
- Runtuboi, F., Bawole, R., Goram, A., Wawiyai, Y., Wambraw, M., Numberi, Y. Z., et al. (2019). Inventarisasi jenis ikan karang dan komposisi jenis ikan ekonomis penting (study kasus Kampung Kornasoren, Saribi dan Syoribo) pulau Numfor Kabupaten Biak Numfor. *Journal of Tropical Fisheries Management*, 2(1), 1–12.
- Sala, R., Bawole, R., Biloru, R. H. H., & Mudjirahayu, M. (2021). Distribusi spasial tutupan karang di taman nasional teluk cenderawasih, Papua. *Jurnal Sumberdaya Akuatik Indopasifik*, 5(2), 201–212.
- Sala, R., Bawole, R., Runtuboi, F., Mudjirahayu, M., Wopi, I. A., Budisetiawan, J., et al. (2018). Population dynamics of the yellowstripe scad (*Selaroides leptolepis* Cuvier, 1833) and Indian mackerel (*Rastrelliger kanagurta* Cuvier, 1816) in Wondama Bay Water, Indonesia. In *IOP Conference Series: Earth and Environmental Science*.
- Shi, J., Li, C., Wang, T., Zhao, J., Liu, Y., & Xiao, Y. (2022). Distribution pattern of coral reef fishes in China. *Sustainability (Switzerland)*, 14(22), 15135. <https://doi.org/10.3390/su142215135>
- Sully, S., Hodgson, G., & van Woeseik, R. (2022). Present and future bright and dark spots for coral reefs through climate change. *Global Change Biology*, 28(15), 4710–4722. <https://doi.org/10.1111/gcb.16482>

- Suwarni, S., & Nurlina, N. (2022). Kelimpahan ikan famili Scaridae berdasarkan tutupan terumbu karang hidup di perairan Pulau Kapoposang Kabupaten Pangkajene dan Kepulauan. *Prosiding Seminar Nasional Politeknik Pertanian Negeri Pangkajene Kepulauan*, 3, 68–76.
- Syms, C., & Jones, G. P. (2000). Disturbance, habitat structure, and the dynamics of a coral-reef fish community. *Ecology*, 81(10), 2710–2729. [https://doi.org/10.1890/0012-9658\(2000\)081\[2710:DHSTAD\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[2710:DHSTAD]2.0.CO;2)
- Syms, C., & Jones, G. P. (2001). Soft corals exert no direct effects on coral reef fish assemblages. *Oecologia*, 127(4), 489–495. <https://doi.org/10.1007/s004420100637>
- Syms, C., & Jones, G. P. (2004). Habitat structure, disturbance and the composition of sand-dwelling goby assemblages in a coral reef lagoon. *Marine Ecology Progress Series*, 268, 271–279. <https://doi.org/10.3354/meps268271>
- van den Elsen, E., Stringer, L. C., De Ita, C., Hessel, R., Kéfi, S., Schneider, F. D., et al. (2020). Advances in understanding and managing catastrophic ecosystem shifts in Mediterranean ecosystems. *Frontiers in Ecology and Evolution*, 8, 1–14. <https://doi.org/10.3389/fevo.2020.00093>
- Wally, K., & Dunning, K. (2023). Climate change, coral bleaching and other threats. In *Democratic management of an ecosystem under threat: The People's Reefs*.
- Walpole, L. C., & Hadwen, W. L. (2022). Extreme events, loss, and grief—an evaluation of the evolving management of climate change threats on the Great Barrier Reef. *Ecology and Society*, 27(1). <https://doi.org/10.5751/ES-12837-270101>
- Wanma, M., Manan, J., Loinenak, F. A., & Kolibongso, D. (2022). Variations and condition of coral lifeforms in the coastal area of Rendani Airport, Manokwari, Indonesia. *Jurnal Sumberdaya Akuatik Indopasifik*, 6(2), 113–120.
- Welsh, J. Q., Goatley, C. H. R., & Bellwood, D. R. (2013). The ontogeny of home ranges: Evidence from coral reef fishes. *Proceedings of the Royal Society B: Biological Sciences*, 280(1773), 20122730. <https://doi.org/10.1098/rspb.2012.2730>
- Williamson, D. H., Ceccarelli, D. M., Evans, R. D., Jones, G. P., & Russ, G. R. (2014). Habitat dynamics, marine reserve status, and the decline and recovery of coral reef fish communities. *Ecology and Evolution*, 4(4), 576–588. <https://doi.org/10.1002/ece3.911>
- Wilson, J., & Green, A. (2009). Biological monitoring methods for assessing coral reef health and management effectiveness of marine protected areas in Indonesia. *The Nature Conservancy*.
- Wilson, S. K., Graham, N. A. J., Pratchett, M. S., Jones, G. P., & Polunin, N. V. C. (2006). Multiple disturbances and the global degradation of coral reefs: Are reef fishes at risk or resilient? *Global Change Biology*, 12, 1294–1305. <https://doi.org/10.1111/j.1365-2486.2006.01213.x>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/124976>